




LENDLEASE COMMUNITIES

SPRINGFIELD RISE - VILLAGE 11 (NORTH)

SITE BASED MANAGEMENT PLAN

01 CONTENTS

1	CONTENTS	2
2	INTRODUCTION	3
3	SITE DESCRIPTION	4
4	ECOLOGICAL VALUES - SUMMARY	5
5	ENVIRONMENTAL MANAGEMENT	6
6	PRE-CLEARANCE VEGETATION MANAGEMENT	7
	PRE-CLEARANCE VEGETATION MANAGEMENT	8
	PRE-CLEARANCE VEGETATION MANAGEMENT	9
7	PRE-CLEARANCE FAUNA MANAGEMENT	10
	PRE-CLEARANCE FAUNA MANAGEMENT	11
	PRE-CLEARANCE FAUNA MANAGEMENT	12
	PRE-CLEARANCE FAUNA MANAGEMENT	13
	PRE-CLEARANCE FAUNA MANAGEMENT	14
8	FAUNA MANAGEMENT CONSTRUCTION	15
9	THREATENED FLORA MANAGEMENT	16
	THREATENED FLORA MANAGEMENT	17
	THREATENED FLORA MANAGEMENT	18
10	FLORA AND FAUNA CHECKLIST	19



02 INTRODUCTION

Introduction

This phase specific Site Based Management Plan (SBMP) has been prepared for the north extent of Village 11 (V11) of Springfield Rise at Spring Mountain Estate and incorporates the management intent, objectives and specifications detailed within the overarching environmental management plans prepared for the development.

The aim of this SBMP-V11 is to set out and guide the implementation of effective measures to ameliorate any impacts, and to ensure and manage the long term sustainability of the project and its natural environment, specifically for Matters of National Environmental Significance (MNES) listed species known to occur within the Spring Mountain project site namely:

- Phascolarctos cinereus (Koala)
- Pteropus poliocephalus (Grey-headed Flying-fox)
- Plectranthus habrophyllus

The document has been developed in accordance with the Spring Mountain SBMP, prepared by Yurrah, as an updated and re-issued phase specific management plan.

The purpose of this SBMP-V11 is to provide a single, consolidated management document which incorporates requirements of numerous ecological management plans prepared for Spring Mountain. From these documents, this SBMP-V11 extracts management objectives, implementation requirements, performance indicators and monitoring and auditing actions relevant to the specific the development of V11 for both construction and operational phases.

Environmental Pre-Start Checklist

This Site Based Management Plan has been prepared to create an on-site working document with easy to find references to management measures without the comprehensive details of the assessment and approval. Core to contractors working under this SBMP is completion of the Spring Mountain Pre-Start Environmental Checklist. Completion and sign off of this checklist, inclusive of attachments should warrant compliance with this SBMP and broader approval parameters.

Details on this SBMP can be found within the following documents:

- Site Based Management Plan for Spring Mountain Community, prepared by **Yurrah** (July 2015)
- Threatened Flora Management Plan for Spring Mountain, prepared by **Yurrah** (July 2015)
- Fauna Management Plan for Spring Mountain, prepared by **Saunders Havill Group** (July 2015)
- Code of practice for Welfare of Animals effected by Land Clearing and Other Habitat Impacts, and Wildlife/ Spotter Catchers (Draft) prepared by **Wildlife Warriors and Voiceless** (2009)
- Offsets Management Plan prepared for Spring Mountain, prepared by **Saunders Havill Group** (July 2015)
- Bushfire Management Plan for Spring Mountain, prepared by **Cardno** (2016)

This SBMP-V11 should be read in conjunction with all V11 approvals and conditions including approved civil, landscape, vegetation management and rehabilitation plans and specifications.

This SBMP-V11 has also been prepared to meet compliance and auditing requirements of the Spring Mountain Commonwealth Department of the Environment and Energy (DEE) approval under the Environment Protection and Biodiversity Conservation Act 1999 (EPBC) (Ref: 2013/7057), specifically Conditions 3-6.

This SBMP-V11 outlines construction measures specific to V11 to manage of impacts to native flora and fauna.

Construction

- Vegetation Management (Clearing & Protection)
- Protection of MNES Fauna (Koala and Grey-headed Flying Fox) and Native Wildlife
- Maintenance of Safe Wildlife Movement Opportunities
- Fauna Habitat Rehabilitation
- Threatened Flora Management
- Pest Management
- Fire Management
- Education and Awareness



03 SITE DESCRIPTION

Site Description

V11 is located centrally within the precinct comprising areas to the north and south of the continuation of Grande Avenue, and maintains a southern boundary adjoining conservation land and western boundary defined by Villages 14 and 15. The village may accommodate some of the proposed Neighbourhood Centre within the northern part of the village. The portion north of Grande Avenue is defined by the confluence of linear open space, the Mountain Creek corridor and the provision of District Recreation Park. Village 11 has a development area of approximately 28ha.

Natural Features

V11 comprises land with undulating topography with subtle ridges emanating from the peak of RL136 in the south western corner of the village exhibiting views over Spring Mountain and Springfield Central. The western-most ridge including a saddle formation typically defines the western boundary with Village 14 adjacent while a minor gully form defines the eastern boundary with Village 10. This boundary will involve earthworks to achieve a consistent interface with adjoining Village 10.

Land Use

V11 will be developed for typical residential development predominately comprising of a range of low rise (1-2 storey) detached dwelling forms. The potential for higher densities and attached dwelling forms towards the northern portions of the village proximate to the Neighbourhood Centre will achieve a suitable integration with this centre.

Interface with Linear Open Space and District Recreation Park (formal Park and Gardens)

The interface with District Recreation Park will ensure a high level of visual permeability to highlight the presentation of the park and non-vehicular access to other parts of the District Recreation Park. Opposite Mountain Creek. Additionally, the provision of access may provide opportunity for a carparking location adjacent to the District Recreation Park.




Photo: Grey-headed Flying-fox (listed as Vulnerable under EPBC Act (Cth))



Photo: Plectranthus habrophyllus (listed as Endangered under EPBC Act (Cth))



Photo: Koala (listed as Vulnerable under EPBC Act (Cth) and NCA (Qld))



Extract: V11 development site from the Spring Mountain Precinct Plan

04 ECOLOGICAL VALUES - SUMMARY


Ecological Values

Numerous ecological surveys were undertaken over the site as part broader concept planning for the Spring Mountain project. In addition, pre-clearance flora and fauna surveys for V11 were undertaken by Saunders Havill Group and Queensland Fauna Consultants, respectively. The following comments summarise the ecological values of the V15 site:

- The V11 site is mapped as containing vegetation comprised of composite Of Concern RE12.9-10.2/12.9-10.7/12.9-10.19 and with Least Concern RE12.9-10.19a.
- Species recorded within the canopy are dominated by *Corymbia citriodora* (Spotted Gum), and *Eucalyptus crebra* (Narrow-leaved Ironbark). Sub-species include scattered *Corymbia henryi* (Large Leaf Spotted Gum), *Eucalyptus seeana* (Narrow Leaf Red Gum), and *Eucalyptus Fibrosa* (Red-leaved Iron-Bark)
- Mapped waterway features were infested with weeds, particularly *Lantana camara* (Lantana) however retained an increased density of *Lophostemon suaveolens* (Swamp Box) and a number of scattered *Eucalyptus tereticornis* (Forest Red Gum) specimens. Areas containing *Lantana camara* recorded very few native flora species within the ground layer.
- The ridgelines and slopes within the assessment area contained a mix of *Corymbia* and *Eucalypt* species with patches of dense understorey of *Acacia* species.
- Rock outcrops were observed along were targeted during the field the survey due to these areas being preferred habitat for a number of the listed flora species including *Plectranthus harbrophyllus* (*Plectranthus*) and *Marsdenia coronata* (Slender Milk Vine). None of these species were recorded on site.
- No State or Commonwealth threatened flora or fauna species were identified within V11 as part of historical and pre-clear surveys.

Regional Ecosystem Descriptions

Least Concern RE 12.9-10.19a	<i>Corymbia henryi</i> +/- <i>Eucalyptus fibrosa</i> subsp. <i>Fibrosa</i> , <i>Corymbia citriodora</i> subsp. <i>Variagate</i> , <i>Eucalyptus siderophloia</i> , <i>Eucalyptus crebra</i> open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
Least Concern RE 12.9-10.2	<i>Corymbia citriodora</i> subsp. <i>Variagate</i> open forest or woodland usually with <i>Eucalyptus crebra</i> . Other species such as <i>Eucalyptus tereticornis</i> , <i>Eucalyptus moluccana</i> , <i>Eucalyptus acmenoides</i> and <i>Eucalyptus siderophloia</i> may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of <i>Lophostemon confertus</i> (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
Of Concern RE 12.9-10.7	<i>Eucalyptus crebra</i> +/- <i>Eucalyptus tereticornis</i> , <i>Corymbia tessellaris</i> , <i>Angophora leiocarpa</i> , <i>Eucalyptus melanophloia</i> woodland. Occurs on Cainozoic and Mesozoic sediments.



Extract: Regional Ecosystem Mapping



Photo: Rock outcrops within the ridgeline areas observed along western facing slope.



05 ENVIRONMENTAL MANAGEMENT


Management – General

This SBMP-V11 sequences through details on a number of site specific outcomes for fauna, vegetation management and operational controls associated with the development of V11. Logically, the document works through construction processes and has been prepared as a sub-plan to the SBMP for Spring Mountain prepared by Yurrah.

Environmental Training

This SBMP-V11 is to be issued to all site contractors (and sub-contractors) and kept within site construction offices. Elements of compliance with the document will form part of the responsibility of the Principle Site Contractor. Training on the management measures outcomes in this SBMP-V11 will occur as part of the broader site environmental management and workplace health and safety procedures. This will include the following steps:

1. Copy of the SBMP-V11 made available to all site contractors (and sub-contractors)
2. Outline of the SBMP-V11 and its requirement relative to the site and / or particular scope of a contract forming part of the site induction requires contractors to read, acknowledge and sign the document prior to commencement of site works.
3. Requirements of the SBMP-V11 to be incorporated into workplace checklists, work method statements and toolbox talks.
4. Weekly review and report on compliance with the SBMP by the Principle Contractor.



Spring Mountain Risk Management Process

Adaptive Management

Adaptive management refers to a way of managing natural resources where management actions are regularly revised and, if necessary, modified based on monitored changes in environmental condition and/or changes in base knowledge which underpins the original management approach. This SBMP has been based on, as far as practical, the current state of knowledge of the species ecology and best practice habitat management approaches. When new facts emerge from future research, they should be immediately integrated into the plan so it remains consistent with the current state of knowledge (and best practice).

Statutory Requirements

Activities associated with this SBMP will comply with the relevant provisions of legislation and regulations and policies of the following:

- Environment Protection and Biodiversity Conservation Act 1999 (Cth) with regard to species listed under the provisions of this Act;
- Nature Conservation Act 1992 (Qld) with regard to species listed under the provisions of this Act;
- Biosecurity Act 2014 (Qld) with regard to weeds and pests; and
- The requirements of the Commonwealth, State and /or Local Government decision notices including any relevant “conditions of approval”.

Roles and Responsibilities

Proponent	Lendlease Communities Pty Ltd	Lendlease Communities Pty Ltd Contact: John Kibble
Contractor	Appointed party or company that performs the construction works on site and included all employees of the Contractor and sub-contractors.	Shadforth Civil Contact: Tony Hopper
Site Supervisor	Appointed party contracted by the Proponent to oversee daily site operations and site management.	Arcadias Contact: Christo Louw
Environmental Representative	Appointed party contracted by the Proponent to oversee environmental compliance.	Saunders Havill Group Contact: Andrew Craig
Fauna/Spotter Catcher	Appointed Contractor employed to implement fauna welfare responsibilities with vegetation clearing operations. The Fauna Spotter Catcher is a person who holds a rehabilitation permit with an extended authority issued by EHP specifying the gilder may take, keep or use an animal whose habitat is about to be destroyed by a human activity.	Queensland Fauna Consulting Contact: Bryan Robinson
Koala Spotter	Appointed Contractor employed to implement Koala welfare responsibilities associated with vegetation clearing operations. The Koala Spotter is a person who holds a tertiary qualification in Biology or Zoology, or who is demonstrably experienced in the identification and location of Koalas in their natural habitat and has an authorisation from EHP to conduct such activities. For example, demonstrably experienced may include a Koala keeper employed by a licensed wildlife exhibitor (i.e. zoo) may be capable of demonstrating competence in locating Koalas.	Queensland Fauna Consulting Contact: Bryan Robinson
Council	Ipswich City Council (ICC)	Ipswich City Council (ICC) Contact: Tim Foote

06 PRE-CLEARANCE - VEGETATION MANAGEMENT

P1– Vegetation Management (General)

Vegetation clearing must be undertaken in accordance with approved plans to ensure protection of areas of ecological significance and agreed retained linear open space corridors. Habitat trees where marked for retention must not be damaged as a result of tree clearing and or are to be removed at the specification and control of the appointed Fauna Spotter.

Table 1 describes the relevant management requirements to address this issue.

Objective

1. To identify clearing in the plans and specification, trees to be retained and trees to be cleared. Areas of retention should be clearly marked and fenced.
2. To ensure that all contractors understand the requirements of protection and retention and install protective devices to ensure no additional clearing occurs.
3. To ensure that the work program is such as to minimise the time between when clearing occurs and the cleared ground is stabilised.
4. To ensure that cleared material is mulches or wood-chipped as appropriate for recycling
5. To protect linear open space from construction damage and run-off.

Management Strategy

- Clearing to be undertaken in accordance with measures outlined in the EPBC Management Plans.
- Install stormwater management devices as per V11 - Stormwater Management Plan.

Performance Indicators

- Integrity of protective devices.
- Existing vegetation and trees retained in good health, with no scars from earthworks machinery and no erosion and sediment deposited within linear open space/retention areas.

Clearing activities should be undertaken in accordance with the with all management plan requirements and associated approval conditions. This SMBP shows the phase 1 clearing of V11 of Spring Mountain. It is acknowledged this clearing line is offset 10m from the ultimately clearing line. Additional lineal clearing will be completed as part of phase 2 works.



Photo: Control clearing of vegetation



Photo: Erosion control to cleared batter



Photo: Tree protection and erosion fence

Table 1: P1: Vegetation Management (Clearing and Protection)

Issue	Vegetation Management – Clearing and Protection	Responsible Person	Timing
Implementation Requirements	Ensure protective devices are installed and maintained in functional condition.	Contractor	During Clearing & Construction
	Monitor and report on the success, protection and retention, and integrity of protective devices such as fences and sediment fences through		
Monitoring	Weekly inspection and log.	Contractor	During Clearing & Construction
Reporting	Monthly (until operation).	Contractor	During Clearing & Construction
Corrective Action	Repair, replace or reinstate protective devices.	Contractor	During Clearing & Construction
	Appropriate treat any damage to trees or vegetation marked for retention as required.	Contractor	During Clearing & Construction

06 PRE-CLEARANCE - VEGETATION MANAGEMENT



06 PRE-CLEARANCE - VEGETATION MANAGEMENT

PAGE INTENTIONALLY BLANK

07 PRE-CLEARANCE - FAUNA MANAGEMENT

P2 – Protection of MNES Fauna (Koala and Grey-headed Flying Fox) and Native Wildlife (Vegetation Clearing)

Clearing of native vegetation has the potential to result in direct injury or death to fauna. Clearing of vegetation for the purposes of preparing development areas also has the potential to result in incidental damage to adjacent habitats to be retained.

Development protocols to respond to injured wildlife must be prepared prior to vegetation clearing operations. It is expected that some of these protocols are likely to be applicable to responses required for all injured fauna (including Koala) and must be included within the Animal Welfare Plan (AWP) to be prepared by the appointed fauna spotter catcher.

Table 2 describes the relevant management requirements to address the protection of terrestrial fauna, specifically Koala, during vegetation clearing and

Objective

- To minimise and mitigate adverse direct and indirect effects of vegetation clearing on terrestrial including Koala and Grey-headed Flying-fox, during clearing and construction.
- Prevent mortality or injury to terrestrial wildlife, specifically Koala.

Management Strategy

- Prevent damage and/or disturbance to native vegetation and associated habitats outside clearing areas.
- Prevent damage and/or disturbance to native vegetation and associated habitats outside clearing areas.
- Clearing and construction operations are employed to maximise animal welfare and reduce fauna mortality.
- Informal all personnel of site environmental responsibility.
- Reuse hollows and large rocks for habitat in retained habitat areas/linear open space.
- Safe fauna movement opportunities are provided within linear open space to prevent fauna moving through construction areas.
- Direct clearing activities from open area to less open areas allowing fauna to natural seek shelter in conservation land and linear open space/retained habitat
- Provision of permanent and temporary fencing in accordance with the V11- Vegetation Management

Clearing Plan
Undertake works in accordance with V11 –Direction of Clearing Plan and install fencing in accordance with V11- VMCP.

Performance Indicators

- Prevent fauna mortality and disturbance to terrestrial fauna.
- No injury or death of Koala.
- No damage to linear open space/retained habitat.
- No disturbance to native vegetation outside permitted clearing footprints.

Fauna Management

Lendlease Communities Pty Ltd commits to the use of leading practice methods and processes for the role of Wildlife Spotter Catchers in the engagement of any contractors for native vegetation clearing works. The standards and requirements outlined in this Specification Note are acknowledged as above minimum requirements in most Local Government areas and are applicable despite lessor requirements listed within individual project approval packages.

As a minimum specification Wildlife Spotter Catchers will retain the following Queensland State Government Permits:

- Animal Ethics
- Scientific Purposes Permit
- Scientific User Registration
- Damage Mitigation Permit
- Rehabilitation Permit

Wherever practical all clearing works will be coordinated in general accordance with applicable site based components of the DRAFT Code of Practice for the welfare of animals affected by land-clearing and other habitat impacts prepared by the Australia Zoo Wildlife Warriors and Voiceless (and or any contemporary Industry based final version of this Draft Code). This includes mandatory controls on the timing and sequencing of clearing works integrated with a regimented series of fauna management protocols implemented by registered Fauna Spotter / Catchers. The following procedural stages listed in the Draft Code are to be applied to clearing works on all Lendlease Communities Pty Ltd projects:

Action 1 – Engagement Wildlife Spotter Catcher

Action requires that the developer (and or the developer's representative through the principal contractor) engage a Wildlife Spotter Catcher with full registrations and licences provided in accordance with the Queensland Government's

National Parks and Wildlife Services. A Registered Wildlife Spotter Catcher engaged shall have the minimum permits listed in this specification.

Action 2 – Wildlife Spotter Catcher to Prepare a Wildlife Protection and Management Plan (WPMP)

The WPMP should be submitted to the Queensland Department of Environment and Heritage Protection (EHP) or relevant authority and or stakeholder. The WPMP should include the following information:

- Description of the project with reference to impacts on wildlife or wildlife habitat;
- Pre development plan of the site showing habitat areas, features, corridors, riparian habitats and adjacent areas;
- Results of any fauna surveys including pre-clearance surveys; and
- A wildlife and habitat impact assessment based on the proposed development works.

Action 3 – Prepare a Wildlife and Habitat Impact Mitigation Plan

Following completion and endorsement of the WPMP the Wildlife Spotter Catcher should prepare a more specific Wildlife and Habitat Impact Mitigation Plan, which will include details on:

- Measures required to be completed to minimise wildlife and habitat impacts during operational works;
- Wildlife capture and removal plan;
- Contingency plan for wildlife requiring euthanasia, other veterinary procedures or captive care;
- Wildlife storage and housing plan;
- Wildlife release and disposal plan; and
- Post works measures to minimise impacts on wildlife.

Lendlease Communities Pty Ltd support the use of innovative leading practice methods minimising and mitigating impacts on all native fauna during clearing operations.

Action 4 – Wildlife Spotter Catcher Role at Pre-Start Meeting

Prior to the commencement of any construction works, a pre-start meeting is to be held between the project manager, site fore-person, plant operators and applicable Local and State Government representatives. At the pre-start meeting, the Wildlife Spotter Catcher is to outline the clearing process and the requirements of the WPMP.

Action 5 – During Construction

The Wildlife Spotter Catcher is to be on-site during all phases

of construction which involve potential impacts on wildlife or habitat (unless otherwise specified by the appointed Wildlife Spotter Catcher. This will enable to the Wildlife Spotter Catcher to make any necessary adjustments to the approved Clearing Management Plans and WPMP to cater for any specific issues encountered during the clearing works.

Action 6 – Post Works Reporting

During the course of all site works, including the pre-clearance surveys, the Wildlife Spotter Catcher is to keep an accurate record of all animals encountered, captured, incidents and disposals for each stage of the project. The records should form part of the Wildlife Management Report to be issued under licence requirements to the State Government. The Wildlife Management Report should consist of the following 3 sections, where they are applicable to the project:

- Wildlife Habitat Management Plan** – Aspects of the planning, design, construction and ongoing operation of the project in which risks to wildlife have been identified. This plan should also include recommendations and outline the type, frequency and timeframes for monitoring

- Wildlife Capture and Disposal Plan** – Should contain the following details for each captured animals:

- Species
- Identification name or number
- Sex (M, F or unknown)
- Approximate Age or Age Class (neonate, juvenile, sub-adult, adult)
- Time and date of capture
- Method of capture
- Exact point of capture (GPS coordinates)
- State of health
- Incidents associated with capture likely to affect health
- Veterinary intervention or treatments
- Time held in captivity
- Disposal method (euthanasia, translocation, re-release)
- Date and time of disposal
- Detailed of disposal (GPS points of release)
- For released animals, location relative to point of capture

- Animal Injury and Euthanasia Report** – similar details for the Wildlife Capture and Disposal Plan should be included in this report.

07 PRE-CLEARANCE - FAUNA MANAGEMENT

Koala Management & Welfare

While clearing activities aim to protect and minimise impacts to all terrestrial fauna, specific management measure for Koala are required as part of the EPBC approval and have been specified within the Fauna Management Plan, prepared by **Saunders Havill Group** which should be read in conjunction with the plan.

Key outcomes within the FMP for Koala include:

- Koalas on site are protected
- Koala habitats are protected, maintained and their integrity enhanced.
- The abilities for Koalas to move into, within and out of the site is maintained.
- All persons involved in construction and operation of the development are aware of the site values, their potential to impact on Koalas and their habitats, and their responsibilities in regard to procedures and strategies within approved management plans.



Koala Signage



Significant Tree Protection Fencing




Fauna Spotter During Tree Clearing



Fauna Spotters Retrieving Fauna



Fauna Exclusion Fencing



Construction fencing detail



Fauna Signage



Fauna Exclusion Fencing

07 PRE-CLEARANCE - FAUNA MANAGEMENT

Table 2: P2 – Protection of MNES Fauna and Native Wildlife (Vegetation Clearing)

Issue	P2 - Protection of MNES Fauna and Native Wildlife	Responsible Person	Timing
Implementation Requirements	No vegetation removal shall occur until relevant approvals have been obtained All permit conditions will be followed	Proponent	Prior to Clearing
	To prevent damage and / or disturbance to native vegetation and associated habitats outside clearing areas: <ul style="list-style-type: none"> a. Clearing boundaries will be delineated on all drawings and in the field to define the authorised clearing extent. b. Installation of vegetation clearance markers (e.g. high visibility poly-web fencing) prior to the commencement of vegetation clearance to identify and protect remnant vegetation for retention. c. Along the interface between clearing precincts and open space / Environmental Corridors, trees are to be felled towards the clearing precinct to avoid damage to these areas. d. Clearing vegetation is to be stockpiled so as not to impede damage to drainage channels. 	Contractor	Prior to Clearing & During Clearing
	No clearing of vegetation is to commence without the presence of an EHP approved Fauna Spotter Catcher, or where clearing includes non-juvenile Koala habitat trees, a Koala Spotter. <ul style="list-style-type: none"> a. An appointed Site Superintendent will be responsible for ensuring that all trees scheduled for removal will be checked on the day of their removal for the presence of fauna by an EHP approved Fauna Spotter Catcher / Koala Spotter as vegetation characteristics dictate. b. The EHP approved Fauna Spotter will check and clear vegetation prior to its felling and, if required, will relocate native wildlife (other than Koala) into appropriate habitat areas within the site which are to be retained. In the case of a Koala being present, translocation of the individual/s must occur in accordance with requirements for Koala. c. Hollow-bearing (habitat) trees are to be identified in the field and by plan prior to commencement of clearing operations. These shall be marked and dismantled using a cherry picker and a suitably qualified arborist and Fauna Spotter Catcher. If fauna is present, the tree will either be left standing overnight to allow the animal to leave via their own volition, or will be encouraged from the tree by shaking or other methods deemed suitable by the fauna spotter. Where no signs of fauna are identified, machinery operators will be instructed to fell trees in a manner directed by the fauna spotter to minimise potential risk to fauna. 	Fauna Spotter Catcher	Prior to Clearing
	All construction personnel shall attend environmental training as part of the site induction process prior to entering the work site. As part of this training, all personnel will be instructed on their obligations in regard to vegetation clearing protocols and to protect native fauna. Areas identified for vegetation clearance are to be clearly defined and detailed in site inductions.	Contractor	Prior to Clearing
	Conduct vegetation clearing in sequential stages for sites with an area of more than 3 hectares. Vegetation clearing is to conform with the following: <ul style="list-style-type: none"> d. The direction of clearing should be away from threatening processes or hostile environments, and towards the clearing precinct to avoid damage to adjacent retained habitat links, ensuring that: <ul style="list-style-type: none"> i. Fauna are not required to cross roads or move through developed areas or disturbed areas. Such as residential areas or areas that require movement of greater than 100m over cleared ground to reach suitable habitat; ii. Fauna area not left occupying an "island" of habitat between hostile environments, such as a road and a cleared area, unless there are no other more suitable habitat areas in which to direct fauna, and iii. Fauna can safely leave the site of clearing and relocate to adjacent habitat. 	Contractor	During Clearing

07 PRE-CLEARANCE - FAUNA MANAGEMENT

Table 2: P2 – Protection of MNES Fauna and Native Wildlife (Vegetation Clearing)

Issue	P2 - Protection of MNES Fauna and Native Wildlife	Responsible Person	Timing
	<ul style="list-style-type: none"> e. Cleared vegetation is to be stockpiled so as not to impede fauna movement. f. Where vegetation to be cleared includes non-juvenile Koala habitat trees, implement sequential clearing as per the requirements for Koala. 		
	Companion animals (e.g. dogs) are to be banned from all construction areas.	Contractor	At all times
	Vehicle access within retained habitat/linear open space will be limited and appropriately signed.	Contractor	Prior to Clearing & During Clearing
	<p>Conduct vegetation clearing in accordance with Section 4 of the Spring Mountain FMP (prepared by Saunders Havill Group dated July 2015) which outlines specific implementation requirements for <u>Koala</u> including clearing in sequential stages for sites. For a site more than 6ha vegetation clearing is to conform with the following:</p> <ul style="list-style-type: none"> a. Is carried out in a way the ensures Koalas on the area being cleared have enough time to move out of the clearing with without human intervention and involves <ul style="list-style-type: none"> i. Ensuring not more than 3ha or 3% of the sites area (whichever is greater) in any one stage ii. Ensuring that between each stage and the next there is at least one period of 12 hours at starts at 6pm on a day and ends at 6am on the following day, during which no trees are cleared on the site b. Is implanted in a way that ensures, while clearing is being carried out, appropriate habitat links are maintained within the clearing site and between the site and its adjacent areas allowing Koalas living on the site to move out of the site c. Ensures that no tree in which a Koala is present, or a tree with a crown overlapping a tree in which a Koala is present, is cleared until the tree is vacated by the Koala. d. Ensures that vegetation clearing is directed away from threatening processes, or hostile environments, and towards any retained vegetation or habitat links, ensuring that: <ul style="list-style-type: none"> i. Koalas are not pressured, through loss of habitat, to cross roads or move through developed or disturbed areas, such as residential areas or areas that require movement of greater than 100m over cleared ground to reach suitable habitat; ii. Koalas are not left occupying an "island" of habitat between hostile environments, such as road and cleared areas, unless there are no other more suitable habitat areas in which direct Koalas; and iii. Koalas can safely leave the site of clearing and relocate to adjacent habitat. e. The Koala spotter is responsible for ensuring, throughout the duration of clearing operations, that no tree in which a Koala is present, or a tree identified as being a risk to Koalas if felled, should not be felled, damaged or interfered with until the Koala has moved from the felling site of its own volitation. f. Where a Koala is present in a tree scheduled for removal, the tree will be marked with distinctive flagging tape (and other advisory means as required) and machinery operators will be briefed on the location of such trees and it will be clearly confirmed with operators that the subject tree(s) are to remain undisturbed until the Koala has moved of its own volition (where the strategy is to allow the Koala to move of its own accord, overnight). On the following day, such trees are to be checked again prior to their removal and, if necessary, the procedure is to be repeated until the Koala has moved g. A Koala spotter is not to be involved in the clearing of vegetation while they are responsible for identify Koalas on site. 	Contractor / Fauna Spotter Catcher/ Koala Spotter	During Clearing

07 PRE-CLEARANCE - FAUNA MANAGEMENT

Table 2: P2 – Protection of MNES Fauna and Native Wildlife (Vegetation Clearing)

Issue	P2 - Protection of MNES Fauna and Native Wildlife	Responsible Person	Timing
	<p>A requirement that a permit to interfere with wildlife from EHP will be mandatory for the wildlife handling activities as will the appropriate Animal Ethics Permit from DAF. Construction personnel shall not attempt to handle any wildlife.</p> <ul style="list-style-type: none"> a. Fauna / Koala handling and relocation activities must only be undertaken by those identified on a current site-specific Damage Mitigation Permit (Removal and Relocation of Wildlife) from EHP. b. Koala Spotter/Fauna Spotter Catchers are required to relocate injured wildlife to the nearest designated veterinary clinic of wildlife hospital. Full contacts will be provided within the AWP. c. A register of fauna incidents / interactions is to be maintained daily during clearing operations. 	Fauna Spotter Catcher/Koala Spotter	During Clearing & Construction
	The timing of vegetation clearance should be selected in order to minimise impacts (direct and indirect) to affected fauna habitats during optimum breeding period.	Contractor	During Clearing
	Avoid clearing of vegetation between the hours of 6pm and 6am.	Contractor	During Clearing
Monitoring	For each day of native vegetation clearing operations, a daily audit log is to be completed by the Contractor either prior to, or on completion of daily operations. Audit of key requirements, e.g. clearing contained within designated limits, integrity of clearing boundary devices, no damage to vegetation outside clearing boundary, Fauna Spotter Catcher present.	Contractor	During Clearing
Reporting	Animal Welfare Plan is prepared prior to clearing operations by the appointed Fauna Spotter Catcher.	Proponent / Fauna Spotter	Prior to Clearing
	Weekly report by the Fauna Spotter Catcher/ Koala Spotter to the Contractor on the clearing of any native vegetation and any animals encountered, injured or relocated is to be submitted.	Contractor	During Clearing
	Monthly report by the Contractor the Site Superintendent on native vegetation operations, including compliance, non-compliance incidents (fauna injury and responses) and corrective actions, outcomes of Fauna Spotter Catcher activities.	Contractor	During Clearing & Construction
	Bi-annual report by the Site Superintendent to the Proponent. Report to consider incident patterns, if any, and provide recommended solutions and a description of the corrective actions taken.	Contractor	During Clearing & Construction
	Annual site audit by the Environmental Representative and report to the Proponent	Environmental Representative	During Clearing & Construction
Corrective Action	In the event that monitoring identifies practices inconsistent with the strategies developed for this FMP, the Contractor shall take the necessary corrective steps and note them in the monthly report to be reviewed by the Site Superintendent.	Contractor	During Clearing & Construction
	In the event that monitoring identifies practices inconsistent with the strategies developed for this SBMP, the Contractor shall take the necessary corrective steps and note them in the monthly report to be reviewed by the Site Superintendent	Contractor	During Clearing & Construction

08 FAUNA MANAGEMENT - CONSTRUCTION

P3 – Maintenance of Safe Wildlife Movement Opportunities (Site Preparation Operations)

The following suite of best practice measures will be employed throughout the site to minimise fauna habitat fragmentation, facilitated fauna movement and reduce related injury and mortality. Management requirements are considered in the context of:

- Site preparation operations (i.e. during vegetation clearing and earthworks phases); and
- Design treatments and strategies for the built phase of the development

Table 3 describes the relevant management requirements in regard to site preparation operations. The following should be read in conjunction with the requirements for Koala design treatments and strategies for the built phase of the development.

Retention and rehabilitation of the Mountain Creek Corridor to the west, in addition to the 293ha of offset land for Conservation to the south, will occur as a result of the Spring Mountain development to maintain fauna movement and connectivity within and between the development site

Objective

1. To avoid the impact of habitat fragmentation by roads and maintain safe movement opportunities for native wildlife (including Koala and Grey-headed Flying-fox) between linear open space.
2. To maintain fauna movement opportunities within retained habitat areas and minimise fauna movement opportunities through site preparations.

Management Strategy

- Develop a track plan for retained habitat areas/linear open space which allows fauna movement to be maintained
- Restrict access to retained habitat areas/linear open space for environmental management only.
- Reduce road speeds
- Increase driver awareness and education

Performance Indicators

- Minimal fauna mortality.

Temporary Fencing

Prior to the commencement of vegetation clearing a temporary fauna exclusion fence will be erected around the area of clearing and works and be maintained until the completion of major civil works. The purpose of the fence is to minimise any native fauna (including koala) from entering into the clearing and or post clearing construction zone during a time when potential risks of impact are at their highest.

The fencing proposed is a “floppy-top” temporary fauna exclusion fencing as per the details and photos shown on this drawing sheet. This fencing type is preferred as it continues to allow any fauna within the impact zone to exit, however prevents new or re-entry once the fence is erected. The fencing type can also be erected along random alignments and relocated to new areas as the clearing areas expand in future clearing and development events. This fencing type has been successfully used as a temporary barrier on other koala related projects within the vicinity of major roads and housing areas.



Fauna exclusion fencing



Controlled clearing access track

Table 3: P3 – Maintenance of Safe Fauna Movement Opportunities – Site Preparation Operations

Issue	P2 – Maintenance of Safe Fauna Movement Opportunities – Site Preparation Operations (Roads and Vehicle Interactions)	Responsible Person	Timing
Implementation Requirements	A site access plan is to be developed for the Environmental Corridors.	Proponent	Prior to Clearing
	Site protocols are to be established which restrict authorised area access to the approved track network identified with the plan.	Contractor	Prior to Clearing
	All construction personnel shall attend environmental training as part of the site induction process prior to entering the work site. As part of this training, all personnel will be instructed on their obligations in regard to vehicle movement restrictions and construction speed limits.	Contractor	Prior to Clearing
	Erect temporary exclusion fencing around the area of clearing and works and be maintained until the completion of major civil works.	Contractor	Prior to Clearing
	Vehicle movements outside designated operational areas (other than for land management purposes) will be prohibited.	Contractor	During Clearing & Construction
	Road speeds throughout construction areas and through retained habitat areas will be restricted to 50km/hr.	Contractor	During Clearing & Construction
	Strategic use of awareness signage is to be implemented along the interface between operational areas and Environmental Corridors and access restriction signage at all track entry points to Environmental Corridors during construction works.	Contractor	During Clearing & Construction
Monitoring	Proposed construction access roads will be subject to design treatments to ensure safe fauna crossing opportunities. Construction of an elevated portion (or portions) in the form of bridging structures (culverts) in associated with guide fencing will be incorporated to ensure the provision of safe crossing opportunities.	Contractor	During Clearing & Construction
	Weekly inspection and log.	Contractor	During Clearing
	Reporting	Monthly report by the Contractor to the Site Superintendent in regard to development / maintenance of structures implemented to facilitate fauna movement, review of fauna / vehicle incident patters, if any, and provide recommended solutions, an a description of corrective actions taken.	Contractor
Bi-annual audit report by the Site Superintendent to the Proponent. Report to include compliance with site access restrictions, integrity of structure implemented to facilitate fauna movement, review of fauna/ vehicle incident patterns, if any, and provide recommended solutions, and a description of corrective actions taken.		Contractor	During Clearing
Annual site audit by Environmental Representative and report to the Proponent.		Environmental Representative	During Clearing & Construction
Corrective Action	In the event that monitoring identifies practices inconsistent with the strategies developed for this SBMP, the contractor shall take the necessary corrective steps and note them in the monthly report to be reviewed by the Site Superintendent.	Contractor	During Clearing & Construction

09 THREATENED FLORA MANAGEMENT

P5 – Threatened Flora Management

Plectranthus habrophyllus, a herb listed as Endangered under the EPBC Act, has been recorded at several locations across the Spring Mountain project site. Core populations have been identified within Core Conservation areas by Yurrah. The majority of these locations are associated with waterways within linear open space and the habitat is to be protected.

Pre-clearance Survey

In accordance with the EPBC approved Threatened Flora Management Plan, prepared by Yurrah, pre-clearance surveys for each development precinct must occur by a suitable qualified person prior to the commencement of clearing. Any additional individuals must be recorded and translocated where necessary.

Translocation

Where plants are located within the development footprint or near the edge of the footprint, and are at risk of impact, these plants will be translocated to establish a new population in suitable habitat within the proposed Linear Open Space. The habitat for both translocated individuals and in situ individuals will be protected within a Core Conservation Area.

As an added habitat protection measures, Buffer Areas, with an offset width of 20m, will be established around Core Conservation Areas. No Go Zones must be marked out by the 20m buffer around known populations within Core Conservation areas. No work apart from conservation management activities is to be permitted within Core Conservation Areas.

Clearing and Construction

Plectranthus habrophyllus is to be protected from impacts of construction. Stormwater Management Plans, Bushfire Management Plans and Weed Management are to address threatened flora management.

Table 5 describes the relevant management requirements to address this issue.

Objective

1. To encourage the locally resident populations of threatened flora species to increase at a natural rate to a desired level on site.

Management Strategy

- Threatened flora habitat to be protected through the approved Threatened Flora Management Plan
- Recognise and protect all linear open space through management of interface between linear open space and development for bushfire, weeds and access issues.
- Establish Core Conservation Areas and Buffer Areas at threatened flora locations to target management activities.
- Design a network for fire-trails to defined spatial blocks to prevent damage caused by uncontrolled fire and allow access for maintenance.
- Awareness and education of threatened flora presence.
- Ensure all responsible persons are aware of the significance of this issue and are fully aware of any likely impacts of scheduled works.

Performance Indicators

- 0% weed cover in Core Conservation Areas and Buffers
- No evidence of damage from stormwater run-off construction
- Recruitment of threatened flora seedlings in Core Conservation Area
- No damage from uncontrolled access
- Condition of protective fencing remains undamaged.

Pre-clearance surveys for V11 were undertaken by Saunders Havill Group in May 2017. No *Plectranthus habrophyllus* individuals were located within the project area.



Photo: *Plectranthus habrophyllus* (listed as Endangered under the EPBC Act (Cth))

09 THREATENED FLORA MANAGEMENT

Table 5: P5 – Threatened Flora Management

Issue	P4 Threatened Flora Management	Responsible Person	Timing
Implementation Requirements	<p>Core Conservation Areas located within 20m of land proposed for uses other than conservation, identified as areas for additional interface management including:</p> <ol style="list-style-type: none"> 1. A detailed survey of threatened plant locations by a registered surveyor. 2. Where interfacing with residential, a fence with a minimum 50% transparency to be erected along interface boundary. Signage to be erected identifying area as 'Significant Ecological Area' and 'Dumping of Rubbish Prohibited' and where further information can be obtained. 3. Where interfacing with road verge or park landscaping, design and plant selection considers and avoids any potential impact upon the threatened flora species. Landscape plant species selected will be non-invasive, existing trees to be retained where possible to maintain microclimate, and clear edge formed that discourages access. Mulch to be preferably sourced from the site and is to be weed free. 	Proponent	Design /Prior to Clearing &
	<p>Undertake pre- clearing surveys.</p> <ol style="list-style-type: none"> 1. Once the line of clearing (including construction of parks, pedestrian tracks and fire trails) is marked out by a registered surveyor, an additional survey for threatened species is to be undertaken within the clearing area, and Linear Open Space within 10m of the clearing line. 2. Additional individuals, or groups of individuals located to be recorded with a GPS, given a unique ID number, and flagged with marking tape. Where necessary individuals will be translocated in accordance with protocols in the Threatened Flora Management Plan. 3. The boundary of the Core Conservation Areas will be adjusted as necessary (if not within construction footprint), to include any additional individuals located during of the pre-clearing survey. 	Proponent	Prior to Clearing
	<p>Establish No Go Zones.</p> <ol style="list-style-type: none"> 1. Core Conservation Areas less than 20m from of the clearing and construction footprint will be identified on construction drawings and through signage on site as 'No Go Zones'. Their associated Buffer Areas will be identified as 'Proceed with Caution Zones'. 2. Work within the Buffer Area will require supervision by the Project Ecologist. 3. No work apart from conservation management activities is to be permitted within the Core Conservation Areas. 	Contractor	Prior to Clearing
	<p>Erect exclusion fencing and signage.</p> <ol style="list-style-type: none"> 1. Where Linear Open Space has not been fenced as part of general vegetation protection, temporary fencing must be installed around the Core Conservation Area, where practical, and necessary (i.e. steep terrain may form natural barrier). The temporary fence shall be a minimum of star pickets with 3 strand wire and high visibility mesh attached to the top wire (with minimum gap of 500mm along the bottom) and erected prior to clearing. 2. The required alignment and extent of the fencing is to be undertaken in consultation by the project ecologist and inspected before the start of clearing. 3. Signage is to be attached to fencing clearly identifying the site as a significant ecological area and a 'No Go Zone', and no entry permitted unless approval given by Proponent. Mapping will be produced identifying location of threatened flora and alignment of protective fencing during detailed design for each Phase of the Spring Mountain 	Contractor	Prior to Clearing

09 THREATENED FLORA MANAGEMENT

Table 5: P5 – Threatened Flora Management

Issue	P4 Threatened Flora Management	Responsible Person	Timing
	Stormwater Management controls to be installed through implementation of an Approved Stormwater Management Plan for Spring Mountain. <ol style="list-style-type: none"> The Stormwater Management Plan will outline management required to ensure water quality and quantity flowing into Core Conservation Areas and all areas of proposed conservation are at predevelopment levels. All stormwater management devices are to be installed and inspected prior to clearing and construction. Stormwater management devices to be regularly checked and maintained to ensure they perform their intended function. 	Contractor	Prior to Clearing
	Instruct all site workers and visitors in the presence and significance of threatened species on site, and on the management measures being implemented at the present time. All personal associated undertaking works within a Buffer Area are to be made aware of the presence of threatened plants, and are to be educated on protective measures in place, prior to entering area. No personnel to enter Core Conservation Area without approval.	Contractor	Prior to Clearing
	Fire trails will be installed in accordance with the Final Bushfire Management Plan with locked gates and structures to prevent access to vehicles, other than emergency and maintenance vehicles, into all Linear Open Space areas.	Contractor	During Clearing
Monitoring	Core Conservation Areas and Buffers will be monitored on a 3 monthly basis for the first year, and annual thereafter for 2 years subject to satisfactory performance including: <ul style="list-style-type: none"> Provide general photographic descriptive record Establish permanent sample quadrats located in each management block, according to an agreed sample strategy Confirm the absence of environmental weeds Measure species richness of the ground layer. Measure abundance of flowering threatened species. Measure abundance of threatened species seedlings General observations. 	Contractor	During Construction / Operation
Reporting	Every 3 months by the Environmental Representative to the Proponent for the first year, every 6 months in the second year and once in the third year/	Environmental Representative	During Clearing & Construction
	Annually by the Proponent to the DoE including non-conformances, corrective actions and assessment of monitoring results.	Proponent	During Clearing
Corrective Action	In the event that monitoring identifies practices inconsistent with the strategies developed for this SBMP, the contractor shall take the necessary corrective steps and note them in the monthly report to be reviewed by the Site Superintendent.	Contractor	During Clearing & Construction

10 FLORA & FAUNA CHECKLIST

Pre-Clearance Checklist:

This Site Based Management Plan (V11) contains only a small portion of information included within existing assessment management plans for Spring Mountain. Subsequently, the volume of requirements remains complex and overlapping. To ensure compliance with approval requirements and provide a record trail for reporting to the Commonwealth Department of the Environment the following pre-clearance checklist is to be completed with each phase of works.




The checklist is to be completed by the principal contractor and requires sign off by the Environmental Coordinator and Fauna Spotter. To complete the checklist a number of items need to be issued from various parties to the principal contractor (eg confirmation of pre-clearance surveys).

The pre-clearance checklist is established in a format which enables direct annual reporting to the Department of the Environment and will include a number of attachments.



Springfield Rise - Environmental Pre-Start Checklist

Project Area: Village 6		Date:			
Contractor: Shadforth's		Construction Stage/ Activity:			
Date work is to start:		Early works bulk earthworks			
Date work is to cease:		Compliance			
#	Control Measure	Yes	No	N/A	Comments
1	Are clearing extents marked out and fenced? (N.B. Fencing is required as per ICC permits unless instructed otherwise by Council, Fauna Spotter or Environmental Coordinator)	✓			Completed by Wolter Consulting on DATE
2	Has the fencing of clearing extents demarcation been inspected by the Environmental Coordinator?	✓			Completed by SHG on DATE
3	Has sign off been provided by the Environmental Coordinator for demarcation areas?	✓			See Attachment 1
4	Has certification for pre-clearance flora been provided? (N.B. Exemptions/permits for protected plants under the NCA must be obtained by EHP where works occur in a High Risk Area). Please provide date and reference.	✓			See Attachment 2. EHP Reference: AR082999 22 January 2016
5	Have pre-clearance checks surveys for <i>Plectanthis habrophyllus</i> been completed over the clearing area?	✓			Completed by SHG on 8 July 2015. See Attachment 3.
6	Are there 'no-go' zones identified within the clearing area?		✓		
7	If yes, have 'no-go' zones been demarcated, fenced, signed and inspected by the Environmental Coordinator and Contractor?			✓	
8	Has the appointed Fauna Spotter completed pre-clearance surveys and reports?				
9	Has the appointed Fauna Spotter identified any				






Springfield Rise - Environmental Pre-Start Checklist

Project Area: V11 Development Area (Northern extent)		Date: 10/07/2017			
Contractor: Shadforth's		Construction Stage/ Activity:			
Date work is to start: 11/07/2017		Phase 1 - Early works bulk earthworks (V11 --northern extent)			
Date work is to cease: 4/08/2017		Compliance			
#	Control Measure	Yes	No	N/A	Comments
1	Are clearing extents marked out and fenced? (N.B. Fencing is required as per ICC permits unless instructed otherwise by Council, Fauna Spotter or Environmental Coordinator)	✓			Completed by Wolter Consulting on 30 th June 2017.
2	Has the fencing of clearing extents demarcation been inspected by the Environmental Coordinator?	✓			Completed by SHG on 3 rd and 5 th July 2017.
3	Has sign off been provided by the Environmental Coordinator for demarcation areas?	✓			See Attachment 1.
4	Has certification for pre-clearance flora been provided? (N.B. Exemptions/permits for protected plants under the NCA must be obtained by EHP where works occur in a High Risk Area). Please provide date and reference.	✓			See Attachment 2. EHP Reference: AR098906 (14 March 2017) & EHP Reference: AR0829996 (22 January 2016).
5	Have pre-clearance checks surveys for <i>Plectanthus habrophyllus</i> been completed over the clearing area?	✓			Completed by SHG on 14 th February 2017. See Attachment 3.
6	Are there 'no-go' zones identified within the clearing area?		✓		
7	If yes, have 'no-go' zones been demarcated, fenced, signed and inspected by the Environmental Coordinator and Contractor?			✓	
8	Has the appointed Fauna Spotter completed pre-clearance surveys and reports?	✓			See Attachment 4. Fauna Spotter Catcher Pre-Clearance and Habitat Values Survey, completed by QFC (July 2017)

9	Has the appointed Fauna Spotter identified any sensitive areas for consideration in clearing methods? Please provide a summary.	✓		See Attachment 4. Fauna Spotter Catcher WHIMP, completed by QFC (July 2017)
10	Have all contractors, subcontractors and associated personnel been instructed on environmental procedures and controls?	✓		See Attachment 5. Environmental Awareness Acknowledgement Notice, signed by Shadforth's (July 2017).
11	Has a Council pre-start been completed?	✓		Confirmation from ICC no pre-start is required. See Attachment 6.

NOTE: if the answer to any question (1-5, 7-11) above is NO then the clearing activity will not proceed.

Name	Company	Position	Signature	Date
Dustyn North	W.M.I.	Clearing Contractor		10/07/2017
Graeme Knox	LL	Client Representative		10/7/17
Christo Louw	Arcadis Australia Pacific Pty Ltd	Project Engineer		10/07/2017

CONTRACTOR COORDINATOR:

Name: SAM SCHROEDER

Position: PROJECT MANAGER

Date: 10/07/2017

Signature: 

ENVIRONMENTAL COORDINATOR:

Name: MURRAY SAUNDERS

Position: DIRECTOR

Date: 10.07.17.

Signature: 

FAUNA SPOTTER COORDINATOR:

Name: BETAN ROBINSON

Position: Director

10/7/17



ATTACHMENT I – Demarcation Flagging Inspection Notification

Date: 5 July 2017
Site: Spring Mountain Precinct (Haul Road)
Client: Lend Lease
EPBC Ref: 2013/7057
SHG Ref: 7243
SHG Contact: Murray Saunders (07 3251 9444)

Attention: Ian Murray

Regional Development Manager, Communities
Level 4, Kings Gate,
King Street
Bowen Hills QLD 4006

Springfield Rise: V11 Development Area – Inspection of flagging for demarcation of clearing extents (Phase 1- early works bulk earthworks, 7002 Grande Avenue, Springfield (Lot 2 on SP291381))


Dear Ian,

The *Environmental Management Division* of **Saunders Havill Group** was engaged by **Lendlease Communities** to carry out an inspection of flagging for demarcation fencing for the Phase 1- Early Works Bulk Earthworks clearing extent associated with the V11 Development Area at Springfield Rise.

Flagging of the V11 Development Area Phase 1 clearing extent was undertaken by the appointed surveys, **Wolter Consulting**, on the 26th June 2017. Ecologists from **Saunders Havill Group** checked and reflagged the clearing extent on the 3rd July and 5th July 2017 to confirm works will be undertaken in accordance with relevant Commonwealth and Council permit requirements.

The GPS track log of the inspection extent shown in the plan provided as **Attachment 1**. A post-inspection notification is provided as **Attachment 2** to be kept for your records.


Kind regards,



Murray Saunders
Director – Saunders Havill Group

Attachment I –

Demarcation Fencing Inspection Track Log



ID	Eastings (m)	Northings (m)	ID	Eastings (m)	Northings (m)	ID	Eastings (m)	Northings (m)
1	489442.18	6936978.96	21	489260.95	6936869.99	41	489111.87	6937055.97
2	489437.42	6936967.59	22	489251.88	6936855.53	42	489113.16	6937061.08
3	489432.61	6936953.01	23	489239.30	6936845.29	43	489117.79	6937068.35
4	489424.70	6936946.66	24	489219.94	6936848.90	44	489128.71	6937073.53
5	489417.71	6936932.02	25	489198.68	6936849.79	45	489142.21	6937083.80
6	489413.55	6936931.94	26	489181.98	6936854.65	46	489154.98	6937091.41
7	489407.41	6936919.44	27	489165.08	6936861.44	47	489174.68	6937092.43
8	489402.01	6936932.60	28	489147.34	6936875.77	48	489199.52	6937091.43
9	489397.85	6936940.35	29	489137.44	6936884.74	49	489213.14	6937090.25
10	489394.57	6936947.05	30	489126.68	6936899.45	50	489229.75	6937089.74
11	489389.79	6936956.76	31	489118.69	6936910.85	51	489246.89	6937089.77
12	489375.38	6936949.91	32	489109.15	6936924.47	52	489267.36	6937087.72
13	489360.32	6936941.77	33	489101.87	6936940.00	53	489287.04	6937086.33
14	489346.20	6936931.36	34	489094.71	6936956.83	54	489303.59	6937086.03
15	489332.89	6936921.84	35	489092.01	6936969.48	55	489321.13	6937085.31
16	489320.48	6936913.10	36	489092.33	6936982.09	56	489340.26	6937089.37
17	489310.61	6936904.52	37	489093.01	6937000.60	57	489350.92	6937092.07
18	489296.15	6936894.13	38	489095.55	6937013.53			
19	489285.82	6936889.26	39	489100.09	6937032.02			
20	489271.33	6936878.95	40	489105.80	6937045.61			

Attachment 2 –

Demarcation Flagging Inspection Notification

Area Inspected:	Springfield Rise – 11 Phase 1 (Early Works Bulk Earthworks)
Location:	7002 Grande Avenue, Springfield (Lot 2 on SP291381)
Date of Inspection:	3 rd July and 5 th July 2017
Appointed Surveyor:	Wolter Consulting – Glenn Hanton
Environmental Representative:	Saunders Havill Group – Andrew Craig
Environmental features:	The 11 adjoins a low order stream order to the east. Dense weed infestations were noted throughout this drainage line.

Photos of flagging prior to demarcation fencing:



ATTACHMENT 2 – NCA Flora Survey Report and Exemption Notification

Keira Grundy

From: PALM <palm@ehp.qld.gov.au>
Sent: Tuesday, 14 March 2017 12:47 PM
To: Keira Grundy
Subject: RE: AR098906 - 7522: Exempt Clearing Notification - Springfield Rise V13 Basin & V15
Attachments: image001.png



Applicant: Lend Lease Communities Pty Ltd

Where clearing is to be conducted –
Street Address: Sinnathamby Boulevard, Springfield
Lot/Plan: Lot 200 on SP283567

EHP Reference: AR098906

Dear Ms Grundy,

Thank you for your request for an Exempt Clearing Notification for protected plants.

Please retain this email as acknowledgement of receipt of a protected plant exemption notification submitted under *Section 261ZA of the Nature Conservation (Wildlife Management) Regulation 2006*.

Clearing of a protected plant under this section must be conducted within two years after the flora survey report was submitted to the Department of Environment and Heritage Protection.

It is strongly recommended that for audit purposes you keep this email together with the relevant flora survey trigger map, flora survey report and any other documentation relating to the clearing in question.

Please visit www.ehp.qld.gov.au for information about available online services.

Kind regards,
Aliza

Aliza Nannicelli

Administration Officer

Regulatory Capability and Customer Service Branch | Environmental Services Regulation

Department of Environment and Heritage Protection

P 1300 130 372 (option 4) F (07) 3330 5875 E palm@ehp.qld.gov.au

Lvl 9, 400 George Street BRISBANE QLD 4000

GPO Box 2454, BRISBANE QLD 4001

www.ehp.qld.gov.au

From: Keira Grundy [<mailto:keiragrundy@saundershavill.com>]

Sent: Thursday, 9 March 2017 10:26 AM

To: PALM

Subject: AR098906 - 7522: Exempt Clearing Notification - Springfield Rise V13 Basin & V15

Hi,

environmental management



Springfield Rise Village V13 Basin & V15 Protected Plants Flora Survey Report

Lendlease Communities
8 March 2017
7522



Document Control

Title	Springfield Rise– Village 13Basin and Village 15 (V13 Basin & V15) –Protected Plants Flora Survey Report
Job Number	7522
Client	Lendlease Communities

Document Issue

Issue	Date	Prepared By	Checked By
Draft	07.03.2017	DH	KG
Final	0.8.03.2017	DH	KG

Disclaimer

This report has been prepared for **Lendlease Communities. Saunders Havill Group** cannot accept responsibility for any use of or reliance upon the contents of this report by any third party.

Reports and/or Plans by Others

Reports and/or plans by others may be included within this report to support the document.



Table of Contents

1. Introduction	4
1.1. Key Site Details	5
2. Desktop Assessment	11
2.1. Nature Conservation Act 1992	11
2.2. Additional legislative instruments	11
3. Flora Survey Methodology	14
3.1. Clearing Impact Area	14
3.2. Survey extent and Limitations	14
3.3. Survey Timing and Limitations	14
3.4. Flora Survey Methodology	14
4. Flora Survey Results	16
4.1. Meander Transect 1	16
4.2. Summary	22
5. Appendices	23



Figures

- Figure 1: Site Context
- Figure 2: Site Aerial
- Figure 2: NCA High Risk Area Map

Plans

- Plan 1: Greater Springfield Structure Plan
- Plan 2: Springfield Rise – Development Proposal
- Plan 3: Regional Ecosystems V8
- Plan 4: Flora Meander Survey

Tables

- Table 1: Wildlife Online Search Results - Flora
- Table 2: Protected Matters Search Results - Flora
- Table 3: Transect Coordinates
- Table 4: Meander survey summary



I. Introduction

The *Environmental Management Division* of the **Saunders Havill Group** was engaged by **Lendlease Communities** to prepare this Protected Plants Flora Survey Report to enable clearing within areas mapped as 'High Risk' under the *Nature Conservation Act 1992* (NCA). Clearing works are associated with early works stages at Springfield Rise master planned development, specifically the development areas known as the Village 13 Basin and Village 15 (V13 Basin & V15). The Springfield Rise development site is located Sinnathamby Boulevard, Springfield Central (Lot 200 on SP283567 and Lot 9000 on SP288670) and is within the jurisdiction of **Ipswich City Council** (ICC).

The **Queensland Government** has adopted a risk-based approach to the regulation of protected plants under the NCA. The regulatory framework captures activities that pose a high risk to plant biodiversity. Regulatory, educational and compliance effort are consequently focused on high risk activities. Under the framework, when a non-exempt clearing activity is proposed within a 'High Risk' area, the proponent of that activity is required to complete a flora survey prior to commencement of clearing.

The main objective of the flora survey is to locate any Endangered, Vulnerable or Near Threatened (EVNT) plants that may be present within the clearing impact area. This is especially important for determining the degree of assessment required for a particular clearing activity. For example, if the survey establishes that EVNT plant species are not present within the clearing impact area, the proposed clearing will be exempt and, following notification to the department, a clearing permit will not be required. Alternatively, if EVNT plant species are identified, and clearing is considered to impact on the EVNT plant (i.e. clearing comes within 100m of the EVNT plant) then an application for a Protected Plant Clearing Permit is required.


Contextually, the Springfield Rise project site is located to the west of Springfield Central, approximately 13km southeast of Ipswich City and approximately 26 km southwest of Brisbane City. The site is bordered by commercial development and educational facilities associated with Springfield Central to the northeast, residential development to the southeast and large vegetated rural properties adjoining White Rock-Spring Mountain Conservation Estate and more broadly the Flinders-Karawatha Bioregional Corridor. The site is bound by Centenary Highway to the north and Sinnathamby Boulevard to the east. The surrounding suburbs of Redbank Plains, Springfield Lakes and Swanbank are highly urbanised and contain a mixture of residential housing, commercial properties and industrial land uses. Refer to **Figures 1 and 2** for site context and aerial. Clearing works proposed within the V13 Basin & V15 as part of early works for the commencement of the Springfield Rise project which forms part of the Greater Springfield urban development area (refer **Plan 1**). It is noted that the Springfield Rise project (refer **Plan 2**) has been approved by the Commonwealth **Department of the Environment and Energy** (DEE) (EPBC 2013/7057).

The flora surveys outlined in this report were conducted where proposed clearing is mapped within 'High Risk' areas under Protected Plants Flora Survey Trigger Mapping (refer **Figure 3**) as per the *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992*. It is noted that previously NCA protected plants surveys have been undertaken and exempt clearing notification issued by the **Department of Environment and Heritage** (EHP) for by for Villages 6, 8 and 13 and the Haul Road (AR082999) and Village 7, DA15 and DA16 (AR095633) and the Village 7 and Village 12 (AR098350).



I.I. Key Site Details

Address	Sinnathamby Boulevard, Springfield Central
RPD	Lot 200 on SP283567 and Lot 9000 on SP288670
Local Government Area	Ipswich City Council
Planning Scheme	Springfield Structure Plan, which forms part of the Ipswich City Council Planning Scheme 2003
Area Classification/Zone	Community Residential
Existing Land Use	Vacant
Proposed Land Use	Residential / Open Space



Legend

 Project impact area

Figure 1 Site Context


File ref. 7522 E Figure 1 NCA4 Site Context A
Date 9/03/2017
Project Springfield Rise Village 15

0 250 500 1,000 1,500 m

Scale (A4): 1:50,000 [GDA 1994 MGA Z56]



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend



-  Project impact area (incl. 100m buffer)
-  Approved NCA exemption areas

Figure 2 Site Aerial

File ref. 7522 E Figure 2 NCA4 Site Aerial A
Date 9/03/2017
Project Springfield Village 7 & 12


0 100 200 400 m



Scale (A4): 1:10,000 [GDA 1994 MGA Z56]



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend



-  Project impact area (incl. 100m buffer)
-  High risk area


Figure 3 NCA Flora Survey Trigger Map

File ref. 7522 E Figure 3 NCA4 NCA A
Date 9/03/2017
Project Springfield Village 7 & DA15/16

0 100 200 400 m
 Scale (A4): 1:10,000 [GDA 1994 MGA Z56]




THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend

- Project impact area (Incl. 100m buffer)
- Community residential
- Town centre
- Regional transport corridor
- Open space
- Conservation



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITIES SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAIL LOCATIONS OF ALL SERVICES.

ISSUES:

Issue	Date	Description	Drawn	Checked
A	9/03/2017	Prelim Draft	AL	MS




NCA Survey - Village 15

Greater Springfield Structure Plan

Date | 9/03/2017
 Scale | 1:32,500 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client | Lend Lease
 Project | NCA
 Address/RPD | Springfield
 Source | DCBD (DNRM, 2013), Aerial (QLD Globe, 2013)
 Layout (LandPartners 2014)


Plan 1

SHG File
 7522 E01 NCA4 Structure Plan A



Legend

- Project impact area
- RESIDENTIAL DEVELOPMENT AREA
- OPTIONAL USE AREA B
- EXISTING SCHOOL
- PROPOSED SCHOOL
- PROPOSED NEIGHBOURHOOD CENTRE
- * ACTIVITY CENTRE
- INFRASTRUCTURE (EXISTING)
- DISTRICT 2 PARK (D2P1)
- MAJOR DISTRICT PARK NORTH (FP1)
- FORMAL PARK WEST (FP2)
- LINEAR CREEKLINE OPEN SPACE
- LP LOCAL RECREATION PARK
- CENTENARY HIGHWAY
- SINNATHAMBY BOULEVARD
- TOWN CENTER ROAD TYPE 2A MODIFIED
- MAJOR COLLECTOR DUAL LANE STREET (NO ACCESS) (Area shown is where road has been defined)
- MAJOR COLLECTOR STREET (ACCESS) (Area shown is where road has been defined)
- COLLECTOR STREET
- VILLAGE BOUNDARY
- SPRING MOUNTAIN PRECINCT BOUNDARY
- 13 INDICATIVE TOWN CENTRE DEVELOPMENT AREAS (SUBJECT TO TOWN CENTRE CONCEPT PLAN)



352 ha

NOTE:
LOCATIONS AND AREAS OF VARIOUS LAND USE ELEMENTS AS SHOWN ARE CONCEPTUAL ONLY AND SUBJECT TO DETAILED DESIGN



ISSUE:	Date	Description	Drawn	Checked
1	9/03/2017	Issue	AL	MS

NCA Survey - Village 15

Springfield Rise - Development Proposal

Date: 9/03/2017
 Scale: 1:15,000 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client: Lend Lease
 Project: NCA
 Address/RPD: Springfield
 Source: Development Layout (Land Partners 2015)

Plan 2

SHG File
7522 E02NCA4 Draft Layout A



2. Desktop Assessment

2.1. Nature Conservation Act 1992

The NCA classifies and protects significant areas (Protected Areas) and protects threatened plant and animal species. The *Nature Conservation (Wildlife) Regulation 1994* (NCWR) lists plant and animal species presumed extinct, endangered, vulnerable, near threatened, least concern, international or prohibited.

The **Queensland Government** has adopted a regulatory framework that captures activities that pose a high risk to plant biodiversity. Under the framework, when a non-exempt clearing activity is proposed within a 'High Risk' area, the proponent of that activity is required to complete a flora survey prior to commencement of clearing. The Protected Plants Flora Survey Trigger Map shows 'High Risk' areas for protected plants and is used to help determine flora survey and clearing permit requirements for a particular location.

A search of the Protected Plants Flora Survey Trigger Mapping indicated proposed clearing areas within the subject site are overlaid as 'High Risk' and so are subject to flora survey requirements (refer **Figure 3**).

Prior to flora surveys, the schedules of the NCWR were considered in this report using a Wildlife Online Database Search with a 10 kilometre radius from the site. Six (6) flora species listed under the NCWR were identified as having the potential to occur on site and are presented in **Table 1**. Refer to **Appendix A** for full search results.

Table 1: Wildlife Online Search Results - Flora

Scientific Name	Common Name	Status
<i>Marsdenia coronata</i>	Slender Milk Vine	Vulnerable
<i>Plectranthus habrophyllus</i>	-	Endangered
<i>Eucalyptus curtisii</i>	Plunkett Mallee	Near Threatened
<i>Melaleuca irbyana</i>	Swamp Tea Tree	Endangered
<i>Notelaea ipsviciensis</i>	-	Endangered
<i>Notelaea lloydii</i>	Lloyd's Native Olive	Vulnerable

2.2. Additional legislative instruments

In order to maximise the scope of the flora survey, a search of protected matters listed as potentially present within 10 km of the sites under the Federal *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) was conducted using the Protected Matters Search Tool. Potential flora EVNT species listed under the EPBC Act are presented in **Table 2**. Refer to **Appendix B** for full search results.



Table 2: EPBC Act Protected Matters Search Results - Flora

Scientific Name	Common Name	Status
<i>Arthraxon hispidus</i>	Hairy Joint Grass	Vulnerable
<i>Bosistoa transversa</i>	Three-leaved Bosistoa	Vulnerable
<i>Cupaniopsis tomentella</i>	Boonah Tuckeroo	Vulnerable
<i>Dichanthium setosum</i>	Bluegrass	Vulnerable
<i>Macadamia integrifolia</i>	Macadamia Nut	Vulnerable
<i>Macadamia tetraphylla</i>	Rough-shelled Bush Nut	Vulnerable
<i>Notelaea ipsviciensis</i>	Cooneana Olive	Critically Endangered
<i>Notelaea lloydii</i>	Lloyd's Olive	Vulnerable
<i>Phaius australis</i>	Lesser Swamp-orchid	Endangered
<i>Planchonella eerwah</i>	Shiny-leaved Condoos	Endangered
<i>Plectranthus habrophyllus</i>	-	Endangered
<i>Samadera bidwillii</i>	Quassia	Vulnerable
<i>Sophora fraseri</i>	-	Vulnerable
<i>Thesium australe</i>	Austral Toadflax	Vulnerable

Regional Ecosystem mapping under the *Vegetation Management Act 1999* (VMA) was utilised to inform flora survey targets and techniques. The broader area where the survey sites occur is mapped under the VMA as Least Concern 12.9-10.17a and composite Of Concern RE12.9-10.2/12.9-10.7/12.9-10.19 (65/20/15) as described below and highlighted in **Plan 3**.

Least Concern RE 12.9 -10.17a

Description	<i>Lophostemon confertus</i> or <i>Lophostemon suaveolens</i> dominated open forest usually with emergent <i>Eucalyptus</i> and/or <i>Corymbia</i> species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments.
--------------------	--

Least Concern RE 12.9-10.2



Description	<i>Corymbia citriodora</i> subsp. <i>variegata</i> open forest or woodland usually with <i>Eucalyptus crebra</i> . Other species such as <i>Eucalyptus tereticornis</i> , <i>Eucalyptus moluccana</i> , <i>Eucalyptus acmenoides</i> and <i>Eucalyptus siderophloia</i> may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of <i>Lophostemon confertus</i> (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
--------------------	---

Of Concern RE 12.9-10.7





Description	<i>Eucalyptus crebra</i> +/- <i>Eucalyptus tereticornis</i> , <i>Corymbia tessellaris</i> , <i>Angophora leiocarpa</i> , <i>Eucalyptus melanophloia</i> woodland. Occurs on Cainozoic and Mesozoic sediments.
--------------------	---


Least Concern RE 12.9-10.19

Description	<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> woodland +/- <i>Corymbia citriodora</i> subsp. <i>variegata</i> , <i>E. acmenoides</i> or <i>E. portuensis</i> , <i>Angophora leiocarpa</i> , <i>E. major</i> . Understorey often sparse.
--------------------	---



Legend

-  Project impact area
- Regional Ecosystems v8**
-  Category A or B area containing endangered regional ecosystems
-  Category A or B area containing of concern regional ecosystems
-  Category A or B area that is a least concern regional ecosystem



APPROVED COMPANY
ISO 9001
Quality Management System
QMS

APPROVED COMPANY
ISO 14001
Environmental Management System
QMS

NCA Survey - Village 15

VMA Regional Ecosystems v8

ISSUES:

No.	Date	Description	Drawn	Checked
A	9/03/2017	Prelim Draft	AL	MS

THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.

CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METERS UNLESS OTHERWISE STATED. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.

PRIOR TO ANY DISMOUNTING, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDERGROUND SERVICES AND DETAILED LOCATIONS OF ALL SERVICES.

Date: 9/03/2017
 Scale: 1:9,000 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client: Lend Lease
 Project: NCA
 Address/RPD: Springfield
 Source: DCBD (DNRM, 2013), Aerial (QLD Globe, 2013)

Plan 3

SHG File
7522 E03 NCA4 Res A



3. Flora Survey Methodology

3.1. Clearing Impact Area

The proposed clearing site (i.e. Village 13 Bain and Village 15) is completely mapped as 'High Risk' areas under Protected Plants Flora Survey Trigger (refer **Figure 3**). The Clearing Impact Area, which is identified the area to be cleared inclusive of a 100m buffer, is shown in **Plan 4**. It is noted that previously NCA protected plants surveys have been undertaken for Villages 6, 8 and 13 and the Haul Road (AR082999), Village 7, DA15 and DA16 (AR095633), and Village 7 and Village 12 (AR098350) and a NCA exempt clearing notification confirmed by **EHP**.

3.2. Survey extent and Limitations

Table 3 and **Plan 4** summarise the Clearing Impact Area and Transect extent. Two (2) transects were undertaken over each of the V13 basin and V15 Clearing Impact Areas. It is noted that the transect over V15 was completed over 2 days due to the time constraints. These transects covered the entire area and buffer. General observations for EVNT flora species were conducted at all times while on-site, both inside and outside designated Clearing Impact Area.

Table 3: Transect Coordinates

Transect	Start	Finish
1	152.888899, -27.690898	152.888442, -27.682486
2	152.897287, 27.680524	152.890468, -27.680726


3.3. Survey Timing and Limitations

The Clearing Impact Areas include the proposed early works sites for the V13 Basin & V15. With the exception of NCA clearance, all other relevant approvals have been obtained to commence clearing works in the first half of 2017. The site has been subject to numerous ecological surveys as part of obtaining relevant Commonwealth, State and Council approvals. No EVNT species have been recorded as part of pre-clearance surveys undertaken by **SHG** as part of Commonwealth condition requirements (EPBC 2013/7075).

3.4. Flora Survey Methodology


The clearing sites were surveyed using the preferred timed meander survey technique as per *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992* by two (2) suitably qualified professionals including (1) Senior Ecologists and one (1) Ecologist (refer to **Appendix C** for curricula vitae). Surveys were carried out as follows:

- 1) The Clearing Impact Area was traversed on foot by project Ecologists (refer to **Plan 4**).
- 2) The start and finish time of each meander was recorded.
- 3) The track log of project Ecologist's transects was recorded using a handheld GPS unit accurate to < 1m.
- 4) The identity of all plant species encountered during each meander was recorded.
- 5) The site and surrounds were photographed.



Legend

- Approved NCA clearing area
- Impact area
- 100m buffer Impact boundary
- NCA MEANDER 1
- NCA MEANDER 2



NCA Survey - Villages 15

Flora Meandering Survey

ISSUES:

Issue	Date	Description	Drawn	Checked
A	9/03/2017	Prelim Draft	TC	MS

Date: 9/03/2017
 Scale: 1:7,000 @ A3
 Coordinate System: GDA 1994 MGA Zone 56
 Projection: Transverse Mercator
 Client: Lend Lease
 Project: Springfield
 Address/RPD: Springfield Village 7 & DA15/16
 Sources: QLD GIS Layers (QLD Gov. Info Services 2015), Aerial (Nearmap, 2015)

Plan 4

SHG File
 7522 E04 NCA4 Flora Meandering Survey A

THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DISMOUNTING, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDERGROUND SERVICES AND DETAILED LOCATIONS OF ALL SERVICES.



4. Flora Survey Results

The Clearing Impact Areas for the V13 Basin and V15 were assessed on the 1st and 2nd March 2017. **No EVNT species were encountered in the clearing impact area.** Given the extent of survey it can be stated with a very high level of confidence that no EVNT species will be cleared by the proposed development.

Two vegetation transects were completed with a combined length of 19, 064 meters by two ecologists using the meander methods. This method allowed for coverage of the entire investigation area and included all habitat niches observed, including rock outcrops, mapped waterways and a variety of regional ecosystem communities.

Table 4 summarises the details of each of the timed meander transect. Meander transect descriptions with photographs are presented in the following pages. A general description for the transect area is provided in this section and respective species lists in **Appendix D**.

Table 4: Meander survey summary

Transect	Date	Start Time	Finish Time	Duration	Distance	Flora Species
1(a)	1.03.2017	12:35	15:40	6 hours, 10 minutes	11088m	65
1(b)	2.03.2017	9:30	10:30	2 hour	3090m	
2	2.03.2017	8:00	9:23	2 hour, 46 minutes	4886m	72

4.1. Meander Transect I (Village I5)

Transect 1 is located within mapped remnant vegetation dominated by Least Concern regional ecosystem 12.9-10.17 and 12.9-10.19a and contains areas of composite Of Concern RE12.9-10.2/12.9-10.7/12.9-10.19. The western boundary of the survey included a mapped waterway containing Least Concern RE12.9-10.17a. These communities are described below:

- **12.9-10.17a:** *Lophostemon confertus* or *Lophostemon suaveolens* dominated open forest usually with emergent *Eucalyptus* and/or *Corymbia* species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments
- **12.9-10.19a:** *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- **12.9-10.2:** *Corymbia citriodora* subsp. *variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *E. moluccana*, *E. acmenoides* and *E. siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments
- **12.9-10.7:** *Eucalyptus crebra* +/- *E. tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *E. melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments



- **12.9-10.19:** *Eucalyptus fibrosa* subsp. *fibrosa* woodland +/- *Corymbia citriodora* subsp. *variegata*, *E. acmenoides* or *E. portuensis*, *Angophora leiocarpa*, *E. major*. Understorey often sparse. Localised occurrences of *Eucalyptus sideroxylon*. Occurs on Cainozoic and Mesozoic sediments.

The transect survey included investigations along 14,178m and included a variety of regional ecosystem communities a mapped waterway as well as environmental niches such as rock outcrops.

This transect traversed throughout the mapped remnant vegetation with the majority of the meander located on the western facing slope, the mapped waterway and the lower eastern facing slope within the same regional ecosystem communities. The mapped waterway is highly infested with weeds, particularly *Lantana camara* (Lantana) however also retains an increased density of *Lophostemon suaveolens* (Swamp Box) and a number of scattered *Eucalyptus tereticornis* (Forest Red Gum) specimens. The area containing *Lantana camara* contained very few native flora species within the ground layer.

The ridgelines and slopes within the assessment area contained a mix of *Corymbia* and *Eucalypt* species with patches of dense understorey of *Acacia* species, particularly *Acacia leiocalyx* (Early Flowering Black Wattle). These increased densities appeared to be a result of past intensive fire periods. Rock outcrops were observed along this ridgeline running north south and were targeted during the field the survey due to these areas being preferred habitat for a number of the listed flora species including *Plectranthus harbrophyllus* (Plectranthus) and *Marsdenia coronata* (Slender Milk Vine).

Sixty five (65) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation.



Photo: Transect 1 waterway areas dominated by dense *Lantana camara* (Lantana)



Photo: Transect 1 rock outcrops within the ridgeline areas observed along western facing slope.



Photo: Transect 1 included patches of dense Acacia leiocalyx as a result of past fire periods.



4.2. Meander Transect 2 (Village 13)

Transect 2 is located within mapped remnant vegetation mapped as containing a composite Of Concern regional ecosystem community containing RE12.9-10.2/12.9-10.7/12.9-10.19. The southern boundary of the survey also included the same mapped waterway as what was surveyed within Transect 1 however further downstream, and is mapped as containing Least Concern RE12.9-10.17a. Vegetation recorded throughout the survey period appeared to be consistent with the current regional ecosystem mapping. These communities are described below:

- **12.9-10.17a:** *Lophostemon confertus* or *Lophostemon suaveolens* dominated open forest usually with emergent *Eucalyptus* and/or *Corymbia* species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments
- **12.9-10.19a:** *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- **12.9-10.2:** *Corymbia citriodora* subsp. *variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *E. moluccana*, *E. acmenoides* and *E. siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments
- **12.9-10.7:** *Eucalyptus crebra* +/- *E. tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *E. melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments
- **12.9-10.19:** *Eucalyptus fibrosa* subsp. *fibrosa* woodland +/- *Corymbia citriodora* subsp. *variegata*, *E. acmenoides* or *E. portuensis*, *Angophora leiocarpa*, *E. major*. Understorey often sparse. Localised occurrences of *Eucalyptus sideroxylon*. Occurs on Cainozoic and Mesozoic sediments.

The transect survey included investigations along 4886 meters by two experienced ecologists.

This transect traversed throughout the mapped remnant vegetation with the majority of the meander located on the southern facing slope and the mapped waterway. Directly south of the investigation area has recently been cleared as part of the broader Springfield development. The construction of Sinnathamby Blvd has also resulted in the inundation of a portion of the investigation area. This is evident with a large number of dead Eucalypt stags still standing within the inundated area. Overland flow paths and drainage features running down the western facing slope are highly infested with introduced species, particularly *Lantana camara* (Lantana).

Vegetation throughout the whole polygon is dominated by Eucalypt and *Corymbia* species with weeds largely confined to the shrub and ground layers. A small area within the central portion of the polygon contains exposed rocky outcrop, with particular attention paid to the listed flora species *Plectranthus harbrophyllus* (*Plectranthus*) and *Marsdenia coronata* (Slender Milk Vine).

The majority of the assessment area contained a mix of *Corymbia* and Eucalypt species and is described as lower slopes. No ridge lines occur within this investigation transect. Small patches of thick *Acacia* including *Acacia leiocalyx* (Early Flowering Black Wattle) and *Acacia fimbriata* (Fringed Wattle) were observed throughout the transect, however, the majority of the polygon contained relatively sparse understorey.



Seventy two (72) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation.



Photo: Transect 2 waterway areas dominated by dense *Lantana camara* (Lantana)



Photo: Transect 2 rocky outcrops within the ridgeline areas.



Photo: Transect 2 Steep south facing slope.



4.3. Summary

Field surveys were carried out within the clearing impact area and buffer of early works precincts (Village 13 Basin and Village15) of the Springfield Rise project site which is mapped as 'High Risk' by Protected Plants Flora Survey Trigger Mapping. The survey utilised the preferred random meander technique as outlined in the *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992* to identify the presence of EVNT species. Coverage included the proposed clearing extents as well as a 100 m buffer. A single meander transect as well as continual observations were completed throughout the investigation area. It is noted that access was heavily impeded by dense Lantana infestations.

The following points provide a summary of the investigation area:

- The vegetation communities observed have been extensively searched and analysed against current regional ecosystem mapping with overall consistencies in the location of each regional ecosystem community. Some minor variations were observed however in the majority of areas these variations are too small to provide for changes to this mapping.
- The majority of the canopy layer of the Clearing Impact Area reflects relatively intact representing an open forest to woodland community. Although evidence of forestry practices was recorded in all transects and throughout observational survey points, the site remains as remnant due to the vegetation community's height and density.
- The sub-canopy layer is relatively sparse throughout the majority of the site and is typical of the mapped vegetation communities represented on site.
- The understorey layer is variable with some patches along ridgelines containing dense acacia regrowth and the VMA mapped waterway containing a relatively sparse native flora layer. The dense infestations of acacia indicate evidence of historical fire events. Old cut stumps throughout the area also indicate historical logging practices.
- Dense weed infestation was observed throughout the mapped waterways and drainage lines. Low to moderate infestation were observed on the lower slopes of the ridgelines with only sparse infestations on the higher part of the ridges.
- Exposed rocky habitat was recorded in isolated patches along ridgelines. Although these areas have been extensively searched, no threatened species were recorded at the time of the assessment within the investigation area.

Surveys **did not identify any EVNT species within the proposed clearing areas or the 100m buffer..**



5. Appendices

Appendix A

Wildlife Online Search Results

Appendix B

Protected Matters Search Results

Appendix C

Curricula Vitae

Appendix D

Species Lists



Appendix B

Wildlife Online Search Results



Queensland Government

Wildlife Online Extract

Search Criteria: Species List for a Specified Point
Species: All
Type: All
Status: Rare and threatened species
Records: All
Date: All
Latitude: -27.6884
Longitude: 152.8983
Distance: 10
Email: keiragrundy@saundershavill.com
Date submitted: Tuesday 07 Mar 2017 17:21:25
Date extracted: Tuesday 07 Mar 2017 17:30:03

The number of records retrieved = 18

Disclaimer

As the DSITIA is still in a process of collating and vetting data, it is possible the information given is not complete. The information provided should only be used for the project for which it was requested and it should be appropriately acknowledged as being derived from Wildlife Online when it is used.

The State of Queensland does not invite reliance upon, nor accept responsibility for this information. Persons should satisfy themselves through independent means as to the accuracy and completeness of this information.

No statements, representations or warranties are made about the accuracy or completeness of this information. The State of Queensland disclaims all responsibility for this information and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you may incur as a result of the information being inaccurate or incomplete in any way for any reason.

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	amphibians	Limnodynastidae	<i>Adelotus brevis</i>	tusked frog		V		10
animals	birds	Cacatuidae	<i>Calyptorhynchus lathami lathami</i>	glossy black-cockatoo (eastern)		V		9
animals	birds	Falconidae	<i>Falco hypoleucos</i>	grey falcon		V		1
animals	birds	Psittacidae	<i>Lathamus discolor</i>	swift parrot		E	CE	3
animals	birds	Rostratulidae	<i>Rostratula australis</i>	Australian painted snipe		V	E	8
animals	birds	Strigidae	<i>Ninox strenua</i>	powerful owl		V		13
animals	birds	Turnicidae	<i>Turnix melanogaster</i>	black-breasted button-quail		V	V	1
animals	mammals	Dasyuridae	<i>Dasyurus maculatus maculatus</i>	spotted-tailed quoll (southern subspecies)		V	E	3
animals	mammals	Macropodidae	<i>Petrogale penicillata</i>	brush-tailed rock-wallaby		V	V	8
animals	mammals	Phascolarctidae	<i>Phascolarctos cinereus</i>	koala		V	V	554
animals	mammals	Vombatidae	<i>Vombatus ursinus</i>	common wombat		NT		1
animals	reptiles	Elapidae	<i>Acanthophis antarcticus</i>	common death adder		V		1
plants	higher dicots	Apocynaceae	<i>Marsdenia coronata</i>	slender milkvine		V		19/19
plants	higher dicots	Lamiaceae	<i>Plectranthus habrophyllus</i>			E	E	16/16
plants	higher dicots	Myrtaceae	<i>Eucalyptus curtisii</i>	Plunkett mallee		NT		13/13
plants	higher dicots	Myrtaceae	<i>Melaleuca irbyana</i>			E		3/3
plants	higher dicots	Oleaceae	<i>Notelaea ipsviciensis</i>			E	CE	10/10
plants	higher dicots	Oleaceae	<i>Notelaea lloydii</i>	Lloyd's native olive		V	V	7/7

CODES

I - Y indicates that the taxon is introduced to Queensland and has naturalised.

Q - Indicates the Queensland conservation status of each taxon under the *Nature Conservation Act 1992*. The codes are Extinct in the Wild (PE), Endangered (E), Vulnerable (V), Near Threatened (NT), Least Concern (C) or Not Protected ().

A - Indicates the Australian conservation status of each taxon under the *Environment Protection and Biodiversity Conservation Act 1999*. The values of EPBC are Conservation Dependent (CD), Critically Endangered (CE), Endangered (E), Extinct (EX), Extinct in the Wild (XW) and Vulnerable (V).

Records – The first number indicates the total number of records of the taxon for the record option selected (i.e. All, Confirmed or Specimens).

This number is output as 99999 if it equals or exceeds this value. The second number located after the / indicates the number of specimen records for the taxon.

This number is output as 999 if it equals or exceeds this value.



Appendix C

Protected Matters Search Results



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 07/03/17 18:31:36

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)


[Acknowledgements](#)



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

[Coordinates](#)

Buffer: 10.0Km





Queensland Government

Wildlife Online Extract

Search Criteria: Species List for a Specified Point
Species: All
Type: All
Status: Rare and threatened species
Records: All
Date: All
Latitude: -27.6884
Longitude: 152.8983
Distance: 10
Email: keiragrundy@saundershavill.com
Date submitted: Tuesday 07 Mar 2017 17:21:25
Date extracted: Tuesday 07 Mar 2017 17:30:03

The number of records retrieved = 18

Disclaimer

As the DSITIA is still in a process of collating and vetting data, it is possible the information given is not complete. The information provided should only be used for the project for which it was requested and it should be appropriately acknowledged as being derived from Wildlife Online when it is used.

The State of Queensland does not invite reliance upon, nor accept responsibility for this information. Persons should satisfy themselves through independent means as to the accuracy and completeness of this information.

No statements, representations or warranties are made about the accuracy or completeness of this information. The State of Queensland disclaims all responsibility for this information and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you may incur as a result of the information being inaccurate or incomplete in any way for any reason.

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	amphibians	Limnodynastidae	<i>Adelotus brevis</i>	tusked frog		V		10
animals	birds	Cacatuidae	<i>Calyptorhynchus lathami lathami</i>	glossy black-cockatoo (eastern)		V		9
animals	birds	Falconidae	<i>Falco hypoleucos</i>	grey falcon		V		1
animals	birds	Psittacidae	<i>Lathamus discolor</i>	swift parrot		E	CE	3
animals	birds	Rostratulidae	<i>Rostratula australis</i>	Australian painted snipe		V	E	8
animals	birds	Strigidae	<i>Ninox strenua</i>	powerful owl		V		13
animals	birds	Turnicidae	<i>Turnix melanogaster</i>	black-breasted button-quail		V	V	1
animals	mammals	Dasyuridae	<i>Dasyurus maculatus maculatus</i>	spotted-tailed quoll (southern subspecies)		V	E	3
animals	mammals	Macropodidae	<i>Petrogale penicillata</i>	brush-tailed rock-wallaby		V	V	8
animals	mammals	Phascolarctidae	<i>Phascolarctos cinereus</i>	koala		V	V	554
animals	mammals	Vombatidae	<i>Vombatus ursinus</i>	common wombat		NT		1
animals	reptiles	Elapidae	<i>Acanthophis antarcticus</i>	common death adder		V		1
plants	higher dicots	Apocynaceae	<i>Marsdenia coronata</i>	slender milkvine		V		19/19
plants	higher dicots	Lamiaceae	<i>Plectranthus habrophyllus</i>			E	E	16/16
plants	higher dicots	Myrtaceae	<i>Eucalyptus curtisii</i>	Plunkett mallee		NT		13/13
plants	higher dicots	Myrtaceae	<i>Melaleuca irbyana</i>			E		3/3
plants	higher dicots	Oleaceae	<i>Notelaea ipsviciensis</i>			E	CE	10/10
plants	higher dicots	Oleaceae	<i>Notelaea lloydii</i>	Lloyd's native olive		V	V	7/7

CODES

I - Y indicates that the taxon is introduced to Queensland and has naturalised.

Q - Indicates the Queensland conservation status of each taxon under the *Nature Conservation Act 1992*. The codes are Extinct in the Wild (PE), Endangered (E), Vulnerable (V), Near Threatened (NT), Least Concern (C) or Not Protected ().

A - Indicates the Australian conservation status of each taxon under the *Environment Protection and Biodiversity Conservation Act 1999*. The values of EPBC are Conservation Dependent (CD), Critically Endangered (CE), Endangered (E), Extinct (EX), Extinct in the Wild (XW) and Vulnerable (V).

Records – The first number indicates the total number of records of the taxon for the record option selected (i.e. All, Confirmed or Specimens).

This number is output as 99999 if it equals or exceeds this value. The second number located after the / indicates the number of specimen records for the taxon.

This number is output as 999 if it equals or exceeds this value.

Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	None
Wetlands of International Importance:	1
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	None
Listed Threatened Ecological Communities:	2
Listed Threatened Species:	59
Listed Migratory Species:	27

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	1
Commonwealth Heritage Places:	1
Listed Marine Species:	38
Whales and Other Cetaceans:	1
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Commonwealth Reserves Marine:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	2
Regional Forest Agreements:	None
Invasive Species:	42
Nationally Important Wetlands:	1
Key Ecological Features (Marine)	None

Details

Matters of National Environmental Significance

Wetlands of International Importance (Ramsar)	[Resource Information]
Name	Proximity
Moreton bay	20 - 30km upstream

Listed Threatened Ecological Communities

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Name	Status	Type of Presence
Lowland Rainforest of Subtropical Australia	Critically Endangered	Community may occur within area
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland	Critically Endangered	Community likely to occur within area

Listed Threatened Species

Name	Status	Type of Presence
------	--------	------------------

Birds		
Anthochaera phrygia Regent Honeyeater [82338]	Critically Endangered	Foraging, feeding or related behaviour may occur within area
Botaurus poiciloptilus Australasian Bittern [1001]	Endangered	Species or species habitat likely to occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Cyclopsitta diophthalma coxeni Coxen's Fig-Parrot [59714]	Endangered	Species or species habitat may occur within area
Dasyornis brachypterus Eastern Bristlebird [533]	Endangered	Species or species habitat likely to occur within area
Diomedea antipodensis Antipodean Albatross [64458]	Vulnerable	Species or species habitat may occur within area
Diomedea antipodensis gibsoni Gibson's Albatross [82270]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
Erythrotriorchis radiatus Red Goshawk [942]	Vulnerable	Species or species habitat likely to occur within area
Geophaps scripta scripta Squatter Pigeon (southern) [64440]	Vulnerable	Species or species

Name	Status	Type of Presence
Grantiella picta Painted Honeyeater [470]	Vulnerable	habitat may occur within area Species or species habitat may occur within area
Lathamus discolor Swift Parrot [744]	Critically Endangered	Species or species habitat likely to occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pachyptila turtur subantarctica Fairy Prion (southern) [64445]	Vulnerable	Species or species habitat likely to occur within area
Poephila cincta cincta Southern Black-throated Finch [64447]	Endangered	Species or species habitat may occur within area
Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat likely to occur within area
Thalassarche cauta cauta Shy Albatross, Tasmanian Shy Albatross [82345]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta steadi White-capped Albatross [82344]	Vulnerable	Species or species habitat likely to occur within area
Thalassarche eremita Chatham Albatross [64457]	Endangered	Species or species habitat may occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche salvini Salvin's Albatross [64463]	Vulnerable	Species or species habitat may occur within area
Turnix melanogaster Black-breasted Button-quail [923]	Vulnerable	Species or species habitat likely to occur within area
Fish		
Epinephelus daemeli Black Rockcod, Black Cod, Saddled Rockcod [68449]	Vulnerable	Species or species habitat may occur within area
Insects		
Phyllodes imperialis smithersi Pink Underwing Moth [86084]	Endangered	Species or species habitat may occur within area
Mammals		

Name	Status	Type of Presence
Chalinolobus dwyeri Large-eared Pied Bat, Large Pied Bat [183]	Vulnerable	Species or species habitat likely to occur within area
Dasyurus hallucatus Northern Quoll, Digul [331]	Endangered	Species or species habitat may occur within area
Dasyurus maculatus maculatus (SE mainland population) Spot-tailed Quoll, Spotted-tail Quoll, Tiger Quoll (southeastern mainland population) [75184]	Endangered	Species or species habitat known to occur within area
Petauroides volans Greater Glider [254]	Vulnerable	Species or species habitat known to occur within area
Petrogale penicillata Brush-tailed Rock-wallaby [225]	Vulnerable	Species or species habitat known to occur within area
Phascolarctos cinereus (combined populations of Qld, NSW and the ACT) Koala (combined populations of Queensland, New South Wales and the Australian Capital Territory) [85104]	Vulnerable	Species or species habitat known to occur within area
Potorous tridactylus tridactylus Long-nosed Potoroo (SE mainland) [66645]	Vulnerable	Species or species habitat may occur within area
Pteropus poliocephalus Grey-headed Flying-fox [186]	Vulnerable	Roosting known to occur within area
Other		
Cycas ophiolitica [55797]	Endangered	Species or species habitat likely to occur within area
Plants		
Arthraxon hispidus Hairy-joint Grass [9338]	Vulnerable	Species or species habitat may occur within area
Bosistoa transversa Three-leaved Bosistoa, Yellow Satinheart [16091]	Vulnerable	Species or species habitat likely to occur within area
Cupaniopsis tomentella Boonah Tuckeroo [3322]	Vulnerable	Species or species habitat likely to occur within area
Dichanthium setosum bluegrass [14159]	Vulnerable	Species or species habitat likely to occur within area
Macadamia integrifolia Macadamia Nut, Queensland Nut Tree, Smooth-shelled Macadamia, Bush Nut, Nut Oak [7326]	Vulnerable	Species or species habitat likely to occur within area
Macadamia tetraphylla Rough-shelled Bush Nut, Macadamia Nut, Rough-shelled Macadamia, Rough-leaved Queensland Nut [6581]	Vulnerable	Species or species habitat may occur within area
Notelaea ipsviciensis Cooneana Olive [81858]	Critically Endangered	Species or species habitat likely to occur within area
Notelaea lloydii Lloyd's Olive [15002]	Vulnerable	Species or species habitat likely to occur within area
Phaius australis Lesser Swamp-orchid [5872]	Endangered	Species or species habitat likely to occur within area

Name	Status	Type of Presence
Planchonella eerwah Shiny-leaved Condoe, Black Plum, Wild Apple [17340]	Endangered	Species or species habitat likely to occur within area
Plectranthus habrophyllus [64589]	Endangered	Species or species habitat likely to occur within area
Samadera bidwillii Quassia [29708]	Vulnerable	Species or species habitat likely to occur within area
Sophora fraseri [8836]	Vulnerable	Species or species habitat likely to occur within area
Thesium australe Austral Toadflax, Toadflax [15202]	Vulnerable	Species or species habitat likely to occur within area

Reptiles

Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Delma torquata Adorned Delma, Collared Delma [1656]	Vulnerable	Species or species habitat likely to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Furina dunmalli Dunmall's Snake [59254]	Vulnerable	Species or species habitat may occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area
Saiphos reticulatus Three-toed Snake-tooth Skink [88328]	Vulnerable	Species or species habitat may occur within area

Listed Migratory Species

[[Resource Information](#)]

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Migratory Marine Birds		
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within

Name	Threatened	Type of Presence area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta Tasmanian Shy Albatross [89224]	Vulnerable*	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Migratory Marine Species		
Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Manta alfredi Reef Manta Ray, Coastal Manta Ray, Inshore Manta Ray, Prince Alfred's Ray, Resident Manta Ray [84994]		Species or species habitat may occur within area
Manta birostris Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray, Oceanic Manta Ray [84995]		Species or species habitat may occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area
Orcaella brevirostris Irrawaddy Dolphin [45]		Species or species habitat known to occur within area
Migratory Terrestrial Species		
Cuculus optatus Oriental Cuckoo, Horsfield's Cuckoo [86651]		Species or species habitat may occur within area
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat known to occur within area
Monarcha melanopsis Black-faced Monarch [609]		Species or species habitat known to occur within area
Monarcha trivirgatus Spectacled Monarch [610]		Species or species habitat known to occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species

Name	Threatened	Type of Presence
Rhipidura rufifrons Rufous Fantail [592]		habitat known to occur within area Species or species habitat known to occur within area
Migratory Wetlands Species		
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Gallinago hardwickii Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Other Matters Protected by the EPBC Act

Commonwealth Land [\[Resource Information \]](#)

The Commonwealth area listed below may indicate the presence of Commonwealth land in this vicinity. Due to the unreliability of the data source, all proposals should be checked as to whether it impacts on a Commonwealth area, before making a definitive decision. Contact the State or Territory government land department for further information.

Name
Defence - GREENBANK TRAINING AREA

Commonwealth Heritage Places [\[Resource Information \]](#)

Name	State	Status
Natural		
Greenbank Military Training Area (part)	QLD	Listed place

Listed Marine Species [\[Resource Information \]](#)

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Birds		
Anseranas semipalmata Magpie Goose [978]		Species or species habitat may occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba Great Egret, White Egret [59541]		Breeding known to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Cuculus saturatus Oriental Cuckoo, Himalayan Cuckoo [710]		Species or species habitat may occur within area

Name	Threatened	Type of Presence
Diomedea antipodensis Antipodean Albatross [64458]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans Wandering Albatross [89223]	Vulnerable	Species or species habitat may occur within area
Diomedea gibsoni Gibson's Albatross [64466]	Vulnerable*	Species or species habitat may occur within area
Gallinago hardwickii Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area
Haliaeetus leucogaster White-bellied Sea-Eagle [943]		Species or species habitat known to occur within area
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat known to occur within area
Lathamus discolor Swift Parrot [744]	Critically Endangered	Species or species habitat likely to occur within area
Macronectes giganteus Southern Giant-Petrel, Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Monarcha melanopsis Black-faced Monarch [609]		Species or species habitat known to occur within area
Monarcha trivirgatus Spectacled Monarch [610]		Species or species habitat known to occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species habitat known to occur within area
Numenius madagascariensis Eastern Curlew, Far Eastern Curlew [847]	Critically Endangered	Species or species habitat likely to occur within area
Pachyptila turtur Fairy Prion [1066]		Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area
Rhipidura rufifrons Rufous Fantail [592]		Species or species habitat known to occur within area

Name	Threatened	Type of Presence
Rostratula benghalensis (sensu lato) Painted Snipe [889]	Endangered*	Species or species habitat likely to occur within area
Thalassarche cauta Tasmanian Shy Albatross [89224]	Vulnerable*	Species or species habitat may occur within area
Thalassarche eremita Chatham Albatross [64457]	Endangered	Species or species habitat may occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche salvini Salvin's Albatross [64463]	Vulnerable	Species or species habitat may occur within area
Thalassarche steadi White-capped Albatross [64462]	Vulnerable*	Species or species habitat likely to occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Reptiles

Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area

Whales and other Cetaceans

[[Resource Information](#)]

Name	Status	Type of Presence
Mammals		
Orcaella brevirostris Irrawaddy Dolphin [45]		Species or species habitat known to occur within area

Extra Information

State and Territory Reserves [\[Resource Information \]](#)

Name	State
Stewartdale	QLD
White Rock	QLD

Invasive Species [\[Resource Information \]](#)

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

Name	Status	Type of Presence
------	--------	------------------

Birds

Acridotheres tristis Common Myna, Indian Myna [387]		Species or species habitat likely to occur within area
--	--	--

Anas platyrhynchos Mallard [974]		Species or species habitat likely to occur within area
-------------------------------------	--	--

Carduelis carduelis European Goldfinch [403]		Species or species habitat likely to occur within area
---	--	--

Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
--	--	--

Lonchura punctulata Nutmeg Mannikin [399]		Species or species habitat likely to occur within area
--	--	--

Passer domesticus House Sparrow [405]		Species or species habitat likely to occur within area
--	--	--

Streptopelia chinensis Spotted Turtle-Dove [780]		Species or species habitat likely to occur within area
---	--	--

Sturnus vulgaris Common Starling [389]		Species or species habitat likely to occur within area
---	--	--

Frogs

Rhinella marina Cane Toad [83218]		Species or species habitat likely to occur within area
--------------------------------------	--	--

Mammals

Bos taurus Domestic Cattle [16]		Species or species habitat likely to occur within area
------------------------------------	--	--

Canis lupus familiaris Domestic Dog [82654]		Species or species habitat likely to occur within area
--	--	--

Equus caballus Horse [5]		Species or species habitat likely to occur within area
-----------------------------	--	--

Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
--	--	--

Name	Status	Type of Presence
Feral deer Feral deer species in Australia [85733]		Species or species habitat likely to occur within area
Lepus capensis Brown Hare [127]		Species or species habitat likely to occur within area
Mus musculus House Mouse [120]		Species or species habitat likely to occur within area
Oryctolagus cuniculus Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
Rattus norvegicus Brown Rat, Norway Rat [83]		Species or species habitat likely to occur within area
Rattus rattus Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
Sus scrofa Pig [6]		Species or species habitat likely to occur within area
Vulpes vulpes Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
Anredera cordifolia Madeira Vine, Jalap, Lamb's-tail, Mignonette Vine, Anredera, Gulf Madeiravine, Heartleaf Madeiravine, Potato Vine [2643]		Species or species habitat likely to occur within area
Asparagus aethiopicus Asparagus Fern, Ground Asparagus, Basket Fern, Sprengi's Fern, Bushy Asparagus, Emerald Asparagus [62425]		Species or species habitat likely to occur within area
Asparagus africanus Climbing Asparagus, Climbing Asparagus Fern [66907]		Species or species habitat likely to occur within area
Asparagus plumosus Climbing Asparagus-fern [48993]		Species or species habitat likely to occur within area
Cabomba caroliniana Cabomba, Fanwort, Carolina Watershield, Fish Grass, Washington Grass, Watershield, Carolina Fanwort, Common Cabomba [5171]		Species or species habitat likely to occur within area
Chrysanthemoides monilifera Bitou Bush, Boneseed [18983]		Species or species habitat may occur within area
Dolichandra unguis-cati Cat's Claw Vine, Yellow Trumpet Vine, Cat's Claw Creeper, Funnel Creeper [85119]		Species or species habitat likely to occur within area
Eichhornia crassipes Water Hyacinth, Water Orchid, Nile Lily [13466]		Species or species habitat likely to occur within area
Genista monspessulana Montpellier Broom, Cape Broom, Canary Broom, Common Broom, French Broom, Soft Broom [20126]		Species or species habitat likely to occur within area
Lantana camara Lantana, Common Lantana, Kamara Lantana, Large-leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White		Species or species habitat likely to occur within area

Name	Status	Type of Presence
Sage, Wild Sage [10892] Opuntia spp. Prickly Pears [82753]		Species or species habitat likely to occur within area
Parkinsonia aculeata Parkinsonia, Jerusalem Thorn, Jelly Bean Tree, Horse Bean [12301]		Species or species habitat likely to occur within area
Parthenium hysterophorus Parthenium Weed, Bitter Weed, Carrot Grass, False Ragweed [19566]		Species or species habitat likely to occur within area
Protasparagus densiflorus Asparagus Fern, Plume Asparagus [5015]		Species or species habitat likely to occur within area
Protasparagus plumosus Climbing Asparagus-fern, Ferny Asparagus [11747]		Species or species habitat likely to occur within area
Sagittaria platyphylla Delta Arrowhead, Arrowhead, Slender Arrowhead [68483]		Species or species habitat likely to occur within area
Salix spp. except S.babylonica, S.x calodendron & S.x reichardtii Willows except Weeping Willow, Pussy Willow and Sterile Pussy Willow [68497]		Species or species habitat likely to occur within area
Salvinia molesta Salvinia, Giant Salvinia, Aquarium Watermoss, Kariba Weed [13665]		Species or species habitat likely to occur within area
Senecio madagascariensis Fireweed, Madagascar Ragwort, Madagascar Groundsel [2624]		Species or species habitat likely to occur within area
Solanum elaeagnifolium Silver Nightshade, Silver-leaved Nightshade, White Horse Nettle, Silver-leaf Nightshade, Tomato Weed, White Nightshade, Bull-nettle, Prairie-berry, Satansbos, Silver-leaf Bitter-apple, Silverleaf-nettle, Trompillo [12323]		Species or species habitat likely to occur within area
Reptiles		
Hemidactylus frenatus Asian House Gecko [1708]		Species or species habitat likely to occur within area
Nationally Important Wetlands		[Resource Information]
Name		State
Greenbank Army Training Area C		QLD

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-27.68849 152.8985

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [-Office of Environment and Heritage, New South Wales](#)
- [-Department of Environment and Primary Industries, Victoria](#)
- [-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [-Department of Environment, Water and Natural Resources, South Australia](#)
- [-Department of Land and Resource Management, Northern Territory](#)
- [-Department of Environmental and Heritage Protection, Queensland](#)
- [-Department of Parks and Wildlife, Western Australia](#)
- [-Environment and Planning Directorate, ACT](#)
- [-Birdlife Australia](#)
- [-Australian Bird and Bat Banding Scheme](#)
- [-Australian National Wildlife Collection](#)
- [-Natural history museums of Australia](#)
- [-Museum Victoria](#)
- [-Australian Museum](#)
- [-South Australian Museum](#)
- [-Queensland Museum](#)
- [-Online Zoological Collections of Australian Museums](#)
- [-Queensland Herbarium](#)
- [-National Herbarium of NSW](#)
- [-Royal Botanic Gardens and National Herbarium of Victoria](#)
- [-Tasmanian Herbarium](#)
- [-State Herbarium of South Australia](#)
- [-Northern Territory Herbarium](#)
- [-Western Australian Herbarium](#)
- [-Australian National Herbarium, Canberra](#)
- [-University of New England](#)
- [-Ocean Biogeographic Information System](#)
- [-Australian Government, Department of Defence Forestry Corporation, NSW](#)
- [-Geoscience Australia](#)
- [-CSIRO](#)
- [-Australian Tropical Herbarium, Cairns](#)
- [-eBird Australia](#)
- [-Australian Government – Australian Antarctic Data Centre](#)
- [-Museum and Art Gallery of the Northern Territory](#)
- [-Australian Government National Environmental Science Program](#)
- [-Australian Institute of Marine Science](#)
- [-Reef Life Survey Australia](#)
- [-American Museum of Natural History](#)
- [-Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [-Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- [-Other groups and individuals](#)

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.



Appendix D

Curricula Vitae – Pen Port



Andrew Craig: Senior Ecologist

Andrew is a senior field ecologist with significant practical experience in the areas of ecological site assessment, weed management programs, large scale revegetation projects, wetland rehabilitation and waterway restoration. His main area of expertise is the identification and classification of flora and fauna including the identification and management of threatened species and communities. Andrew has significant experience in some of Queensland's largest infrastructure projects including coordinating on-ground flora assessments and development of weed management and rehabilitation strategies for the Southern Regional Water Pipeline.




Andrew's background in managing revegetation, translocation and forestry establishment projects brings a wealth of experience in the practical management, rehabilitation and offsetting across numerous projects. These skills linked with strong scientific and analytical site survey methods ensures Saunders Havill Group complies with all necessary state and federal government sampling procedures.

Qualifications

Bachelor of Science (Zoology), [The University of Queensland](#) (1997)

Maree Clancy: Ecologist

Maree has extensive ecological field and desktop research experience gained while working in the forestry industry and with the Australian Koala Foundation. In previous roles she assisted with quarterly and annual reporting of rehabilitation/revegetation works at residential development reserves, habitat translocation sites and the Bruce Highway upgrade project, and also with annual fauna surveying and reporting on various projects. She has a wealth of experience with preliminary desktop assessments of potential species at survey sites and the identification of flora and fauna species present during surveys.



At the Australian Koala Foundation, Maree was involved in the Koala habitat mapping project which included the use of GIS and determining habitat values for regional ecosystems and mosaics based on canopy species rankings and percentage composition.

Maree has additional skills in native seed propagation and growing of seedlings for large scale revegetation and farm forestry projects, ongoing monitoring of propagation methods and plant health status and adaptive approaches to improving methods.

Qualifications

Bachelor of Environmental Science, [University of the Sunshine Coast](#) (2014)



Appendix E

Species Lists



Flora Species Transect 1 (Village 15)	
Species	Common Name
<i>Acacia concurrens</i>	Black Wattle
<i>Acacia disparrima</i>	Hickory Wattle
<i>Acacia fimbriata</i>	Fringed Wattle
<i>Acacia leiocalyx</i>	Early Flowering Black Wattle
<i>Adiantum atroviride</i>	Maiden Hair Fern
<i>Ageratum houstonianum</i>	Blue Billy Goat Weed
<i>Allocasuarina littoralis</i>	Black She Oak
<i>Alphitonia excelsa</i>	Soap Tree
<i>Angophora leiocarpa</i>	Smooth Bark Apple
<i>Aristida latifolia</i>	Many Head Wire Grass
<i>Celtis sensis</i>	Chinese Elm
<i>Chloris gayana</i>	Rhodes Grass
<i>Chrysocephalum apiculatum</i>	Yellow Buttons
<i>Conyza sumatrensis</i>	Tall Fleabane
<i>Corymbia citriodora</i>	Spotted Gum
<i>Corymbia intermedia</i>	Pink Bloodwood
<i>Corymbia tessellaris</i>	Moreton Bay Ash
<i>Corymbia torrelliana</i>	Cadaghi
<i>Corymbia trachyphloia</i>	Brown Bloodwood
<i>Cuscuta campestris</i>	Dodder Vine
<i>Cymbopogon refractus</i>	Barbed Wire Grass
<i>Dianella careulea</i>	Blue Lilly
<i>Dichondra repens</i>	Kydney Weed
<i>Dodonaea triquetra</i>	Forest Hop Bush
<i>Dodonaea viscosa</i>	Hop Bush
<i>Eragrostis brownii</i>	Browns Love Grass
<i>Eremophila debilis</i>	Winter Apple
<i>Eucalyptus acmenoides</i>	White Mahogany
<i>Eucalyptus fibrosa</i>	Broad Leaf Ironbark
<i>Eucalyptus microcorys</i>	Tallowwood
<i>Eucalyptus seeana</i>	Narrow Leaf Red Gum
<i>Eucalyptus siderophloia</i>	Grey Ironbark
<i>Eucalyptus tereticornis</i>	Forest Red Gum



<i>Eustrephus latifolius</i>	Wombat Berry
<i>Gahnia asperia</i>	Saw Sedge
<i>Glochidion ferdinandi</i>	Cheese Tree
<i>Glossocardia bidens</i>	Native Cobblers Pegs
<i>Goodenia rotundifolia</i>	Goodenia
<i>Grewia latifolia</i>	Dogs Balls
<i>Hardenbergia violacea</i>	Native Sarsparilla
<i>Hetropogon conturtus</i>	Black Spear Grass
<i>Imperata cylindrica</i>	Blady Grass
<i>Jacksonia scoparia</i>	Dogwood
<i>Lantana camara</i>	Lantana
<i>Lantana montevidensis</i>	Creeping Lantana
<i>Lepidosperma</i>	Lemon Scented Myrtle
<i>Leucopogon juniperinus</i>	Prickly Heath
<i>Lobelia purpurascens</i>	White Root
<i>Lomandra longifolia</i>	Mat Rush
<i>Lomandra multiflora</i>	Many Flowered Mat Rush
<i>Lophostemon confertus</i>	Brush Box
<i>Lophostemon suaveolens</i>	Swamp Box
<i>Melinis repens</i>	Red Natal Grass
<i>Neonotonia wightii</i>	Glycine
<i>Nephrolepis</i>	Fish Bone Fern
<i>Passiflora suberosa</i>	Corky Passion Vine
<i>Petalostigma pubescens</i>	Quinine Bush
<i>Plectranthus parviflorus</i>	Plectranthus
<i>Pteridium esculentum</i>	Bracken Fern
<i>Setaria sphacelata</i>	Setaria
<i>Sida cordifolia</i>	Flannel Weed
<i>Smilax australis</i>	Barbed Wire Vine
<i>Themeda triandra</i>	Kangaroo Grass
<i>Vigna vexillata</i>	Wild Cowpea
<i>Xanthorrhoea glauca</i>	Grass Tree



Flora Species Transect 2 (Village 13)	
Species	Common Name
<i>Acacia fimbriata</i>	Fringed Wattle
<i>Acacia leiocalyx</i>	Early Flowering Black Wattle
<i>Ageratum houstonianum</i>	Blue Billygoat Weed
<i>Alphitonia excelsa</i>	Soap Tree
<i>Angophora subvelintina</i>	Rough Barked Apple
<i>Aristida latifolia</i>	Many Flowering Wire Grass
<i>Baccharis halimifolia</i>	Groundsel
<i>Bidens pilosa</i>	Cobbler's Pegs
<i>Brenna oblongifolia</i>	Coffee Bush
<i>Carex lasiocarpa</i>	Slender Sedge
<i>Cheilanthes distans</i>	Bristle Cloak Fern
<i>Chloris gayana</i>	Rhodes Grass
<i>Cirsium vulgare</i>	Spear Thistle
<i>Conyza sumatrensis</i>	Tall Fleabane
<i>Corymbia citriodora</i>	Spotted Gum
<i>Corymbia torrelliana</i>	Cadaghi
<i>Cuscuta campestris</i>	Dodder Vine
<i>Cymbopogon refractus</i>	Barbed Wire Grass
<i>Cyperus polystachyos</i>	Bunchy Sedge
<i>Dianella caerulea</i>	Blueberry Lily
<i>Dichondra repens</i>	Kydney Weed
<i>Digitaria didactyla</i>	Queensland Couch
<i>Dodonaea triquetra</i>	Forest Hop Bush
<i>Dodonaea viscosa</i>	Hop Bush
<i>Eragrostis brownii</i>	Browns Love Grass
<i>Eucalyptus acmenoides</i>	White Mahogany
<i>Eucalyptus fibrosa</i>	Broad Leaf Ironbark
<i>Eucalyptus microcorys</i>	Tallowwood
<i>Eucalyptus mollucana</i>	Gum Topped Box
<i>Eucalyptus siderophloia</i>	Grey ironbark
<i>Eucalyptus tereticornis</i>	Forest Red Gum
<i>Eustrephus latifolius</i>	Wombat Berry
<i>Glochidion ferdinandi</i>	Cheese Tree



<i>Glochidion sumatranum</i>	Umbrella Cheese Tree
<i>Gomphocarpus physocarpus</i>	Balloon Cotton Bush
<i>Imperata cylindrica</i>	Blady Grass
<i>Juncus sp.</i>	Juncus
<i>Lantana camara</i>	Lantana
<i>Lantana montevidensis</i>	Creeping Lantana
<i>Lepidosperma laterale</i>	Variable Sword Sedge
<i>Leucopogon juniperinus</i>	Prickly Heath
<i>Lobelia purpurascens</i>	White Root
<i>Lomandra longifolia</i>	Mat Rush
<i>Lomandra multiflora</i>	Many Flowered Mat Rush
<i>Lophostemon confertus</i>	Brush Box
<i>Lophostemon suaveolens</i>	Swamp Box
<i>Megathyrsus maximus</i>	Guinea Grass
<i>Melaleuca quinquenervia</i>	Broad Leaf Paperbark
<i>Melinis repens</i>	Red Natal Grass
<i>Oplismenus aemulus</i>	Creeping Beard Grass
<i>Opuntia tomentosa</i>	Prickly Pear
<i>Parsonsia straminea</i>	Monkey Rope
<i>Passiflora suberosa</i>	Corky Passion Flower
<i>Pennisetum purpureum</i>	Elephant Grass
<i>Persicaria decipiens</i>	Slender Knot Weed
<i>Philydrum lanuginosum</i>	Woolly Frogmouth
<i>Plectranthus parviflorus</i>	Plectranthus
<i>Pomax umbellata</i>	Pomax
<i>Pteridium esculentum</i>	Bracken Fern
<i>Rubusparvifolius</i>	Pink Flowered Native Raspberry
<i>Sesbania cannabina</i>	Yellow Pea Bush
<i>Setaria sphacelata</i>	Setaria
<i>Smilax australis</i>	Barbed Wire Vine
<i>Solanum mauritianum</i>	Wild Tobacco
<i>Solanum sp.</i>	Potato Bush
<i>Solanum torvum</i>	Devils Fig
<i>Sporobolus pyramidalis</i>	Giant Rat's Tail Grass
<i>Stephania japonica</i>	Tape Vine



<i>Themeda triandra</i>	Kangaroo Grass
<i>Trema tomentosa</i>	Poison Peach
<i>Typha sp.</i>	Bull Rush
<i>Vigna vexillata</i>	Wild Cowpea

On behalf of Lendlease Communities, please accept this exempt clearing notification (protected plants) for the site area known as Springfield Rise – Village 13 Basin & Village 15. Attached are the following documents:

- Notification form completed and signed
- Protected Plants Flora Survey Report

If you have any questions, please do not hesitate to contact me.

Kind regards,

Keira Grundy Environmental Planner **Saunders Havill Group**
direct line (07) 3251 9468 mobile 0437 822 880 email keiragrundy@saundershavill.com
phone 1300 123 SHG web www.saundershavill.com head office 9 Thompson St Bowen Hills Q 4006

Brisbane / Emerald / Rockhampton

Surveying / Town Planning / Urban Design / Environmental Management / Landscape Architecture

The information transmitted is for the use of the intended recipient only and may contain confidential and/or legally privileged material. Any review, re-transmission, disclosure, dissemination or other use of, or taking of any action in reliance upon, this information by persons or entities other than the intended recipient is prohibited. If you have received this email in error please delete all copies of this transmission together with any attachments and notify the sender. Opinions, conclusions and other information in this email that do not relate to the official business of Saunders Havill Group shall be understood as neither given nor endorsed by it. We have taken precautions to minimise the risk of transmitting software viruses, but we advise you to carry out your own virus checks on any attachment to this message. We cannot accept liability for any loss or damage caused by software viruses.

The information in this email together with any attachments is intended only for the person or entity to which it is addressed and may contain confidential and/or privileged material. There is no waiver of any confidentiality/privilege by your inadvertent receipt of this material.

Any form of review, disclosure, modification, distribution and/or publication of this email message is prohibited, unless as a necessary part of Departmental business.

If you have received this message in error, you are asked to inform the sender as quickly as possible and delete this message and any copies of this message from your computer and/or your computer system network.

Keira Grundy

Subject: FW: 7522: FW: AR082999 Exempt Clearing Email Lot 33 on SP269190

From: PALM [<mailto:palm@ehp.qld.gov.au>]

Sent: Friday, 22 January 2016 3:27 PM

To: Keira Grundy <keiragrundy@saundershavill.com>

Subject: RE: AR082999 Exempt Clearing Email Lot 33 on SP269190

Dear Mr Ian Murray

Applicant: Lend Lease Communities (Springfield) Pty Ltd

Exempt clearing notification (protected plants)

Where clearing is to be conducted –

Street Address: Sinnathamby Boulevard, Springfield

Lot/Plan: Lot 22 on SP234042 and Lot 33 on SP269190

EHP Reference: AR082999

Thank you for your email. Please retain this email as acknowledgement of receipt of a protected plant exemption notification submitted under section 261ZA of the Nature Conservation (Wildlife Management) Regulation 2006. Clearing of a protected plant under this section must be conducted within two years after the flora survey report was submitted to the Department of Environment and Heritage Protection.

It is strongly recommended for audit purposes that you keep this email together with the relevant flora survey trigger map, flora survey report and any other documentation relating to the clearing in question.

Please visit www.ehp.qld.gov.au for information about available online services.

Kind regards

Katrina



Katrina Theilemann

Administration Officer

Customer Service Team I Regulatory Capability and Customer Service

Department of Environment and Heritage Protection

P 1300 130 372 (option 4) F (07) 3330 5875 E Palm@ehp.qld.gov.au

400 George Street BRISBANE QLD 4000

GPO Box 2454, BRISBANE QLD 4001

environmental management



Spring Mountain Villages 6, 8, 13 & Haul Road Protected Plants Flora Survey Report

Lendlease
15th December 2015
7522



Document Control

Title	Spring Mountain – Villages 6, 8, 13 & Haul Road –Protected Plants Flora Survey Report
Job Number	7522
Client	Lendlease

Document Issue

Issue	Date	Prepared By	Checked By
Draft	15.12.2015	David Havill	Keira Grundy
Final			

Disclaimer

This report has been prepared for **Lendlease. Saunders Havill Group** cannot accept responsibility for any use of or reliance upon the contents of this report by any third party.

Reports and/or Plans by Others

Reports and/or plans by others may be included within this report to support the document.



Table of Contents

1. Introduction	4
1.1. Key Site Details	5
2. Desktop Assessment	11
2.1. Nature Conservation Act	11
2.2. Additional legislative instruments	11
3. Flora Survey Methodology	14
3.1. Clearing Impact Areas	14
3.2. Survey extent	14
3.3. Flora Survey Methodology	14
4. Flora Survey Results	16
4.1. Meander Transect 1	17
4.2. Meander transect 2	19
4.3. Meander transect 3	21
4.4. Meander transect 4	23
4.5. Meander transect 5	25
4.6. Meander transect 6	27
4.7. Meander transect 7	29
4.8. Meander transect 8	31
4.9. Meander transect 9	33
4.10. Meander transect 10	35
4.11. Meander transect 11	37
4.12. Meander transect 12	39
4.13. Summary	41
5. Appendices	42

Figures

Figure 1: Site Context

Figure 2: Site Aerial



Plans

Plan 1: Clearing Impact Area and Transect locations

Tables

Table 1: Wildlife Online Search Results - Flora
Table 2: Protected Matters Search Results - Flora
Table 3: Transect Coordinates
Table 4: Meander survey summary



I. Introduction

The *Environmental Management Division* of the **Saunders Havill Group** was engaged by **Lendlease** to prepare this Protected Plants Flora Survey Report to enable clearing within areas mapped as 'High Risk' under the *Nature Conservation Act 1992* (NCA). Clearing works are associated with early works stages at Spring Mountain, specifically Villages 6, 8 and 13 and a primary road connection known as the Haul Road. The Spring Mountain development site is located Sinnathamby Boulevard, Springfield Central (Lot 22 on SP234042 and Lot 33 on SP269190) and is within the jurisdiction of **Ipswich City Council** (ICC).

The **Queensland Government** has adopted a risk-based approach to the regulation of protected plants under the NCA. The regulatory framework captures activities that pose a high risk to plant biodiversity. Regulatory, educational and compliance effort are consequently focused on high risk activities. Under the framework, when a non-exempt clearing activity is proposed within a 'High Risk' area, the proponent of that activity is required to complete a flora survey prior to commencement of clearing.

The main objective of the flora survey is to locate any Endangered, Vulnerable or Near Threatened (EVNT) plants that may be present within the clearing impact area. This is especially important for determining the degree of assessment required for a particular clearing activity. For example, if the survey establishes that EVNT plant species are not present within the clearing impact area, the proposed clearing will be exempt and, following notification to the department, a clearing permit will not be required. Alternatively, if EVNT plant species are identified, and clearing is considered to impact on the EVNT plant (i.e. clearing comes within 100m of the EVNT plant) then an application for a Protected Plant Clearing Permit is required.


Contextually, the Spring Mountain project site is located to the west of Springfield Central, approximately 13km southeast of Ipswich City and approximately 26 km southwest of Brisbane City. The site is bordered by commercial development and educational facilities associated with Springfield Central to the northeast, residential development to the southeast and large vegetated rural properties adjoining White Rock-Spring Mountain Conservation Estate and more broadly the Flinders-Karawatha Bioregional Corridor. The site is bound by Centenary Highway to the north and Sinnathamby Boulevard to the east. The surrounding suburbs of Redbank Plains, Springfield Lakes and Swanbank are highly urbanised and contain a mixture of residential housing, commercial properties and industrial land uses. Refer to **Figures 1 and 2** for site context and aerial. Clearing works proposed within Village 6, 8, 13 and the Haul Road form part of early works for the commencement of the Spring Mountain project which forms part of the Greater Springfield urban development area (refer **Plan 1**). It is noted that the Spring Mountain project (refer **Plan 2**) has been approved by the Commonwealth **Department of the Environment** (DoE) (EPBC 2013/7057).

The flora surveys outlined in this report were conducted where proposed clearing is mapped within 'High Risk' areas under Protected Plants Flora Survey Trigger Mapping (refer **Figure 3**) as per the *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992*.



I.I. Key Site Details

Address	Sinnathamby Boulevard
RPD	Lot 22 on SP234042, Lot 33 on SP269190
Local Government Area	Ipswich City Council
Planning Scheme	Springfield Structure Plan, which forms part of the Ipswich City Council Planning Scheme 2003
Area Classification/Zone	Community Residential
Existing Land Use	Vacant
Proposed Land Use	Residential / Road



© OpenStreetMap (and) contributors, CC-BY-SA

Legend


 Project impact area

Figure 1 Site Context


File ref. 7522 E 01 Site Context A
Date 5/01/2016
Project Springfield Villages 6 & 8

0 250 500 1,000 1,500 m

Scale (A4): 1:50,000 [GDA 1994 MGA Z56]



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend


 Project impact area


Figure 2 Site Aerial

File ref. 7522 E 02 Site Aerial A
Date 5/01/2016
Project Springfield Villages 6 & 8

0 100 200 400 600 m
Scale (A4): 1:15,837 [GDA 1994 MGA Z56]



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend



-  Project impact area
-  High Risk


Figure 3 NCA Flora Survey Trigger Map

File ref. 7522 E 03 NCA A
Date 5/01/2016
Project Springfield Villages 6 & 8

0 100 200 400 600 m
 Scale (A4): 1:15,837 [GDA 1994 MGA Z56]




THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWING BY ANY THIRD PARTY.



Legend

- Project impact area
- Community residential
- Town centre
- Regional transport corridor
- Open space
- Conservation



APPROVED COMPANY
ISO 9001
Quality Management Systems
QMS
Certification Services

APPROVED COMPANY
ISO 14001
Environmental Management Systems
QMS
Certification Services

Spring Mountain- Villages 6 , 8 & 13 & Haul Road

Greater Springfield Structure Plan

Plan 1


ISSUES:

Issue	Date	Description	Drawn	Checked
A	5/01/2016	Prelim Draft	AL	MS

Date | 5/01/2016
 Scale | 1:32,500 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client | Lend Lease
 Project | NCA
 Address/RPD | Springfield
 Source | DCBD (DNRM, 2013), Aerial (QLD Globe, 2013)
 Layout (LandPartners 2014)

SHG File
7522 E 02 Structure Plan A


THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAILED LOCATIONS OF ALL SERVICES.



Legend

- Project impact area
- RESIDENTIAL DEVELOPMENT AREA
- OPTIONAL USE AREA B
- EXISTING SCHOOL
- PROPOSED SCHOOL
- PROPOSED NEIGHBOURHOOD CENTRE
- ACTIVITY CENTRE
- INFRASTRUCTURE (EXISTING)
- DISTRICT 2 PARK (D2P1)
- MAJOR DISTRICT PARK NORTH (FP1)
- FORMAL PARK WEST (FP2)
- LINEAR CREEKLINE OPEN SPACE
- LOCAL RECREATION PARK
- CENTENARY HIGHWAY
- SINNATHAMBY BOULEVARD
- TOWN CENTER ROAD TYPE 2A MODIFIED
- MAJOR COLLECTOR DUAL LANE STREET (NO ACCESS)
(Area shown is where road has been defined)
- MAJOR COLLECTOR STREET (ACCESS)
(Area shown is where road has been defined)
- COLLECTOR STREET
- VILLAGE BOUNDARY
- SPRING MOUNTAIN PRECINCT BOUNDARY 352 ha
- INDICATIVE TOWN CENTRE DEVELOPMENT AREAS (SUBJECT TO TOWN CENTRE CONCEPT PLAN)

NOTE:
LOCATIONS AND AREAS OF VARIOUS LAND USE ELEMENTS AS SHOWN ARE CONCEPTUAL ONLY AND SUBJECT TO DETAILED DESIGN



Spring Mountain - Villages 6, 8 13 & Haul Road

Spring Mountain Development Proposal

Date: 5/01/2016
 Scale: 1:15,000 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client: Lend Lease
 Project: NCA
 Address/ RPD: Springfield
 Source: Development Layout (LandPartners 2015)

Plan 2

SHG File
7522 E 01 Draft Layout A

ISSUES:	Issue	Date	Description	Drawn	Checked
A	5/01/2016	Prelim Draft		AL	MS



THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN MILLIMETRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAILED LOCATIONS OF ALL SERVICES.



2. Desktop Assessment

2.1. Nature Conservation Act

The NCA classifies and protects significant areas (Protected Areas) and protects threatened plant and animal species. The *Nature Conservation (Wildlife) Regulation 1994* (NCWR) lists plant and animal species presumed extinct, endangered, vulnerable, near threatened, least concern, international or prohibited.

The **Queensland Government** has adopted a regulatory framework that captures activities that pose a high risk to plant biodiversity. Under the framework, when a non-exempt clearing activity is proposed within a 'High Risk' area, the proponent of that activity is required to complete a flora survey prior to commencement of clearing. The Protected Plants Flora Survey Trigger Map shows 'High Risk' areas for protected plants and is used to help determine flora survey and clearing permit requirements for a particular location.

A search of Projected Plants Flora Survey Trigger Mapping indicated proposed clearing areas within the subject site are overlaid as 'High Risk' and so are subject to flora survey requirements (refer **Figure 3**).

Prior to flora surveys, the schedules of the NCWR were considered in this report using a Wildlife Online Database Search with a 10 kilometre radius from the site. Six (6) flora species listed under the NCWR were identified as having the potential to occur on site and are presented in **Table 1**. Refer to **Appendix A** for full search results.

Table 1: Wildlife Online Search Results - Flora

Scientific Name	Common Name	Status
<i>Marsdenia coronata</i>	Slender Milk Vine	Vulnerable
<i>Plectranthus habrophyllus</i>	-	Endangered
<i>Eucalyptus curtisii</i>	Plunkett Mallee	Near Threatened
<i>Melaleuca irbyana</i>	Swamp Tea Tree	Endangered
<i>Notelaea ipsviciensis</i>	-	Endangered
<i>Notelaea lloydii</i>	Lloyd's Native Olive	Vulnerable

2.2. Additional legislative instruments

In order to maximise the scope of the flora survey, a search of protected matters listed as potentially present within 10 km of the sites under the Federal *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) was conducted using the Protected Matters Search Tool. Potential flora EVNT species listed under the EPBC Act are presented in **Table 2**. Refer to **Appendix B** for full search results.



Table 2: EPBC Act Protected Matters Search Results - Flora

Scientific Name	Common Name	Status
<i>Arthraxon hispidus</i>	Hairy Joint Grass	Vulnerable
<i>Bosistoa transversa</i>	Three-leaved Bosistoa	Vulnerable
<i>Cupaniopsis tomentella</i>	Boonah Tuckeroo	Vulnerable
<i>Notelaea ipsviciensis</i>	Cooneana Olive	Critically Endangered
<i>Notelaea lloydii</i>	Lloyd's Olive	Vulnerable
<i>Phaius australis</i>	Lesser Swamp-orchid	Endangered
<i>Phebalium distans</i>	My Berryman Phebalium	Critically Endangered
<i>Planchonella eerwah</i>	Shiny-leaved Condoo	Endangered
<i>Plectranthus habrophyllus</i>	-	Endangered
<i>Sophora fraseri</i>	-	Vulnerable
<i>Thesium australe</i>	Austral Toadflax	Vulnerable

Regional Ecosystem mapping under the *Vegetation Management Act, 1999* (VMA) was utilised to inform flora survey targets and techniques. The broader area where the survey sites occur is mapped under the VMA as Least Concern 12.9-10.19a, 12.9-10.17a, 12.9-10.2, and 12.9-10.7 as described below and highlighted in **Plan 3**.

Least Concern RE 12.9-10.19a

Description	<i>Corymbia henryi</i> +/- <i>Eucalyptus fibrosa</i> subsp. <i>Fibrosa</i> , <i>Corymbia citriodora</i> subsp. <i>Variegata</i> , <i>Eucalyptus siderophloia</i> , <i>Eucalyptus crebra</i> open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
--------------------	---

Least Concern RE 12.12.17a



Description	<i>Lophostemon confertus</i> or <i>Lophostemon suaveolens</i> dominated open forest usually with emergent <i>Eucalyptus</i> and/or <i>Corymbia</i> species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments.
--------------------	--

Least Concern RE 12.9-10.2





Description	<i>Corymbia citriodora</i> subsp. <i>Variegata</i> open forest or woodland usually with <i>Eucalyptus crebra</i> . Other species such as <i>Eucalyptus tereticornis</i> , <i>Eucalyptus moluccana</i> , <i>Eucalyptus acmenoides</i> and <i>Eucalyptus siderophloia</i> may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of <i>Lophostemon confertus</i> (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
--------------------	---


Of Concern RE 12.9-10.7

Description	<i>Eucalyptus crebra</i> +/- <i>Eucalyptus tereticornis</i> , <i>Corymbia tessellaris</i> , <i>Angophora leiocarpa</i> , <i>Eucalyptus melanophloia</i> woodland. Occurs on Cainozoic and Mesozoic sediments.
--------------------	---



Legend

-  Project impact area
- Regional Ecosystem v8**
-  Category A or B area containing endangered regional ecosystems
-  Category A or B area containing of concern regional ecosystems
-  Category A or B area that is a least concern regional ecosystem



Spring Mountain- Villages 6, 8 & 13 & Haul Road

Regional Ecosystems v8

Plan 3

THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.

CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN MILLIMETRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.

PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAIL LOCATIONS OF ALL SERVICES.

Issue	Date	Description	Drawn	Checked
A	5/01/2016	Prelim Draft	AL	MS

Date | 5/01/2016
 Scale | 1:14,962 @ A3
 Data Information:
 Universal Transverse Mercator
 GDA 1994 MGA Zone 56
 Client | Lend Lease
 Project | NCA
 Address/RPD | Springfield
 Source | DCBD (DNRM, 2013), Aerial (QLD Globe, 2013)

SHG File
 7522 E 03 REs A



3. Flora Survey Methodology

3.1. Clearing Impact Areas

The proposed clearing sites (i.e. Villages 6, 8, 13 and the Haul Road) are mostly mapped as 'High Risk' areas under Protected Plants Flora Survey Trigger (refer **Figure 3**). The Clearing Impact Areas, which are identified the areas to be cleared inclusive of a 100m buffer, are shown in **Plan 4**.

3.2. Survey extent

Table 3 and **Plan 4** summarise the Clearing Impact Areas and Transect extents. General observations for EVNT flora species were conducted at all times while on-site, including while traversing roads and vegetated area both inside and outside designated Clearing Impact Areas. The 100m buffer areas was assessed where access was possible.




Table 3: Transect Coordinates

Transect	Start	Finish
1	-27.703174° / 152.909798°	-27.702452° / 152.909382°
2	-27.702202° / 152.906698°	-27.698432° / 152.905453°
3	-27.695617° / 152.905829°	-27.693931° / 152.903303°
4	-27.694879° / 152.901439°	-27.699177° / 152.900416°
5	-27.700895° / 152.902626°	-27.700693° / 152.909101°
6	-27.683117° / 152.895659°	-27.681752° / 152.894641°
7	-27.683179° / 152.892057°	-27.683791° / 152.890378°
8	-27.686838° / 152.890317°	-27.688842° / 152.892838°
9	-27.689488° / 152.891223°	-27.688196° / 152.889467°
10	-27.691064° / 152.892680°	-27.692380° / 152.895896°
11	-27.688213° / 152.894579°	-27.685155° / 152.895197°
12	-27.684803° / 152.894378°	-27.688865° / 152.894291°

3.3. Flora Survey Methodology

The clearing sites were surveyed using the preferred timed meander survey technique as per *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992* by three (3) suitably qualified professionals including (1) Senior Ecologists and two (2) Ecologists (refer to **Appendix C** for curricula vitae). Surveys were carried out as follows:

- 1) The Clearing Impact Areas were traversed on foot by project Ecologists (refer to **Plan 4**).
- 2) The start and finish time of each meander was recorded.
- 3) The track log of project Ecologist's transects was recorded using a handheld GPS unit accurate to < 1m.
- 4) The identity of all plant species encountered during each meander was recorded.
- 5) The site and surrounds were photographed.



APPROVED COMPANY
ISO 9001
Quality Management Systems
QMS
APPROVED COMPANY
ISO 14001
Environmental Management Systems
QMS

Spring Mountain - Villages 6, 8 & 13 & Haul Road

Flora Meandering Survey Transects

ISSUES:

Issue	Date	Description	Drawn	Checked
A	5/01/2016	Prelim Draft	TC	MS

Date | 5/01/2016
 Scale | 1:12,500 @ A3
 Coordinate System | GDA 1994 MGA Zone 56
 Projection | Transverse Mercator
 Client | Lend Lease
 Project | Springfield
 Address/RPD | Springfield Villages 6 to 8
 Sources | QLD GIS Layers (QLD Gov. Info Services 2015),
 Aerial (Nearmap, 2015)

Plan 4

SHG File
7522 E 01 Flora Meandering Survey B

THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAIL LOCATIONS OF ALL SERVICES.



4. Flora Survey Results

The Clearing Impact Areas were assessed on 8 and 9 July 2015. **No EVNT species were encountered in any of the proposed clearing areas** however a population of translocated *Marsdenia coronate* (Slender Milk Vine) was recorded within the 100m buffer. Given the extent of survey it can be stated with a high level of confidence that no EVNT species will be cleared by the proposed development.

A total of one hundred and thirty seven (137) species were identified throughout the survey period. This included fourteen (14) herbs, thirteen (13) vines, three (3) orchids and epiphyte species, forty six (46) ground layer species, twenty five (25) shrubs, twenty (20) sub-canopy species and sixteen (16) canopy species. The transect length varied however a total of 11.813 kilometres were searched for threatened species by three ecologists using the meander methods. Each transect was located in areas which represented each mapped vegetation community verified through extensive site surveys.

It is noted however that *Marsdenia coronate* (Slender Milk Vine) has been recorded within the buffer area adjacent to Transect 1 (refer **Plan 1**). These specimens form part of a previously approved translocation program and are located within a Council Park known as Marsdenia Park, within the existing residential development to the east. The proposed works will not impact on these specimens which are separated from the project area by an existing bitumen road.

Table 4 summarises the details of each of the timed meander transects. Meander transect descriptions with photographs are presented in the following pages. A general description for each transect area is provided in this section and respective species lists in **Appendix D**.

Table 4: Meander survey summary

Site	Date	Start Time	Finish Time	Duration	Distance	Flora Species
1	1.12.2015	11.05am	12.45pm	100 minutes	1.161km	55
2	1.12.2015	12.25pm	13.45	80 minutes	1.117km	39
3	1.12.2015	13.46	15.08	92 minutes	888m	52
4	1.12.2015	14.55	16.18	83 minutes	1.149km	46
5	1.12.2015	16.00	17.15	75 minutes	1.189km	42
6	2.12.2015	9.31am	10.40am	69 minutes	480m	79
7	2.12.2015	10.23am	11.47am	85 minutes	982m	45
8	2.12.2015	11.25am	12.27pm	62 minutes	756m	47
9	2.12.2015	12.31	13.55	86 minutes	1.019km	58
10	2.12.2015	13.42	14.42	60 minutes	696m	44
11	14.12.2015	13.36	15.08	92 minutes	1.019m	51
12	14.12.2015	14.55	16.27	92 minutes	1.357km	64



4.1. Meander Transect I

Transect 1 is located within mapped remnant vegetation dominated by Least Concern Regional Ecosystem community 12.9-10.19a. This community is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variagate*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments. Transect searches extended along 1.1.61 kilometres. Canopy species recorded are consistent with current regional ecosystem mapping. *Corymbia henryi* (Large Leaf Spotted Gum) and *Eucalyptus fibrosa* (Broad Leaf Ironbark) were the dominant species recorded.

The Transect 1 Area is located towards the edge of the existing residential development, on North West facing slopes. The canopy and sub-canopy tree layers are largely intact with disturbances confined to some minimal historic tree removal including evidence of logging practices. The shrub and ground layer are dominated by native species with the majority of introduced species confined to the occasional small clump of *Lantana camara* (Lantana) and patches of introduced grasses and weeds along the edge of the vegetated patch and within the cleared easement track which runs to an existing water tower directly south of the transect. The shrub layer is very sparse with the ground layer patchy in areas amongst areas of exposed earth and leaf litter.

Only a small area of exposed rock surface was observed along a portion of the ridge line adjacent to the cleared track throughout the transect area. This area was thoroughly searched specifically for both *Marsdenia coronata* (Slender Milk Vine) and *Plectranthus habrophyllus* (Plectranthus) both of which have habitat niches suited to this terrain. The remaining area retained an open understorey and ground layer.



Photo: Transect 1 dominated by *Corymbia henryi* and *Eucalyptus fibrosa*.



Photo: Exposed rocky terrain observed along the ridge line.

Fifty five (55) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. Flora diversity consisted of two (2) herbs, four (4) vines, twenty six (26) ground layer, twelve (12) shrub, seven sub-canopy and four (4) canopy species.



4.2. Meander Transect 2

Transect 2 is located within mapped remnant vegetation dominated by Least Concern Regional Ecosystem community 12.9-10.19a. This community is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variagate*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments. Transect searches extended along 1.117 kilometres.

Species recorded within the canopy are dominated by *Eucalyptus fibrosa* (Broad Leaf Ironbark). This dominant species is scattered amongst *Corymbia henryi* (Large Leaf Spotted Gum) and the occasional *Corymbia intermedia* (Pink Bloodwood) and *Eucalyptus acmenoides* (White Mahogany). This transect is consistent with the current remnant regional ecosystem mapping. A patchy shrub layer was recorded throughout the transect area however overall is considered relatively sparse. The ground layer also varied from relatively sparse amongst the areas with exposed rock along the ridge lines with greater densities recorded on slopes and towards the lower portion of the transect.

Disturbances within this transect are restricted to some introduced species within the ground layer which are mainly concentrated along the edges of vehicle access tracks. Some evidence of logging and fire is also noted throughout the survey.

Some exposed rocky outcrops, limited to along the ridgeline, were recorded by field survey. The remaining area is typical of Landzone 9-10, containing fine to coarse grained sedimentary rocks.



Photo: Transect 2 dominated by *Eucalyptus fibrosa* and *Corymbia henryi*



Photo: Minimal exposed rock outcrops.

Thirty nine (39) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included one (1) herb, four (4) vines, twelve (12) ground layer, ten (10) shrub, five (5) sub-canopy and seven (7) canopy species.



4.3. Meander Transect 3

Transect 3 is located within mapped remnant vegetation dominated by least concern regional ecosystem 12.9-10.17. This community is described as *Lophostemon confertus* or *Lophostemon suaveolens* dominated open forest usually with emergent *Eucalyptus* and/or *Corymbia* species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments. The transect survey included investigations along 888m.

This transect is located within vegetation that is typical of lower gully lines with increase densities of *Lophostemon suaveolens* (Swamp Box). There is a greater density of weed species recorded throughout this transect which occurred along the edges of the cleared adjacent development area directly south as well as throughout the mapped waterway. It is noted that thick patches of *Lantana camara* (Lantana) was recorded along the edges of this mapped waterway. The ground layer was relatively dense with leaf litter and bare earth confined to isolated small patches.



Photo: *Eucalyptus* and *Corymbia* species dominated the hill side with *Lophostemon suaveolens* dominated the lower embankment area.



Photo: Steep south west facing slope

Although canopy species recorded are consistent with current regional ecosystem mapping, the age structure appeared to be reduced with the number of large trees remaining previously removed through historical logging practices. The height of this vegetation community however remains at remnant status.

Fifty two (52) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity consisted of five (5) herb, three (3) vines, nineteen (19) ground layer, eleven (11) shrub, six (6) sub-canopy and eight (8) canopy species.



4.4. Meander Transect 4

Transect 4 is located within mapped remnant vegetation dominated by Least Concern Regional Ecosystem community 12.9-10.19a. This community is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variagate*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments. The transect included investigations along 1.149 kilometres.

Although elements of Least Concern Regional Ecosystem 12.9-10.19 were recorded throughout this transect, some species representing Least Concern Regional Ecosystem 12.9-10.2 and Of Concern Regional Ecosystem 12.9-10.7 were also observed however were too small to separate through mapping amendments. The shrub layer is dominated by *Acacia* species including *Acacia leiocalyx* (Early Flowering Black Wattle), *Acacia disparrima* (Hickory Wattle) and *Acacia concurrens* (Black Wattle). This appeared to be a result of fire activity which was evident towards the canopy of some of the established canopy trees. The ground layer is recorded as being dense and is dominated by *Themeda triandra* (Kangaroo Grass) and *Imperata cylindrica* (Blady Grass).

The majority of this transect is located on a western facing slope with weeds confined to the lower portion of the hill and is dominated by *Lantana camara* (Lantana). The site also retained evidence of fire and some past logging activities.



Photo: Fire evidence throughout the transect area typical with *Acacia* regrowth within the shrub layer.



Photo: Species recorded typical of the current regional ecosystem mapping.

The vegetation community retains a canopy height and structure which retains its remnant status. Also observed were a number of well-established specimens which appeared to be less favourable for past logging practices.

Forty six (46) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included two (2) herb, four (4) vines, nineteen (19) ground layer, eight (8) shrub, seven (7) sub-canopy and six (6) canopy species.



4.5. Meander Transect 5

Transect 5 is located within two (2) regional ecosystem communities both of which are categorised as Least Concern regional ecosystems. The areas outside of the mapped waterway is described as RE12.9-10.19a whereas the vegetation associated with the drainage line is described as RE12.9-10.17a. The transect survey included investigations along 1.189 kilometres.

The majority of Transect 5 is located within mapped remnant vegetation dominated by Least Concern Regional Ecosystem community 12.9-10.19a. This community is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variagate*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments. Transect 5 is also traverses across a mapped drainage line and is located within mapped remnant vegetation dominated by least concern regional ecosystem 12.9-10.17. This community is described as *Lophostemon confertus* or *Lophostemon suaveolens* dominated open forest usually with emergent *Eucalyptus* and/or *Corymbia* species. Occurs in gullies and southern slopes on Cainozoic and Mesozoic sediments.

The changes between the two regional ecosystem communities appeared evident with the increase in density of *Lophostemon suaveolens* (Swamp Box) associated with the drainage line or lower lying areas. The Landzone between these two communities is mapped the same, however it is noted that a very small portion of this drainage feature contains some deposited material and contains characteristics of Landzone 3. This portion of the Least Concern RE12.9-10.17 area is too small to map within the regional ecosystem framework. Apart from the occasional small patch of *Juncus* sp, there were limited changes in flora species recorded.



Photo: Majority of transect located within RE12.9-10.19a.



Photo: Transect intersected mapped drainage feature mapped as RE12.9-10.17.

The majority of this transect contained very little shrub layer coverage with the density of the ground layer relatively high. Patches of bare earth and leaf litter were confined to some isolated small patches.

Forty three (43) species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included two (2) herb, four (4) vines, twelve (12) ground layer, twelve (12) shrub, seven (7) sub-canopy and five (5) canopy species.



4.6. Meander Transect 6

Transect 6 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey included investigations along 480 metres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Transect located within a composite Regional Ecosystem community.



Photo: Greater density of weed invasion towards the lower slopes of the transect area.

The majority of Transect 6 is located on a north facing slope and on the southern side of a mapped waterway. Disturbances were confined to selective canopy thinning through logging practices, cleared vehicle tracks as well as weed infestations. The whole of transect area contained evidence of fire with patches of *Imperata cylindrica* (Blady Grass) dominating the ground layer as well as a shrub layer dominated by *Acacia* species.

The diversity of species recorded within this transect is a result of the mapped composite regional ecosystem community. Patches of vegetation were dominated by species representing each of the regional ecosystem communities however the understorey, including the shrub and ground layer remained relatively consistent throughout the entire transect area. Small changes in species were recorded within areas containing exposed rocky outcrops and within the low lying areas associated with overland flow paths.

Seventy nine (79) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included five (5) herb, six (6) vines, three (3) orchids/epiphytes, thirty two (32) ground layer, twelve (12) shrub, thirteen (13) sub-canopy and eight (8) canopy species.



4.7. Meander Transect 7

Transect 7 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey included investigations along 982 metres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.

The dominant regional ecosystem observed throughout the transect area is recorded as the Least Concern RE12.9-10.2 however elements of RE12.9-10.19 and Of Concern RE12.9-10.7 were observed within small patches within and adjacent to this transect. Small habitat variations were recorded within areas containing exposed rock however these patches were confined to isolated areas towards the ridgeline. Limited diversity was recorded within the shrub layer limited to three native species amongst patches of *Lantana camara* (Lantana).

Greater disturbances were recorded within the canopy layer within this portion of the site resulting in greater weed invasion and higher density of ground layer species dominated by *Imperata cylindrica* (Blady Grass).



Photo: Transect located within a mapped composite regional ecosystem community.



Photo: Transect located within a mapped composite regional ecosystem community.

Forty five (45) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included five (5) herb, five (5) vines, eighteen (18) ground layer, four (4) shrub, six (6) sub-canopy and seven (7) canopy species.



4.8. Meander Transect 8

Transect 8 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey included investigations along 786 metres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Some exposed rocky outcrops thoroughly searched for threatened plants



Transect 8 contained some exposed rock outcrop areas however the majority of the investigation area contained a thick dense a grass layer with a scattered or sparse shrub layer amongst a woodland community. Although some introduced species were observed within the ground layer, these specimens were generally associated with old vehicle access tracks most likely as a result of logging activities.

Species recorded within the canopy are dominated by *Corymbia citriodora* (Spotted Gum), and *Eucalyptus siderophloia* (Grey Ironbark). This dominant species is recorded amongst scattered *Corymbia henryi* (Large Leaf Spotted Gum) and the occasional *Eucalyptus seeana* (Narrow Leaf Red Gum) and *Angophora leiocarpa* (Smooth Bark Apple). This transect is consistent with the current remnant regional ecosystem mapping. A patchy shrub layer was recorded throughout the transect area however overall was relatively sparse. The ground layer also varied from relatively sparse amongst the areas with exposed rock along the ridge lines with greater densities recorded on slopes and towards the lower portion of the transect.



Photo: Very few introduced species recorded throughout the transect

Forty seven (47) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included five (5) herb, five (5) vines, twenty (20) ground layer, three (3) shrub, six (6) sub-canopy and eight (8) canopy species.



4.9. Meander Transect 9

Transect 9 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey included investigations along 1.019 kilometres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Transect dominated by *Corymbia citriodora*



The dominant regional ecosystem community recorded within the transect area is Least Concern RE12.9-10.2 with *Corymbia citriodora* (Spotted Gum) being the dominant species recorded within the canopy layer. Other canopy species included *Angophora leiocarpa* (Smooth Bark Apple), *Corymbia intermedia* (Pink Bloodwood), *Corymbia trachyphloia* (Brown Bloodwood), *Eucalyptus acmenoides* (White Mahogany), *Eucalyptus seeana* (Narrow Leaf Red Gum) and *Eucalyptus siderophloia* (Grey Ironbark). Disturbances within this transect were restricted to some introduced species within the ground layer which were mainly concentrated along the vehicle access tracks. Some evidence of logging and fire were also recorded throughout the survey.

Fifty eight (58) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included six (6) herb, six (6) vines, twenty two (22) ground layer, nine (9) shrub, eight (8) sub-canopy and seven (7) canopy species.



4.10. Meander Transect IO

Transect 10 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey included investigations along 696 metres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Evidence of fire with greater densities of Acacia regrowth.



Canopy species recorded throughout transect 10 include scattered occurrences of *Angophora leiocarpa* (Smooth Bark Apple), *Corymbia intermedia* (Pink Bloodwood), *Corymbia trachyphloia* (Brown Bloodwood), *Eucalyptus microcorys* (Tallowwood), *Eucalyptus seeana* (Narrow Leaf Red Gum), and *Eucalyptus siderophloia* (Grey Ironbark).

Disturbances within this transect were restricted to some introduced species within the ground layer which were mainly concentrated along the vehicle access tracks. Some evidence of logging and fire were also recorded throughout the survey. Species recorded within the shrub layer were dominated by *Acacia* species including *Acacia leiocalyx* (Early Flowering Black Wattle), *Acacia concurrens* (Black Wattle) and *Acacia disparrima* (Hickory Wattle).

The ground layer was relatively dense with the occasional rocky outcrop and small patches of leaf litter and bare earth.

Forty four (44) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included three (3) herb, four (4) vines, seventeen (17) ground layer, eight (8) shrub, six (6) sub-canopy and six (6) canopy species.



4.II. Meander Transect II

Transect 11 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey length included investigations along 1.019 kilometres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Site dominated by *Corymbia citriodora*



Photo: Minimal exposed rock.

The canopy layer is dominated by *Corymbia citriodora* (Spotted Gum), with occasional occurrences of *Angophora leiocarpa* (Smooth Bark Apple), *Corymbia henryi* (Large Leaf Spotted Gum), *Corymbia intermedia* (Pink Bloodwood), *Eucalyptus acmenoides* (White Mahogany), *Eucalyptus seeana* (Narrow Leaf Red Gum), *Eucalyptus siderophloia* (Grey Ironbark) and *Eucalyptus tereticornis* (Forest Red Gum).

Disturbances within this transect were restricted to some introduced species within the ground layer which were mainly concentrated along the vehicle access tracks. Some evidence of logging and fire were also recorded throughout the survey.

Fifty one (51) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included five (5) herb, six (6) vines, sixteen (16) ground layer, nine (9) shrub, seven (7) sub-canopy and eight (8) canopy species.



4.12. Meander Transect 12

Transect 12 is located within mapped remnant vegetation dominated by a composite regional ecosystem community including 65% Least Concern RE12.9-10.2, 20% Of Concern RE12.9-10.7 and 15% Least Concern RE12.9-10.19. The transect survey length included investigations along 1.357 kilometres.

- Least Concern Regional Ecosystem community 12.9-10.19a is described as *Corymbia henryi* +/- *Eucalyptus fibrosa* subsp. *Fibrosa*, *Corymbia citriodora* subsp. *Variegata*, *Eucalyptus siderophloia*, *Eucalyptus crebra* open forest. Occurs in coastal areas on Cainozoic and Mesozoic sediments.
- Least Concern Regional Ecosystem 12.9-10.2 is described as *Corymbia citriodora* subsp. *Variegata* open forest or woodland usually with *Eucalyptus crebra*. Other species such as *Eucalyptus tereticornis*, *Eucalyptus moluccana*, *Eucalyptus acmenoides* and *Eucalyptus siderophloia* may be present in scattered patches or in low densities. Understorey can be grassy or shrubby. Shrubby understorey of *Lophostemon confertus* (whipstick form) often present in northern parts of bioregion. Occurs on Cainozoic and Mesozoic sediments.
- Of Concern Regional Ecosystem 12.9-10.7 is described as *Eucalyptus crebra* +/- *Eucalyptus tereticornis*, *Corymbia tessellaris*, *Angophora leiocarpa*, *Eucalyptus melanophloia* woodland. Occurs on Cainozoic and Mesozoic sediments.



Photo: Site vegetation consistent with current regional ecosystem mapping.



Photo: Sparse understorey typical of the regional ecosystem communities searches.

Disturbances within this transect were restricted to some introduced species within the ground layer which were mainly concentrated along the vehicle access tracks. Some evidence of logging and fire were also recorded throughout the survey.

Sixty four (64) flora species were recorded throughout the transect area, all of which are listed as common under state and federal legislation. This diversity included five (5) herb, six (6) vines, twenty seven (27) ground layer, seven (7) shrub, eleven (11) sub-canopy and eight (8) canopy species.



4.13. Summary

Field surveys were carried out within the clearing impact area and buffer of early works precincts (Village 6, 8, 13 and the Haul Road) of the Spring Mountain project site which is mapped as 'High Risk' by Protected Plants Flora Survey Trigger Mapping. The surveys utilised the preferred random meander technique as outlined in the *Flora Survey Guidelines – Protected Plants Nature Conservation Act 1992* to identify the presence of EVNT species. Coverage included the clearing extents as well as a 100 m buffer with each Clearing Impact Area almost entirely traversed during the timed meander transects. Twelve (12) meander transects as well as continual observations were completed throughout the investigation area.

The following points provide a summary of the investigation area:

- The vegetation communities observed have been extensively searched and analysed against current regional ecosystem mapping with overall consistency in the location of reach regional ecosystem community. Some minor variations were observed however in the majority of areas these variations are too small to provide for changes to this mapping.
- The majority of the clearing site's canopy is relatively in-tact representing an open forest to woodland community. Although evidence of forestry practices were recorded in all transects and throughout observational survey points, the site remains as remnant due to the vegetation community's height and density.
- The sub-canopy layer is relatively sparse throughout the majority of the site and is typical of the mapped vegetation communities represented on site.
- The shrub layer is relatively sparse and in some areas is almost completely absent, which is typical of the mapped regional ecosystem communities. However evidence of fire and some vegetation clearing was recorded throughout the majority of all transects.
- Weed invasion in most areas was largely confined to areas that have been cleared including vehicle access tracks and easements as well as greater densities recorded within overland flow paths and mapped waterways and drainage lines.
- Exposed rocky habitat was recorded in isolated patches along ridge lines as well as along major creek lines. Although these areas have been extensively searched, no threatened species were recorded at the time of the assessment within the investigation area.
- *Marsdenia coronate* (Slender Milk Vine) has been recorded within the buffer area adjacent to Transect 1. These specimens form part of a previously approved translocation program and are located within a Council Park known as Marsdenia Park, within the existing residential development to the east. The proposed works will not impact on these specimens which are separated from the project area by a 20m wide existing bitumen road.

Surveys **did not identify any EVNT species within the proposed clearing areas** however a population of translocated *Marsdenia coronate* (Slender Milk Vine) was located within the 100m buffer. While this protected species is located with the 'Clearing Impact Area' as defined by the *Flora Survey Guidelines*, as no impacts to EVNT species will occur as a result of the proposed clearing, an 'Exempt Clearing Notification' form should be lodged with the *Department of Environment and Heritage Protection* prior to any clearing taking place.



5. Appendices

Appendix A

Wildlife Online Search Results

Appendix B

Protected Matters Search Results

Appendix C

Curricula Vitae

Appendix D

Species Lists



Appendix A

Wildlife Online Search Results



Queensland Government

Wildlife Online Extract

Search Criteria: Species List for a Specified Point

Species: All

Type: All

Status: Rare and threatened species

Records: All

Date: All

Latitude: -27.6906

Longitude: 152.8996

Distance: 10

Email: davidhavill@saundershavill.com

Date submitted: Monday 30 Nov 2015 15:16:35

Date extracted: Monday 30 Nov 2015 15:20:08

The number of records retrieved = 19

Disclaimer

As the DSITIA is still in a process of collating and vetting data, it is possible the information given is not complete. The information provided should only be used for the project for which it was requested and it should be appropriately acknowledged as being derived from Wildlife Online when it is used.

The State of Queensland does not invite reliance upon, nor accept responsibility for this information. Persons should satisfy themselves through independent means as to the accuracy and completeness of this information.

No statements, representations or warranties are made about the accuracy or completeness of this information. The State of Queensland disclaims all responsibility for this information and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you may incur as a result of the information being inaccurate or incomplete in any way for any reason.

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	amphibians	Limnodynastidae	<i>Adelotus brevis</i>	tusked frog		V		10
animals	birds	Cacatuidae	<i>Calyptorhynchus lathami lathami</i>	glossy black-cockatoo (eastern)		V		10
animals	birds	Falconidae	<i>Falco hypoleucos</i>	grey falcon		V		1
animals	birds	Maluridae	<i>Stipiturus malachurus</i>	southern emu-wren		V		1
animals	birds	Psittacidae	<i>Lathamus discolor</i>	swift parrot		E	E	3
animals	birds	Rostratulidae	<i>Rostratula australis</i>	Australian painted snipe		V	E	8
animals	birds	Strigidae	<i>Ninox strenua</i>	powerful owl		V		13
animals	birds	Turnicidae	<i>Turnix melanogaster</i>	black-breasted button-quail		V	V	1
animals	mammals	Dasyuridae	<i>Dasyurus maculatus maculatus</i>	spotted-tailed quoll (southern subspecies)		V	E	3
animals	mammals	Macropodidae	<i>Petrogale penicillata</i>	brush-tailed rock-wallaby		V	V	8
animals	mammals	Phascolarctidae	<i>Phascolarctos cinereus</i>	koala		V	V	541
animals	mammals	Vombatidae	<i>Vombatus ursinus</i>	common wombat		NT		1
animals	reptiles	Elapidae	<i>Acanthophis antarcticus</i>	common death adder		V		1
plants	higher dicots	Apocynaceae	<i>Marsdenia coronata</i>	slender milkvine		V		19/19
plants	higher dicots	Lamiaceae	<i>Plectranthus habrophyllus</i>			E	E	11/11
plants	higher dicots	Myrtaceae	<i>Eucalyptus curtisii</i>	Plunkett mallee		NT		13/13
plants	higher dicots	Myrtaceae	<i>Melaleuca irbyana</i>			E		1/1
plants	higher dicots	Oleaceae	<i>Notelaea ipsviciensis</i>			E	CE	12/12
plants	higher dicots	Oleaceae	<i>Notelaea lloydii</i>	Lloyd's native olive		V	V	6/6

CODES

I - Y indicates that the taxon is introduced to Queensland and has naturalised.

Q - Indicates the Queensland conservation status of each taxon under the *Nature Conservation Act 1992*. The codes are Extinct in the Wild (PE), Endangered (E), Vulnerable (V), Near Threatened (NT), Least Concern (C) or Not Protected ().

A - Indicates the Australian conservation status of each taxon under the *Environment Protection and Biodiversity Conservation Act 1999*. The values of EPBC are Conservation Dependent (CD), Critically Endangered (CE), Endangered (E), Extinct (EX), Extinct in the Wild (XW) and Vulnerable (V).

Records – The first number indicates the total number of records of the taxon for the record option selected (i.e. All, Confirmed or Specimens).

This number is output as 99999 if it equals or exceeds this value. The second number located after the / indicates the number of specimen records for the taxon.

This number is output as 999 if it equals or exceeds this value.



Appendix B

Protected Matters Search Results



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 30/11/15 16:16:24

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

[Coordinates](#)

[Buffer: 10.0Km](#)



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	None
Wetlands of International Importance:	1
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	None
Listed Threatened Ecological Communities:	2
Listed Threatened Species:	54
Listed Migratory Species:	34

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	1
Commonwealth Heritage Places:	1
Listed Marine Species:	36
Whales and Other Cetaceans:	1
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Commonwealth Reserves Marine:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	3
Regional Forest Agreements:	None
Invasive Species:	35
Nationally Important Wetlands:	1
Key Ecological Features (Marine)	None

Details

Matters of National Environmental Significance

Wetlands of International Importance (Ramsar)	[Resource Information]
Name	Proximity
Moreton bay	20 - 30km upstream

Listed Threatened Ecological Communities

 [Resource Information]

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Name	Status	Type of Presence
Lowland Rainforest of Subtropical Australia	Critically Endangered	Community may occur within area
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland	Critically Endangered	Community likely to occur within area

Listed Threatened Species

 [Resource Information]

Name	Status	Type of Presence
Birds		
Anthochaera phrygia Regent Honeyeater [82338]	Critically Endangered	Foraging, feeding or related behaviour may occur within area
Botaurus poiciloptilus Australasian Bittern [1001]	Endangered	Species or species habitat likely to occur within area
Cyclopsitta diophthalma coxeni Coxen's Fig-Parrot [59714]	Endangered	Species or species habitat may occur within area
Dasyornis brachypterus Eastern Bristlebird [533]	Endangered	Species or species habitat likely to occur within area
Diomedea exulans antipodensis Antipodean Albatross [82269]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans exulans Tristan Albatross [82337]	Endangered	Species or species habitat may occur within area
Diomedea exulans gibsoni Gibson's Albatross [82271]	Vulnerable	Species or species habitat may occur within area
Diomedea exulans (sensu lato) Wandering Albatross [1073]	Vulnerable	Species or species habitat may occur within area
Erythrotriorchis radiatus Red Goshawk [942]	Vulnerable	Species or species habitat known to occur within area
Geophaps scripta scripta Squatter Pigeon (southern) [64440]	Vulnerable	Species or species

Name	Status	Type of Presence
Grantiella picta Painted Honeyeater [470]	Vulnerable	habitat may occur within area Species or species habitat may occur within area
Lathamus discolor Swift Parrot [744]	Endangered	Species or species habitat likely to occur within area
Macronectes giganteus Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Pachyptila turtur subantarctica Fairy Prion (southern) [64445]	Vulnerable	Species or species habitat likely to occur within area
Poephila cincta cincta Black-throated Finch (southern) [64447]	Endangered	Species or species habitat may occur within area
Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat likely to occur within area
Thalassarche cauta cauta Shy Albatross, Tasmanian Shy Albatross [82345]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta salvini Salvin's Albatross [82343]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta steadi White-capped Albatross [82344]	Vulnerable	Species or species habitat likely to occur within area
Thalassarche eremita Chatham Albatross [64457]	Endangered	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche melanophris impavida Campbell Albatross [82449]	Vulnerable	Species or species habitat may occur within area
Turnix melanogaster Black-breasted Button-quail [923]	Vulnerable	Species or species habitat likely to occur within area
Fish		
Epinephelus daemeli Black Rockcod, Black Cod, Saddled Rockcod [68449]	Vulnerable	Species or species habitat may occur within area
Insects		
Phyllodes imperialis smithersi Pink Underwing Moth [86084]	Endangered	Species or species habitat may occur within area
Mammals		
Chalinolobus dwyeri Large-eared Pied Bat, Large Pied Bat [183]	Vulnerable	Species or species habitat likely to occur within area

Name	Status	Type of Presence
Dasyurus hallucatus Northern Quoll [331]	Endangered	Species or species habitat may occur within area
Dasyurus maculatus maculatus (SE mainland population) Spot-tailed Quoll, Spotted-tail Quoll, Tiger Quoll (southeastern mainland population) [75184]	Endangered	Species or species habitat known to occur within area
Petrogale penicillata Brush-tailed Rock-wallaby [225]	Vulnerable	Species or species habitat known to occur within area
Phascolarctos cinereus (combined populations of Qld, NSW and the ACT) Koala (combined populations of Queensland, New South Wales and the Australian Capital Territory) [85104]	Vulnerable	Species or species habitat known to occur within area
Potorous tridactylus tridactylus Long-nosed Potoroo (SE mainland) [66645]	Vulnerable	Species or species habitat may occur within area
Pteropus poliocephalus Grey-headed Flying-fox [186]	Vulnerable	Roosting known to occur within area
Other		
Cycas ophiolitica [55797]	Endangered	Species or species habitat likely to occur within area
Plants		
Arthraxon hispidus Hairy-joint Grass [9338]	Vulnerable	Species or species habitat may occur within area
Bosistoa transversa Three-leaved Bosistoa, Yellow Satinheart [16091]	Vulnerable	Species or species habitat likely to occur within area
Cupaniopsis tomentella Boonah Tuckeroo [3322]	Vulnerable	Species or species habitat likely to occur within area
Notelaea ipsviciensis Cooneana Olive [81858]	Critically Endangered	Species or species habitat may occur within area
Notelaea lloydii Lloyd's Olive [15002]	Vulnerable	Species or species habitat likely to occur within area
Phaius australis Lesser Swamp-orchid [5872]	Endangered	Species or species habitat likely to occur within area
Phebalium distans Mt Berryman Phebalium [81869]	Critically Endangered	Species or species habitat may occur within area
Planchonella eerwah Shiny-leaved Condoo, Black Plum, Wild Apple [17340]	Endangered	Species or species habitat likely to occur within area
Plectranthus habrophyllus [64589]	Endangered	Species or species habitat likely to occur within area
Sophora fraseri [8836]	Vulnerable	Species or species habitat likely to occur within area
Thesium australe Austral Toadflax, Toadflax [15202]	Vulnerable	Species or species habitat likely to occur within area

Name	Status	Type of Presence
Reptiles		
Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Coeranoscincus reticulatus Three-toed Snake-tooth Skink [59628]	Vulnerable	Species or species habitat may occur within area
Delma torquata Collared Delma [1656]	Vulnerable	Species or species habitat likely to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Furina dunmali Dunmall's Snake [59254]	Vulnerable	Species or species habitat may occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area

Listed Migratory Species [\[Resource Information \]](#)

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Migratory Marine Birds		
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Diomedea antipodensis Antipodean Albatross [64458]	Vulnerable*	Species or species habitat may occur within area
Diomedea dabbenena Tristan Albatross [66471]	Endangered*	Species or species habitat may occur within area
Diomedea exulans (sensu lato) Wandering Albatross [1073]	Vulnerable	Species or species habitat may occur within area
Diomedea gibsoni Gibson's Albatross [64466]	Vulnerable*	Species or species habitat may occur within area
Macronectes giganteus Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Thalassarche cauta (sensu stricto) Shy Albatross, Tasmanian Shy Albatross [64697]	Vulnerable*	Species or species habitat may occur within

Name	Threatened	Type of Presence area
Thalassarche eremita Chatham Albatross [64457]	Endangered	Species or species habitat may occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable*	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche salvini Salvin's Albatross [64463]	Vulnerable*	Species or species habitat may occur within area
Thalassarche steadi White-capped Albatross [64462]	Vulnerable*	Species or species habitat likely to occur within area
Migratory Marine Species		
Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Manta alfredi Reef Manta Ray, Coastal Manta Ray, Inshore Manta Ray, Prince Alfred's Ray, Resident Manta Ray [84994]		Species or species habitat may occur within area
Manta birostris Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray, Oceanic Manta Ray [84995]		Species or species habitat may occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area
Orcaella brevirostris Irrawaddy Dolphin [45]		Species or species habitat known to occur within area
Migratory Terrestrial Species		
Cuculus optatus Oriental Cuckoo, Horsfield's Cuckoo [86651]		Species or species habitat may occur within area
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat known to occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Monarcha melanopsis Black-faced Monarch [609]		Species or species

Name	Threatened	Type of Presence
Monarcha trivirgatus Spectacled Monarch [610]		habitat known to occur within area Species or species habitat known to occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species habitat known to occur within area
Rhipidura rufifrons Rufous Fantail [592]		Species or species habitat known to occur within area

Migratory Wetlands Species

Ardea alba Great Egret, White Egret [59541]		Breeding known to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Gallinago hardwickii Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area

Other Matters Protected by the EPBC Act

Commonwealth Land

[\[Resource Information \]](#)

The Commonwealth area listed below may indicate the presence of Commonwealth land in this vicinity. Due to the unreliability of the data source, all proposals should be checked as to whether it impacts on a Commonwealth area, before making a definitive decision. Contact the State or Territory government land department for further information.

Name
Defence - GREENBANK TRAINING AREA

Commonwealth Heritage Places

[\[Resource Information \]](#)

Name	State	Status
Natural		
Greenbank Military Training Area (part)	QLD	Listed place

Listed Marine Species

[\[Resource Information \]](#)

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Birds		
Anseranas semipalmata Magpie Goose [978]		Species or species habitat may occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba Great Egret, White Egret [59541]		Breeding known to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Cuculus saturatus Oriental Cuckoo, Himalayan Cuckoo [710]		Species or species

Name	Threatened	Type of Presence
Diomedea antipodensis Antipodean Albatross [64458]	Vulnerable*	habitat may occur within area Species or species habitat may occur within area
Diomedea dabbenena Tristan Albatross [66471]	Endangered*	Species or species habitat may occur within area
Diomedea exulans (sensu lato) Wandering Albatross [1073]	Vulnerable	Species or species habitat may occur within area
Diomedea gibsoni Gibson's Albatross [64466]	Vulnerable*	Species or species habitat may occur within area
Gallinago hardwickii Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area
Haliaeetus leucogaster White-bellied Sea-Eagle [943]		Species or species habitat known to occur within area
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat known to occur within area
Lathamus discolor Swift Parrot [744]	Endangered	Species or species habitat likely to occur within area
Macronectes giganteus Southern Giant Petrel [1060]	Endangered	Species or species habitat may occur within area
Macronectes halli Northern Giant Petrel [1061]	Vulnerable	Species or species habitat may occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Monarcha melanopsis Black-faced Monarch [609]		Species or species habitat known to occur within area
Monarcha trivirgatus Spectacled Monarch [610]		Species or species habitat known to occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species habitat known to occur within area
Pachyptila turtur Fairy Prion [1066]		Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Species or species habitat known to occur within area
Rhipidura rufifrons Rufous Fantail [592]		Species or species habitat known to occur

Name	Threatened	Type of Presence within area
Rostratula benghalensis (sensu lato) Painted Snipe [889]	Endangered*	Species or species habitat likely to occur within area
Thalassarche cauta (sensu stricto) Shy Albatross, Tasmanian Shy Albatross [64697]	Vulnerable*	Species or species habitat may occur within area
Thalassarche eremita Chatham Albatross [64457]	Endangered	Species or species habitat may occur within area
Thalassarche impavida Campbell Albatross, Campbell Black-browed Albatross [64459]	Vulnerable*	Species or species habitat may occur within area
Thalassarche melanophris Black-browed Albatross [66472]	Vulnerable	Species or species habitat may occur within area
Thalassarche salvini Salvin's Albatross [64463]	Vulnerable*	Species or species habitat may occur within area
Thalassarche steadi White-capped Albatross [64462]	Vulnerable*	Species or species habitat likely to occur within area

Reptiles

Caretta caretta Loggerhead Turtle [1763]	Endangered	Species or species habitat known to occur within area
Chelonia mydas Green Turtle [1765]	Vulnerable	Species or species habitat known to occur within area
Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768]	Endangered	Species or species habitat known to occur within area
Eretmochelys imbricata Hawksbill Turtle [1766]	Vulnerable	Species or species habitat known to occur within area
Lepidochelys olivacea Olive Ridley Turtle, Pacific Ridley Turtle [1767]	Endangered	Species or species habitat known to occur within area
Natator depressus Flatback Turtle [59257]	Vulnerable	Species or species habitat known to occur within area

Whales and other Cetaceans

Name	Status	Type of Presence
Mammals		
Orcaella brevirostris Irrawaddy Dolphin [45]		Species or species habitat known to occur within area

[Resource Information]

Extra Information

State and Territory Reserves [\[Resource Information \]](#)

Name	State
Mount Perry 1	QLD
Stewartdale	QLD
White Rock	QLD

Invasive Species [\[Resource Information \]](#)

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

Name	Status	Type of Presence
------	--------	------------------

Birds

Acridotheres tristis Common Myna, Indian Myna [387]		Species or species habitat likely to occur within area
Anas platyrhynchos Mallard [974]		Species or species habitat likely to occur within area
Carduelis carduelis European Goldfinch [403]		Species or species habitat likely to occur within area
Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Lonchura punctulata Nutmeg Mannikin [399]		Species or species habitat likely to occur within area
Passer domesticus House Sparrow [405]		Species or species habitat likely to occur within area
Streptopelia chinensis Spotted Turtle-Dove [780]		Species or species habitat likely to occur within area
Sturnus vulgaris Common Starling [389]		Species or species habitat likely to occur within area

Frogs

Rhinella marina Cane Toad [83218]		Species or species habitat likely to occur within area
--------------------------------------	--	--

Mammals

Bos taurus Domestic Cattle [16]		Species or species habitat likely to occur within area
Canis lupus familiaris Domestic Dog [82654]		Species or species habitat likely to occur within area
Equus caballus Horse [5]		Species or species habitat likely to occur within area
Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area

Name	Status	Type of Presence
Feral deer Feral deer species in Australia [85733]		Species or species habitat likely to occur within area
Lepus capensis Brown Hare [127]		Species or species habitat likely to occur within area
Mus musculus House Mouse [120]		Species or species habitat likely to occur within area
Oryctolagus cuniculus Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
Rattus norvegicus Brown Rat, Norway Rat [83]		Species or species habitat likely to occur within area
Rattus rattus Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
Sus scrofa Pig [6]		Species or species habitat likely to occur within area
Vulpes vulpes Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
Anredera cordifolia Madeira Vine, Jalap, Lamb's-tail, Mignonette Vine, Anredera, Gulf Madeiravine, Heartleaf Madeiravine, Potato Vine [2643]		Species or species habitat likely to occur within area
Cabomba caroliniana Cabomba, Fanwort, Carolina Watershield, Fish Grass, Washington Grass, Watershield, Carolina Fanwort, Common Cabomba [5171]		Species or species habitat likely to occur within area
Chrysanthemoides monilifera Bitou Bush, Boneseed [18983]		Species or species habitat may occur within area
Eichhornia crassipes Water Hyacinth, Water Orchid, Nile Lily [13466]		Species or species habitat likely to occur within area
Genista monspessulana Montpellier Broom, Cape Broom, Canary Broom, Common Broom, French Broom, Soft Broom [20126]		Species or species habitat likely to occur within area
Lantana camara Lantana, Common Lantana, Kamara Lantana, Large-leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White Sage, Wild Sage [10892]		Species or species habitat likely to occur within area
Opuntia spp. Prickly Pears [82753]		Species or species habitat likely to occur within area
Parkinsonia aculeata Parkinsonia, Jerusalem Thorn, Jelly Bean Tree, Horse Bean [12301]		Species or species habitat likely to occur within area
Parthenium hysterophorus Parthenium Weed, Bitter Weed, Carrot Grass, False Ragweed [19566]		Species or species habitat likely to occur within area
Salix spp. except S.babylonica, S.x calodendron & S.x reichardtii Willows except Weeping Willow, Pussy Willow and Sterile Pussy Willow [68497]		Species or species habitat likely to occur

Name	Status	Type of Presence within area
Salvinia molesta Salvinia, Giant Salvinia, Aquarium Watermoss, Kariba Weed [13665]		Species or species habitat likely to occur within area
Senecio madagascariensis Fireweed, Madagascar Ragwort, Madagascar Groundsel [2624]		Species or species habitat likely to occur within area
Solanum elaeagnifolium Silver Nightshade, Silver-leaved Nightshade, White Horse Nettle, Silver-leaf Nightshade, Tomato Weed, White Nightshade, Bull-nettle, Prairie-berry, Satansbos, Silver-leaf Bitter-apple, Silverleaf-nettle, Trompillo [12323]		Species or species habitat likely to occur within area
Reptiles		
Hemidactylus frenatus Asian House Gecko [1708]		Species or species habitat likely to occur within area

Nationally Important Wetlands	[Resource Information]
Name	State
Greenbank Army Training Area C	QLD

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

For species where the distributions are well known, maps are digitised from sources such as recovery plans and detailed habitat studies. Where appropriate, core breeding, foraging and roosting areas are indicated under 'type of presence'. For species whose distributions are less well known, point locations are collated from government wildlife authorities, museums, and non-government organisations; bioclimatic distribution models are generated and these validated by experts. In some cases, the distribution maps are based solely on expert knowledge.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-27.6906 152.89956

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [-Office of Environment and Heritage, New South Wales](#)
- [-Department of Environment and Primary Industries, Victoria](#)
- [-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [-Department of Environment, Water and Natural Resources, South Australia](#)
- [-Parks and Wildlife Commission NT, Northern Territory Government](#)
- [-Department of Environmental and Heritage Protection, Queensland](#)
- [-Department of Parks and Wildlife, Western Australia](#)
- [-Environment and Planning Directorate, ACT](#)
- [-Birdlife Australia](#)
- [-Australian Bird and Bat Banding Scheme](#)
- [-Australian National Wildlife Collection](#)
- Natural history museums of Australia
- [-Museum Victoria](#)
- [-Australian Museum](#)
- [-South Australian Museum](#)
- [-Queensland Museum](#)
- [-Online Zoological Collections of Australian Museums](#)
- [-Queensland Herbarium](#)
- [-National Herbarium of NSW](#)
- [-Royal Botanic Gardens and National Herbarium of Victoria](#)
- [-Tasmanian Herbarium](#)
- [-State Herbarium of South Australia](#)
- [-Northern Territory Herbarium](#)
- [-Western Australian Herbarium](#)
- [-Australian National Herbarium, Atherton and Canberra](#)
- [-University of New England](#)
- [-Ocean Biogeographic Information System](#)
- [-Australian Government, Department of Defence Forestry Corporation, NSW](#)
- [-Geoscience Australia](#)
- [-CSIRO](#)
- Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.




Appendix C

Curricula Vitae – Pen Port



David Havill: Senior Ecologist

David Havill has significant practical experience in the areas of ecological site assessments (flora and fauna), weed management programs, large scale revegetation projects, wetland rehabilitation and waterway restoration. He has a strong understanding of the intricate workings of the Vegetation Management Act 1999 and the complex codes and policies which influence site vegetation constraints.



David's expertise relates to the on-site identification and spatial mapping of fauna and flora species including endangered, rare and vulnerable plants and animals. He has an accurate understanding of site survey processes and standards developed by the State and Commonwealth Governments. This provides the ability to challenge the various inaccuracies that occur within broad scale vegetation mapping developed by these Government agencies.

David works closely with our in house team of GIS, environmental planning, and landscape rehabilitation specialists to document findings of ecological survey and prepare targeted restoration and rehabilitation strategies. He has a strong understanding of construction techniques associated with development projects and has the ability to prepare practical flora and fauna management plans to assist in guiding the construction process within sensitive areas.

Qualifications

Bachelor of Applied Science (Natural Systems and Wildlife Management), University of Queensland (1998)

Angela Little: Ecologist

Angela is a member of our Environmental Management team, with an academic background in the fields of marine and environmental science, and significant experience within the university and government research setting. Angela's environmental management experience is widespread, ranging from GIS support, ecological assessment in aquatic and terrestrial environments, compliance assessments for state departments, and reporting to meet regulatory requirements for resource sector, infrastructure and land development projects.



Angela has substantial technical expertise in water quality monitoring and baseline assessments, which includes equipment calibration, field sampling, and data management. Her recent completion of a qualification in Environment and Sustainability has enabled Angela to develop skills in community engagement.

Qualifications

Graduate Certificate in Environment, [Griffith University](#) (2013)

Bachelor of Science with Honours (Marine Science) Class I, [James Cook University](#) (2004)



Maree Clancy: Ecologist

Maree has extensive ecological field and desktop research experience gained while working in the forestry industry and with the Australian Koala Foundation. In previous roles she assisted with quarterly and annual reporting of rehabilitation/revegetation works at residential development reserves, habitat translocation sites and the Bruce Highway upgrade project, and also with annual fauna surveying and reporting on various projects. She has a wealth of experience with preliminary desktop assessments of potential species at survey sites and the identification of flora and fauna species present during surveys.



At the Australian Koala Foundation, Maree was involved in the Koala habitat mapping project which included the use of GIS and determining habitat values for regional ecosystems and mosaics based on canopy species rankings and percentage composition.

Maree has additional skills in native seed propagation and growing of seedlings for large scale revegetation and farm forestry projects, ongoing monitoring of propagation methods and plant health status and adaptive approaches to improving methods.

Qualifications

Bachelor of Environmental Science, [University of the Sunshine Coast](#) (2014)



Appendix D

Species Lists



Site Flora - Transect Meander Results													
Species	Common Name	Transect 1	Transect 2	Transect 3	Transect 4	Transect 5	Transect 6	Transect 7	Transect 8	Transect 9	Transect 10	Transect 11	Transect 12
HERB													
<i>Brunoniella australis</i>	Blue Trumpet							"	"	"	"		
<i>Chrysocephalum apiculatum</i>	Yellow Buttons	"		"	"		"	"	"	"	"	"	"
<i>Commelina diffusa</i>	Wandering Jew			"		"	"		"	"	"	"	"
<i>Crassula sieberiana</i>	Australian Crassula						"						
<i>Einadia nutans</i>	Einadia						"						
<i>Glossocardia bidens</i>	Native Cobbler's Pegs											"	"
<i>Lobelia purpurascens</i>	White Root	"				"	"	"	"	"		"	"
<i>Oxalis corniculata</i>	Yellow Wood-sorrel			"			"	"	"	"	"	"	"
<i>Murdannia graminea</i>	Slug Herb								"	"			
<i>Phyllanthus virgatus</i>	Phyllanthus		"		"			"	"	"		"	"
<i>Plectranthus parviflorus</i>	Plectranthus			"						"			
<i>Poranthera microphylla</i>	Poranthera						"						
<i>Hybanthus stellarioides</i>	Spade Flower							"			"		
<i>Wahlenbergia gracilis</i>	Small-flowered Bluebell			"		"		"	"	"	"	"	"
Total Number of Herbs Recorded		2	1	5	2	2	5	5	5	6	3	5	5
VINES													
<i>Asparagus africanus</i>	Climbing Asparagus Fern						"						
<i>Cassytha glabella</i>	Dodder Laurel							"	"	"		"	"
<i>Eustrephus latifolius</i>	Wombat Berry	"	"	"	"				"	"		"	"
<i>Geitonoplesium cymosum</i>	Scrambling Lily							"			"		
<i>Glycine microphylla</i>	Glycine			"	"	"		"	"			"	
<i>Hardenbergia violacea</i>	Native Sarsaparilla		"							"		"	"
<i>Ipomoea cairica</i>	Mile-a-minute						"						
<i>Neonotonia wightii</i>	Glycine	"			"		"			"			"
<i>Parsonsia straminea</i>	Monkey Rope Vine					"							
<i>Passiflora suberosa</i>	Corky Passion Vine	"	"	"	"	"	"	"	"	"	"	"	"
<i>Smilax australis</i>	Barbed Wire Vine					"					"		

environmental management
protected plants survey report



<i>Thunbergia alata</i>	Black-eyed Susan							"					
<i>Vigna vexillata</i>	Wild Cowpea	"	"					"	"	"	"	"	"
Total Number of Vines Recorded		4	4	3	4	4	6	5	5	6	4	6	6
ORCHIDS / EPIPHYTES													
<i>Dockrillia linguiformis</i>	Tongue Orchid							"					
<i>Platyterium bifurcatum</i>	Elkhorn							"					
<i>Platyterium suberbum</i>	Staghorn							"					
Total Number of Orchids / Epiphytes Recorded		0	0	0	0	0	3	0	0	0	0	0	0
GROUND													
<i>Adiantum aethiopicum</i>	Maidenhair Fern							"					
<i>Ageratum houstonianum</i>	Blue Billygoat Weed	"							"		"		"
<i>Ambrosia artemisiifolia</i>	Annual Ragweed	"						"					
<i>Andropogon virginicus</i>	Whicky Grass												"
<i>Aristida sp.</i>	Many Head Wire Grass	"		"	"	"	"	"	"	"	"	"	"
<i>Asclepias curassavica</i>	Red-head Cotton Bush							"					
<i>Bidens pilosa</i>	Cobbler's Pegs	"			"			"		"	"	"	
<i>Cassytha pubescens</i>	Dodder Laurel							"					
<i>Cayratia clematidea</i>	Slender Grape							"					
<i>Centella asiatica</i>	Pennywort							"					
<i>Cheilanthes distans</i>	Bristle Cloak Fern	"	"	"	"	"	"	"	"	"	"	"	"
<i>Chloris gayana</i>	Rhodes Grass	"			"				"		"		
<i>Chloris virgata</i>	Feathertop Rhodes Grass	"								"			
<i>Conyza bonariensis</i>	Flaxleaf Fleabane	"						"		"			
<i>Cymbopogon refractus</i>	Barbed Wire Grass	"		"	"				"	"	"	"	"
<i>Cyperus polystachyos</i>	Bunchy Sedge	"	"	"	"			"	"	"		"	
<i>Dianella caerulea</i>	Blueberry Lilly	"	"	"	"	"	"	"	"	"	"	"	"
<i>Dianella longifolia</i>	Blueberry Lilly	"	"	"	"	"	"	"	"	"	"	"	"
<i>Drynaria rigidula</i>	Basket Fern							"					
<i>Entolasia stricta</i>	Wiry Panic	"	"	"	"	"	"	"	"	"	"	"	"
<i>Gahnia aspera</i>	Saw Sedge	"		"	"	"	"	"	"	"	"	"	"
<i>Goodenia rotundifolia</i>	Goodenia	"		"	"	"				"	"	"	"
<i>Heliotropium amplexicaule</i>	Blue Heliotrope							"		"			"
<i>Heteropogon contortus</i>	Black Spear Grass		"	"	"			"	"		"	"	"
<i>Eremophila debilis</i>	Winter Apple			"	"								
<i>Imperata cylindrica</i>	Blady Grass	"	"	"	"				"	"	"	"	"

environmental management
protected plants survey report



<i>Juncus usitatus</i>	Common Rush							"					"
<i>Lantana montevidensis</i>	Creeping Lantana			"				"		"	"	"	"
<i>Lepidosperma laterale</i>	Variable Sword Sedge	"		"		"		"		"	"		"
<i>Lomandra longifolia</i>	Mat Rush			"				"	"	"	"	"	"
<i>Lomandra multiflora</i>	Many-flowering Mat Rush	"			"	"		"	"	"	"		"
<i>Megathyrsus maximus</i>	Guinea Grass		"					"				"	"
<i>Melinis repens</i>	Red Natal Grass			"	"			"		"	"	"	"
<i>Panicum sp.</i>	Panicum	"	"	"	"			"		"	"	"	"
<i>Paspalum conjugatum</i>	Sourgrass	"						"	"	"		"	"
<i>Poa labillardieri</i>	Tussock Grass	"	"			"		"	"	"	"		"
<i>Pomax umbellata</i>	Pomax	"	"		"					"			
<i>Pteridium esculentum</i>	Bracken											"	"
<i>Senecio madagascariensis</i>	Fireweed	"				"		"					
<i>Sida cordifolia</i>	Flannel Weed			"						"			
<i>Solanum nigrum</i>	Blackberry Nightshade	"						"					
<i>Solanum seafortianum</i>	Brazilian Nightshade	"						"					
<i>Sporobolus pyramidalis</i>	Giant Rat's Tail Grass							"			"		"
<i>Themeda triandra</i>	Kangaroo Grass	"	"	"	"	"		"	"	"	"	"	"
<i>Urochloa mutica</i>	Para Grass							"					
<i>Xyris complanata</i>	Hat Pins									"			"
Total Number of Ground Layer Species Recorded		26	12	19	19	12	32	18	20	22	17	16	27
SHRUB													
<i>Acacia complanata</i>	Flat Stem Wattle				"					"			
<i>Acacia fimbriata</i>	Fringed Wattle	"	"	"	"	"	"	"	"	"	"	"	"
<i>Baccharis halimifolia</i>	Groundsel Bush							"					
<i>Breynia oblongifolia</i>	Coffee Bush							"	"		"	"	"
<i>Bursaria spinosa</i>	Black Thorn		"	"		"						"	
<i>Daviesia villifera</i>	Daviesia	"	"			"							
<i>Dodonaea lanceolata</i>	Hop Bush	"	"	"	"	"			"	"	"		
<i>Dodonaea triangularis</i>	Small-leaved Hop	"											
<i>Gomphocarpus physocarpus</i>	Balloon Cotton Bush	"						"				"	
<i>Grewia latifolia</i>	Dog's Balls		"		"	"							
<i>Jacksonia scoparia</i>	Dogwood	"	"					"	"	"	"	"	"
<i>Lantana camara</i>	Lantana	"		"	"	"	"	"	"		"	"	"
<i>Leucaena leucocephala</i>	Leucaena							"					

environmental management
protected plants survey report



<i>Leucopogon juniperinus</i>	Prickly Heath	"	"	"	"	"			"	"	"	"	"
<i>Ochna serrulata</i>	Ochna			"									
<i>Opuntia stricta</i>	Prickly Pear	"		"		"				"			"
<i>Persoonia sericea</i>	Persoonia	"	"	"		"				"	"	"	"
<i>Pultenaea euchila</i>	Orange Pultenaea	"	"	"	"	"				"		"	
<i>Schinus terebinthifolius</i>	Broadleaved Pepper Tree							"					
<i>Senna pendula</i>	Easter Cassia							"					
<i>Solanum mauritianum</i>	Wild Tobacco Tree							"					
<i>Solanum torvum</i>	Devil's Fig							"					
<i>Tithonia diversifolia</i>	Japanese Sunflower							"					
<i>Trema tomentosa</i>	Poison Peach			"		"							
<i>Xanthorrhoea johnsonii</i>	Forest Grass Tree	"	"	"	"	"				"	"	"	"
Total Number of Shrub Species Recorded		12	10	11	8	12	12	4	3	9	8	9	7
SUB-CANOPY													
<i>Acacia concurrens</i>	Black Wattle		"	"	"			"		"	"	"	"
<i>Acacia disparrima</i>	Hickory Wattle												
<i>Acacia leiocalyx</i>	Early Flowering Black Wattle	"		"	"	"		"	"	"	"	"	"
<i>Acacia salicina</i>	Sally Wattle	"	"	"		"							"
<i>Allocasuarina littoralis</i>	Black She-oak	"	"	"	"	"		"	"	"	"	"	"
<i>Allocasuarina torulosa</i>	Forest Oak							"					"
<i>Alphitonia excelsa</i>	Soap Tree	"	"	"	"	"		"	"	"	"	"	"
<i>Celtis sinensis</i>	Chinese Elm							"					
<i>Cinnamomum camphora</i>	Camphor Laurel							"					
<i>Cupaniopsis anacardioides</i>	Tuckeroo	"			"	"		"		"			"
<i>Glochidion ferdinandi</i>	Cheese Tree												"
<i>Jagera pseudorhus</i>	Foambark Tree							"				"	
<i>Lophostemon confertus</i>	Brush Box			"		"				"	"	"	"
<i>Lophostemon suaveolens</i>	Swamp Box				"	"		"	"	"	"	"	"
<i>Melaleuca quinquenervia</i>	Broad Leaf Paperbark							"					
<i>Melaleuca saligna</i>	Willow Bottlebrush					"		"					
<i>Melia azedarach</i>	White Cedar							"					
<i>Petalostigma pubescens</i>	Quinine Bush	"	"		"			"	"	"	"	"	"
<i>Schefflera actinophylla</i>	Umbrella Tree	"						"					
<i>Tecoma stans</i>	Yellow Bells							"					
Total Number of Sub-canopy Species Recorded		7	5	6	7	7	13	6	6	8	6	7	11

environmental management
protected plants survey report



CANOPY													
<i>Angophora leiocarpa</i>	Smooth Bark Apple	"	"		"		"	"	"	"	"	"	"
<i>Angophora woodsiana</i>	Rough Bark Apple						"						
<i>Corymbia citriodora</i>	Spotted Gum		"	"	"	"	"	"	"	"	"	"	"
<i>Corymbia henryi</i>	Large Leaf Spotted Gum	"	"	"	"	"	"	"	"	"	"	"	"
<i>Corymbia intermedia</i>	Pink Bloodwood			"		"	"	"	"	"	"	"	"
<i>Corymbia tessellaris</i>	Moreton Bay Ash						"						"
<i>Corymbia trachyphloia</i>	Brown Bloodwood	"	"		"			"		"	"		
<i>Eucalyptus acmenoides</i>	White Mahogany		"	"	"			"		"		"	"
<i>Eucalyptus crebra</i>	Narrow Leaf Ironbark												
<i>Eucalyptus fibrosa</i>	Broad Leaf Ironbark	"	"	"									
<i>Eucalyptus major</i>	Grey Gum						"		"				
<i>Eucalyptus micorcorcys</i>	Tallowood										"		
<i>Eucalyptus moluccana</i>	Gum Topped Box			"									
<i>Eucalyptus seeana</i>	Narrow Leaf Red Gum			"	"	"			"	"	"	"	"
<i>Eucalyptus siderophloia</i>	Grey Ironbark		"	"		"	"	"	"	"	"	"	"
<i>Eucalyptus tereticornis</i>	Forest Red Gum				"				"			"	"
Total Number of Canopy Species Recorded		4	7	8	6	5	8	7	8	7	6	8	8
Total Species Recorded		55	39	52	46	42	79	45	47	58	44	51	64

ATTACHMENT 3 –

Plectanthus habrophyllus Pre-clearance

Survey Notification

Date: 3 July 2017
Site: Springfield Rise
Client: Lendlease Communities
EPBC Ref: 2013/7057
SHG Ref: 7243
SHG Contact: Murray Saunders (07 3251 9444)

Attention: Ian Murray

Regional Development Manager, Communities
Level 4, Kings Gate,
King Street
Bowen Hills QLD 4006

Springfield Rise: V11 *Plectanthus habrophyllus* pre-clearance survey, 7002 Grande Avenue, Springfield (Lot 2 on SP291381)

Dear Ian,

This letter provides confirmation that the *Environmental Management Division* of **Saunders Havill Group** was engaged by **Lendlease Communities** to undertake a pre-clearance survey for *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) threatened flora species *Plectanthus habrophyllus* within the proposed clearing extent for the Village 11 (V11) Development Area – Phase 1 preliminary bulk earthworks area to meet Condition 6 of the EPBC Act approval (Ref: 2013/7057).

While *Plectanthus habrophyllus* populations were previously recorded in close proximity to the works area as part of the Spring Mountain EPBC survey by **Yurrah** (refer to **Attachment 2**), no *Plectanthus habrophyllus* specimens were recorded within the V11 Development Area clearing extent (refer to **Attachment 1** for a copy of the clearing extent).

The following provides relevant details of the survey:

Applicant: Lend Lease Communities (Springfield) Pty Ltd
Site Details: 7002 Grande Avenue, Springfield (Lot 2 on SP291381)
Development Area: Springfield Rise V11

***Plectanthus habrophyllus* Pre-Clearance Survey Results:**

Survey Completed by: David Havill (Senior Ecologist) & Maree Clancy (Ecologist)

Survey Completion Date: 14th February 2017

Was the survey undertaken in accordance with EPBC Act survey guidelines? Yes

Were any *Plectanthus habrophyllus* specimens identified within the clearing area? No




Kind regards,



Murray Saunders

Director – Saunders Havill Group

Attachment I –
Plectranthus habrophyllus Pre-clearance Survey Extent



Springfield - Villages 6 & 8

Flora Meandering Survey Transects

ISSUES:

Issue	Date	Description	Drawn	Checked
A	5/01/2016	Prelim Draft	TC	MS

Date | 5/01/2016
 Scale | 1:12,500 @ A3
 Coordinate System | GDA 1994 MGA Zone 56
 Projection | Transverse Mercator
 Client | Lend Lease
 Project | Springfield
 Address/RPD | Springfield Villages 6 to 8
 Sources | QLD GIS Layers (QLD Gov. Info Services 2015), Aerial (Nearmap, 2015)

Plan 1

SHG File
 7522 E 01 Flora Meandering Survey B


THESE PLANS HAVE BEEN PREPARED FOR THE EXCLUSIVE USE OF THE CLIENT. SAUNDERS HAVILL GROUP CANNOT ACCEPT RESPONSIBILITY FOR ANY USE OF OR RELIANCE UPON THE CONTENTS OF THESE DRAWINGS BY ANY THIRD PARTY.
 CONFIRM ALL DIMENSIONS ON SITE PRIOR TO CONSTRUCTION AND DO NOT SCALE FROM THE DRAWINGS. ALL DIMENSIONS ARE IN METRES. ANY DISCREPANCIES SHOULD BE CLARIFIED IN WRITING WITH SAUNDERS HAVILL GROUP PRIOR TO THE COMMENCEMENT OF WORK.
 PRIOR TO ANY DEMOLITION, EXCAVATION OR CONSTRUCTION ON SITE, THE RELEVANT AUTHORITY SHOULD BE CONTACTED FOR FURTHER UNDER-GROUND SERVICES AND DETAIL LOCATIONS OF ALL SERVICES.

Attachment 2 –


Plectranthus habrophyllus Surevy by Yurrah

CONCEPT MANAGEMENT PLAN


GPS Locations of <i>Plectranthus habrophyllus</i> populations UTM Zone 56 J		
ID	Latitude	Longitude
Plec 1	489651	6937126
Plec 2	489534	6937058
Plec 3	490045	6937140
Plec 4	488935	6937742
Plec 5	489700	6938233
Plec 6	489823	6937058



Approximate extent of *Plectranthus habrophyllus* sub-population (ID - Plec 5). Approximately 5 mature individuals within approximately 500m². 127m from development footprint to the south.




Approximate extent of *Plectranthus habrophyllus* ID - Plec 4. Approximately 5 mature individuals within approximately 400m². Population will require translocation into Linear Open Space. See Insert C. The road is located in this alignment to minimise earthworks within the linear open space and development areas. The southeast regional pipeline is located along the ridge and controls the level of the road as it crosses this linear open space area.



Scale - 1:20 000 @ A3

Approximate extent of *Plectranthus habrophyllus* ID - Plec 2. Approximately 5 mature individuals within approximately 200m². Development footprint, proposed residential, 5m to northwest.


Approximate extent of *Plectranthus habrophyllus* ID - Plec 1. Approximately 10 mature individuals within approximately 500m². Development footprint, proposed residential, 20m to northwest.



Scale 1:2000 @ A3

Approximate extent of *Plectranthus habrophyllus* ID - Plec 6. Approximately 20 mature individuals within approximately 500m². Population will require translocation into Linear Open Space to the east.

Approximate extent of *Plectranthus habrophyllus* ID - Plec 3. Approximately 50 mature individuals within approximately 3000m². Adjacent to residential development.



INSERT D

Scale 1:2000 @ A3

LEGEND

- E2 Precinct Boundary
- Proposed Development Layout**
 - Development footprint - use other than for conservation purposes
 - Linear Open Space - managed for conservation purposes
- Management Plan Core Conservation Areas - *Plectranthus habrophyllus* population location**
 - Where adjacent to an area identified for 'Interface Management' additional management actions required during clearing and construction. Refer Threatened Flora Management Plan Section 3.2.2. Refer Section 3.3 for ongoing habitat management.
 - In-situ population.
 - Receive area - translocated population.
- Management Plan Buffer Area**
 - Buffer Area overlapping development area. Considered detailed design required. Refer Threatened Flora Management Plan Section 3.2.1 for more information.
 - Buffer Area within Linear Open Space. Any Buffer Area adjacent an area identified for 'Interface Management' will require targeted management actions for protection of threatened flora during clearing and construction. Refer Threatened Flora Management Plan Section 3.2.2 for more information.
- Management Plan Additional Management Actions**
 - Approximate population extent of *P. habrophyllus* to be translocated. Refer Threatened Flora Management Plan Section 3.1 for actions.
 - Threatened flora Interface management required. Refer Threatened Flora Management Plan Section 3.3.1 for actions.

ATTACHMENT 4 – Fauna Spotter Catcher Pre-clearance WHIMP and WPMP

Fauna Spotter Catcher Wildlife and Habitat Impact Mitigation Plan

Springfield Rise – Village 11
Spring Mountain, Queensland
Report prepared for Shadforths Civil Contractors



Report prepared by
QLD Fauna Consultancy Pty Ltd
Phone: (07) 3376 9780
Fax: (07) 3376 9740
Email: fauna@qfc.com.au

Date:	04/07/17
Title:	Fauna Spotter Catcher Wildlife and Habitat Impact Mitigation Plan Springfield Rise – Village 11, Spring Mountain, Queensland
Author/s:	Bryan Robinson, Camille Palmer, Ramona Rohwedder
Reviewed by:	Bryan Robinson
Status:	Final Report
Filed as:	QFC WHIMP Shadforth's Springfield Rise Village 11 June 2017.doc

The contents of this report and its appendices may not be used in any form by any party other than the Client and this project specifically. The reproduction, adaptation, use or communication of the information contained within this report may not be used without the written permission of Queensland Fauna Consultancy Pty Ltd. Neither the author/s nor the company (QFC Pty Ltd) accepts any liability or responsibility for the unauthorised use of any part of this document.

Contents

1. Introduction	4
1.1 Project Background	4
1.2 Project Location and Site Description	4
1.3 Current Permits and Authorities	6
2. Mitigation Strategies	7
2.1 Fauna Spotter	7
2.2 Clearing Methodologies	7
2.3 Fauna Fencing.....	7
2.4 Felling Procedures	8
2.5 Macropods.....	8
2.6 Aquatic Fauna.....	8
2.7 General Terrestrial and Arboreal Fauna.....	9
2.8 EVNT Fauna	9
3. Wildlife Capture & Removal Plan	12
4. Wildlife Contingency Plan	17
4.1 Basic Wildlife Care	17
4.2 First Aid.....	20
4.3 Euthanasia	21
5. Wildlife Storage & Housing Plan.....	22
6. Wildlife Release & Disposal Plan	24
7. Post Works Impact Minimisation	25
8. Assessment, Conclusion and Fauna Management Recommendations	26
9. References.....	27
10. Appendix A: Intended stages of clearing.....	28
11. Appendix B: Intended Release Site for Wildlife.....	29

1. Introduction

1.1 Project Background

Queensland Fauna Consultancy Pty Ltd has been engaged by Shadforth's Civil Contractors to prepare a Fauna Spotter Catcher Wildlife and Habitat Impact Mitigation Plan for Village 11 as part of the Springfield Rise Project, Spring Mountain, Queensland.

The objective of this report is to summarise the existing fauna values presented in the Fauna Spotter Catcher Pre-clearance Survey and Wildlife Protection and Management Plan (WPMP) and assign mitigatory strategies applicable to probable species likely to be encountered during the clearing of identified habitats throughout or within specific localities of the site. Fauna species both common and of elevated conservation value have been considered within the parameters of onsite investigations and, where provided to QFC, include review of current fauna and floristic reports that may influence the assemblages expected to utilise the microhabitats evident within the site.

This review encompasses species identified under the provisions of the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* and the Queensland *Nature Conservation Act 1992*. Further consideration is given, where applicable, to species of iconic, cultural and/or regional significance identified under commonwealth, state or local planning instruments aimed at the persistence of biodiversity values within the area.

1.2 Project Location and Site Description

Village 11 is centrally located within the Springfield Rise precinct and adjoins the previously cleared areas of Village 12 to the north-east and Village 15 to the north-west. Village 11 will adjoin the conservation area to the south of the precinct.

Existing features exhibit a woodland vegetative complex with drainage features present due to an undulating topography. Dominant trees species across several vegetation types include *Corymbia henryi*, *C. citriodora*, *Eucalyptus crebra*, *E. siderophloia*, *Lophostemon confertus* and *Angophora leiocarpa*.



Map 1: Project Location

(Image extracted from Springfield Rise at Spring Mountain Concept Masterplan, LandPartners 2016)

2. Mitigation Strategies

2.1 Fauna Spotter

It is advised that all identified fauna habitats onsite be inspected by a licensed Fauna Spotter prior to vegetation clearing, and all vegetation removal activities be supervised during the clearing process.

2.2 Clearing Methodologies

In accordance to the *Nature Conservation (Koala) Conservation Plan 2006 and Management Program 2006-2016* the following sequential clearing conditions are required to be adhered to:

- Clearing of trees is carried out in a way that ensures koalas living in or near the area being cleared (the clearing site) have enough time to move out of the clearing site without human intervention, including in particular, for a clearing site with an area of more than 6ha, by:
 - Carrying out the clearing in stages; and
 - Ensuring not more than the following is cleared in any one stage:
 - for a clearing site with an area of 6 ha or less—50 percent of the site’s area;
 - for a clearing site with an area of more than 6ha—3ha or 3 percent of the site’s area, whichever is the greater; and
 - Ensuring that between each stage there is at least one period of 12 hours that starts at 6 p.m. on a day and ends at 6 a.m. on the following day, during which no trees are cleared on the site;

In addition to these measures it is recommended that clearing activities be undertaken in a directional manner specified by the fauna spotter/catcher. This is done so as to reduce the likelihood of negative interactions between fauna and potential hazards e.g. roads and traffic, prevent isolation of fauna through habitat fragmentation, and to ensure that natural dispersal of wildlife away from clearing activities is not impeded.

A plan detailing the clearing direction within Village 11 can be viewed in Appendix A. Saunders Havill Group has proposed a plan detailing the direction and clearing locations within the Site Based Management Plan (Refer to Appendix A). This involves directional clearing towards the Spring Mountain Offset zone in the south, which has been earmarked as a safe haven zone for fauna movement and connectivity (Saunders Havill Group, 2016). This approach is supported by QFC as the most applicable response to managing highly mobile fauna.

2.3 Fauna Fencing

Temporary fencing has already been installed in various localities and will aid in minimizing the movement of large fauna including highly mobile macropods. The addition of further fauna fencing may be required if site conditions change and fauna considerations are presented by the fauna spotter catcher.

2.4 Felling Procedures

Trees identified as having potential fauna values (such as hollows, arboreal termitaria and exfoliating bark) will be clearly identified and subsequently marked for supervision during felling and inspected once felled. Efforts will be made to determine potentially occupant species by way of investigations for indicative signs (scats, scratchings and tracks) on the day(s) of clearing. Where no signs are found or potentially occupant species are undeterminable, machinery operators will be instructed to fell trees in a manner directed at minimising the potential risk of injury to fauna.

All identified microhabitats will be inspected via ground based observation and the direction of felling will be determined considering the safety of personnel, machinery and potentially occupant fauna. Felling procedures will see implementation of a soft felling technique specifically constructed by QFC to achieve minimal deceleration and impact upon felling. This will be achieved under direction of the Fauna Spotter present directly communicating with the plant operator(s).

2.5 Macropods

Macropods have been observed on adjacent sites and other signs including macropod scat and footprints have been located throughout the clearing precinct, as well as in areas adjacent to site.

The area of proposed clearing activities exhibits direct connectivity to notable habitat values along the southern border, therefore, if clearing commences in a directional and incremental fashion any macropods potentially encountered on site may move on of their own volition. In this event, it is recommended that clearing proceed as already recommended with continual reassessment by the onsite fauna spotters.

2.6 Aquatic Fauna

In the event aquatic dewatering activities will be required within the proposed clearing area; pooled water and drainage features will be inspected during terrestrial load reduction activities ahead of the clearing front. The following recommendations are made to mitigate impacts to potentially occupant fauna:

- Inspection of banks, peripheral vegetation and other immediate terrestrial microhabitats;
- Identification of potential fauna values including: logs, rocks, artificial structures, discarded rubbish and burrows;
- Targeted searched for frog egg deposition sites on debris, bank edges, water surface and vegetation.

2.7 General Terrestrial and Arboreal Fauna

Overall the site contains medium value refugial opportunities for arboreal and terrestrial fauna species. The species expected within the site are likely to primarily reflect common fauna assemblages for the region however provisions are proposed directly for common fauna and species of conservation significance.

It is advised that all identified fauna habitats onsite be inspected by a DEHP approved Fauna Spotter prior to vegetation clearing and all vegetation removal activities be supervised during the clearing process. Terrestrial load reduction activities will be conducted ahead of the clearing front where possible. Fauna captured will be relocated to adjacent habitat consistent with the life history requirements of the species requiring translocation.

2.8 EVNT Fauna

It is not envisaged that any species, listed under the provisions of the *Environment Protection and Biodiversity Conservation Act 1999* or the *Nature Conservation Act 1992*, other than those listed in the WPMP, will require specific management during vegetation clearing activities.

However, specific management for those identified EVNT species will include targeted investigations immediately prior to vegetation removal activities on each day of clearing and subsequently whilst clearing takes place. Preliminary investigations will be supported by additional monitoring applied during clearing activities with a designated fauna spotter operating with each machine actively involved in vegetation or identified habitat disturbance. These should include the following:

Koala:

As favoured Koala food trees on site exceed a diameter of 100mm at 1.3 metres from the ground, requirements under the Koala Plan's 'Koala Habitat Area' provisions trigger the need for inspection and monitoring during vegetation clearing by a qualified Fauna Spotter.

Historically known to occur within the area the Koala will feature highly in daily search efforts with a dedicated and detailed methodology employed as follows:

- Pre clearing (preliminary) investigations to be conducted specifically for Koala detection by one experienced fauna spotter a minimum half hour prior to works each day. The investigation will embrace all designated clearing zones identified for that day inclusive of a 25 metre buffer around that zone;
- Once clearing commences a fauna spotter will accompany each machine providing continuous verification of habitat values and potential identification of undetected koalas ahead of operating plant. This will also account for potentially transient Koalas that may enter the site after preliminary investigations are complete.

Direct observational methodology will include the following components

- Use of binoculars to inspect the crown, forks and trunk of trees for individuals currently occupying the site;
- 'Drip zone' searches at the base of known food trees for the presence of scats to a radius equal to that of the crown of individual trees;
- Inspection of trunks for scratchings indicative of use by Koalas;
- Repeat observations made of single trees from numerous angles at repeated times throughout the clearing activities by the assigned fauna spotter.

In the event a Koala is detected, the Fauna Spotter will determine the appropriate course of action with exclusion zones implemented and alterations to the clearing plan discussed with the Site Supervisor. Once defined, these directions will be communicated to the plant operators and clearing will proceed in accordance with the recommendations made.

Changes to Koala management strategies highlighted in the *Nature Conservation (Koala) Conservation Plan 2006 and Management Program 2006-2016* have resulted in particular conditions placed on vegetation clearance involving the removal of Koala food trees. These provisions entail an increased responsibility by developers and land clearance operators alike to ensure the welfare of potentially present Koalas in areas identified as having significance for the persistence of this species.

Where significance under planning instruments is assigned provisions may include the restriction of all clearance that directly interferes with any tree a Koala is residing in or surrounding trees that, when felled, may impact on the crown of the host tree. Koalas are to leave via their own volition through a corridor designated by the Fauna Spotter to the closest remaining suitable habitat.

Throughout this time the Koala may not be interfered with by any means unless special dispensation has been sought through the appropriate government body or where the Koala is evidently in a state of compromised health. Only when Koalas have vacated a tree can clearance operations include the identified host tree and surrounding vegetation which composes the established exclusion zone. Recommendations made by the Fauna Spotter on site will embrace these provisions.

Grey-headed Flying Fox:

Although no Flying Fox camps or roosts were noted during the site survey, the transient nature of this species and the abundance of available feeding resources would see probability for the species to intermittently utilise the site.

The following recommendations are made for management of potentially occurring Grey-headed Flying Fox:

- Daily Inspection of trees assigned for removal be conducted to detect potential roosting Flying Foxes;
- Trees found to contain roosting Flying Foxes to be left standing and re assessed at the end of each days clearing. Being a transient species, the disturbance associated by the surrounding clearing is likely to see individuals fly off via its own volition come nightfall and not return the following morning, thus negating the need for direct disturbance.

Powerful Owl:

The site contains hollowing bearing trees with the potential to support nesting localities for the Powerful Owl. Diurnal roosting opportunities are afforded however these are considered only moderately favourable. Feeding resources would be available as highly targeted species such as glider and possum species are common throughout the region.

The following recommendations are made for management of potentially occurring Powerful Owl;

- Inspection daily of trees assigned for removal in areas of likely occurrence to detect potentially roosting birds;
- Identification of hollows exhibiting suitable dimensions for use as a nesting resource;
- Ground searches for casts and faecal accumulates indicative of the presence of Powerful Owl roosting and nesting sites;
- Implementation of a soft felling technique where trees are determined to have potential for occupancy.

Spotted-tail Quoll:

Although no dens or further evidence of Spotted-tail Quoll activity was detected during the survey, the species is known to occur historically in low densities in proximity to the site. Geomorphic structure and topography are considered favourable resulting in the following recommendations for further mitigation during the clearing activity:

- Inspection daily of identified geomorphic structure such as large boulders and rock accumulates, large hollow ground logs and log stock piles;
- Monitored dismantling of identified microhabitats by fauna spotters with machinery assistance.

Greater Glider:

The site contains hollow-bearing trees with the potential to support den localities for the Greater Glider. Suitable feeding resources are highly available given the availability of *Eucalyptus* leaves; on which the Greater Glider almost exclusively feeds on. The following recommendations are made for management of potentially occurring Greater Glider;

- Basal and drip zone searches for scats indicative of the presence of Greater Glider;
- Inspection daily of trees assigned for removal in areas of likely occurrence to detect Great Glider;
- Implementation of a soft felling technique where trees are determined to have potential for occupancy.

Collared Delma:

The presence of rocky habitat combined with *Eucalyptus* dominated woodlands presents known favorable habitat for the Collared Delma. The following recommendations are made for mitigation during clearing activity:

- Inspection daily of identified geomorphic structures including rocky outcrops, surface rock, leaf litter and bark exfoliates;
- Monitored dismantling of identified microhabitats by fauna spotters with machinery assistance.

3. Wildlife Capture & Removal Plan

Relocation of native fauna is a strategy that may be required during the course of developmental works to up-hold the project's required nature conservation, animal welfare and human safety objectives.

In all circumstance where native fauna are required to be relocated it must be done so, or under the direct supervision of, a suitably licensed fauna spotter/catcher. A summary of the fauna capture, handling and relocations strategies to be implemented by the fauna spotter/catcher for fauna groups deemed likely, or possible, to occur on site are presented in *Table 2*.

Table 2: Fauna capture, handling and relocation strategy table

Animal Group	Capture and handling	Relocation
Lizards Geckoes Dragons Monitors	<ul style="list-style-type: none"> Place one hand behind the head at the base of the quadrates and the other at the base of the tail behind the hind limbs; Be cautious when handling smaller skinks and legless lizards as they may discard their tail; Lizards and geckoes can be placed inside suitably sized calico bags In the case of large monitor lizards keep the animal's ventral surface directly away from the body with the tail between the upper arm and torso. Dragons and small monitors can be placed in suitably sized calico bags. Larger monitors to be placed in suitably sized crate 	<ul style="list-style-type: none"> Place the lizard head first into a suitable holding crate for later release. <ul style="list-style-type: none"> Dragons & monitors– release up trees or into heavy vegetation; Water dragons – in the vicinity of riparian areas; Skinks, Geckoes, Legless lizards – around creek margins.
Snakes	<ul style="list-style-type: none"> Due to their mobile nature, large snakes generally do not require to be handled or relocated, with the exception of slow moving species (i.e. pythons) or smaller species; Snakes should be identified and only moved if competent and safe to do so (see SOP006 Handling Venomous Snakes Procedure); Do not attempt to catch a snake if you're not competent; Injured snakes should be handled with suitable equipment. 	<ul style="list-style-type: none"> Release in suitable habitat e.g. along creek lines for python and tree snakes If feasible take them well away from clearance site to a suitable release location Release discreetly away from high density suburban areas
Small Mammals	<ul style="list-style-type: none"> Place a gloved hand around the whole animal in the case of small mammals (melomys or rats), Do not handle rodents by the tail as this will cause damage to the tail sheath Place the animal in calico bag in a cool place for later relocation. Minimise holding time to avoid animal gnawing through bags and escaping 	<ul style="list-style-type: none"> Release animal into area suitable to its habitat requirements. Ensure plenty of cover is available.

Animal Group	Capture and handling	Relocation
<p>Glider Family</p>	<ul style="list-style-type: none"> Place gloved hands around the animal at initial capture; Place the glider(s) into a calico bag or suitable animal crate ensuring family groups are kept together for all inclusive release; Place in a cool dry area during the day. When using calico bags ensure the bag is hung and well ventilated Where possible contain gliders within hollow by plugging openings with a towel or calico bag 	<ul style="list-style-type: none"> Release glider into habitat with natural hollows and canopy cover; When releasing a family group with more than one furred young (being carried on the back) either: <ul style="list-style-type: none"> Divide young between parents as a mother is unlikely to carry more than one young, Place young in elevated hollow with parents and allow them to move away in their own time. Place animal in bag at the base of the selected tree, opening the bag wide and allowing the animal to leave the bag when it is ready. Relocate hollow (with gliders inside) to suitable habitat and cover lightly with foliage so that the gliders can move away of their own accord and are protected from predators.
<p>Amphibians</p>	<ul style="list-style-type: none"> Amphibians should be handled only when necessary and handling times should be kept to a minimum to help prevent: <ul style="list-style-type: none"> Removal of the protective mucous layer covering the skin of amphibians; To prevent handling stress induced by changes in their body temperature; Risk of spreading pathogens and parasites. Amphibians from different sites need to be kept isolated from each other, and need to be kept in different containers or bags; Any dead or sick amphibians need to be quarantined from other amphibians. <p>Amphibians can be handled utilising one of the following methodologies:</p> <ul style="list-style-type: none"> Bare handed – ensure hands are sterilized before handling and free from lotions, sunscreen etc Gloves – disposable gloves desirable or disinfect gloves between handling different animals; Plastic bags – Single use lightweight plastic bags can be used to pick up and handle frogs; again plastic bags should be disposed of before handling amphibians form a different site. All staff should be knowledgeable and familiar with the <i>Interim Hygiene Protocol for Handling Amphibians – Technical Manual (DEHP)</i> 	<ul style="list-style-type: none"> Always ensure that amphibians are kept moist until release. This can include storing in a designated container with moist soil or toweling or in a wet calico bag; Release into suitable adjacent vegetation that is typical of the species requirements; Suitable release locations include riparian vegetation, low-lying wetlands, alongside creek lines, hollow logs, dams and ponds; Amphibians from different sites need to be released in separate locations; Disinfection procedures in relation to amphibians need to be followed.

Animal Group	Capture and handling	Relocation
Macropods	<ul style="list-style-type: none"> • Capture and restraint of macropods carries a high risk of injury and fatal hyperthermia/myopathy syndrome, and must not be performed by inexperienced personnel, or without appropriate equipment and sedation. • Capture and restraint of healthy macropods (other than pouch young) must be performed using sedation or anaesthesia due to the high risk of developmental myopathy, and other capture and restraint-associated conditions. Sedative and anaesthetic drugs may only be used under direct supervision of a registered veterinarian, or by appropriately licensed persons (Hanger & Nottidge, 2009). 	<ul style="list-style-type: none"> • Release animal into suitable to its habitat requirements. Ensure plenty of cover is available. • Macropods are to be released within the range of normal movement from their place of origin. E.g. a Kangaroo can be released within 100 km of its origin, based on its capacity to travel long distances. • Monitor animals to ensure adequate recovery if sedated.
Microbats	<ul style="list-style-type: none"> • Only vaccinated persons are to handle bats • If possible plug the hollow opening with a bag or towel and ask the operator to cut the hollow from the tree; • Always wear gloves when handling bats. • If not contained within a hollow, place bats inside a calico bag and hang upright in a cool place 	<ul style="list-style-type: none"> • Relocate hollow (with bats inside) to suitable habitat and cover lightly with foliage so that the bats can move away of their own accord and are protected from predators. • Bats not contained within a hollow should be released as late as possible at the end of the day.
Possums	<ul style="list-style-type: none"> • Use thick elbow length gloves when handling possums; • Try to grip the animal behind the head near the shoulder blades and around the tail so that you have control of the animal; • Keep fingers away from the mouth of the animal; • Keep the animal's body facing away at all times; • Transfer into a thick calico bag and then into a kitty crate. Place in a safe and shady place until you can relocate the animal. 	<ul style="list-style-type: none"> • Release the possum into habitat with adequate hollows and cover; • Place animal in bag at the base of a select tree, opening the bag and allow the animal to leave the bag when it is ready; • When releasing a Ringtail Possum mother with more than one furred young (being carried on her back) it is unlikely that she will carry both young if highly stressed; <ul style="list-style-type: none"> ○ Choose a smaller shrubby tree with vines or heavy foliage (so the adult can construct a drey easily) ○ Watch the adult ascend the tree, it is possible she will only carry one young and so any additional young may be pushed from her back ○ It may be necessary to take one or more of the young to a wildlife carer ○ If possible place mother and young in a suspended hollow, cover lightly with foliage and allow the animals to move on their own accord. This way the mother can ferry young one at a time to a more suitable location.

Animal Group	Capture and handling	Relocation
Birds	<ul style="list-style-type: none"> • Use gloves when handling larger birds • Use a towel to cover the bird and simultaneously restrain the bird and transfer into calico bag • With larger parrots and raptors, restrain head and legs and transfer into a kitty crate • Wrap chicks loosely in a towel and transfer to kitty crate, keep in a warm location. 	<ul style="list-style-type: none"> • Relocate adult birds in suitable habitat • Chicks should be referred to wildlife carer
Koalas	<p>Movement of Koalas is heavily legislated in South East Queensland. Koalas are not to be captured or relocated without the prior consent of Department Environment and Natural Resource Management (DERM). Koalas should be left to move away of their own volition and trees are not to be felled while a Koala remains in occupancy. See SOP003 Koala Management Procedure for further information.</p>	

4. Wildlife Contingency Plan

In the event sick, injured or orphaned protected animals are encountered during the course of the project they shall be administered to in accordance with the *Code of Practice Care of Sick, Injured or Orphaned Protected Animals in Queensland* under the *Nature Conservation Act 1992*.

The stages in which injuries or illness are described under the code are as follows:

Critical: Injuries or illnesses that are life-threatening; for example, an animal that has been struck by a car and has serious head injuries.

Serious: Injuries or illnesses that might reasonably be expected to cause moderate pain (but are not immediately life-threatening), and the animal is not showing obvious signs of distress or pain, or significantly reduced mental activity; for example an animal with a closed fracture but no other apparent injuries and that is alert and responsive.

Mild: The injuries or illness of an animal appear to cause little discomfort, pain or function loss and are not life-threatening (even without immediate vet treatment); for example superficial cuts, superficial bruising or orphaned animals suffering from mild dehydration.

4.1 Basic Wildlife Care

If wildlife requiring care are encountered by the fauna spotter/catcher they will be attended to in the manner set out by the guidelines provided in *Table 4*. Supplementary advice will be sought from a wildlife carer and/or veterinarian where required. QFC have previously utilised experienced local carer groups and vets. These are listed in *Table 3*.

Table 3: List of Local Vets & Wildlife Carer Groups

Vets			
Name	Location	Contact Number	Comments
RSPCA Wildlife Hospital	139 Wacol Station Road, Wacol	07 3426 9999	24 Hours/7days
Carers			
Name	Location	Contact Number	Comments
RSPCA Wildlife Hospital	139 Wacol Station Road, Wacol	07 3426 9999	24 Hours/7days
Ipswich Koala Protection Society	Ipswich	Ruth: 07 5464 6274 / 0419 760 127 Helen: 07 3282 5035 / 0417 604 761	Specialize in koalas however rescue all wildlife

Table 4: Basic Wildlife Care

Birds	Reptiles & Amphibians	Mammals
<p>Egg</p> <p>Viable eggs must be kept warm until transferred to a suitable wildlife carer. It is necessary that the orientation of the eggs be maintained as fixed embryos may be lost. Keep wrapped in a pouch and on a heat source (where available). An ideal temperature is between 25-27° (DEHP 2013); where possible attempt to identify the species so the carer can be informed as the management of eggs can vary in accordance with species and stage of development.</p>	<p>Egg</p> <p>Viable eggs must be kept warm and stable until transferred to a wildlife carer. It is necessary that the orientation of the eggs be maintained as fixed embryos may be lost. Keep wrapped in pouch or towel and place into an animal crate in a safe location.</p>	<p>Neonate</p> <p>Unfurred animals need to be kept warm until transferred to a carer. Place into a pouch and onto a heat pad. Ideal temperature is between 31-34°. 25-27° is appropriate in most other cases (DEHP 2013). Regularly check the animal to ensure it is not overheating by observing for obvious signs of distress (i.e. panting, very warm to the touch, red blotched skin). Adjust the temperature where required. Seek further advice from the carer if you are unsure.</p>
<p>Chick</p> <p>Make sure the animal is correctly identified as different species often have very different requirements. Place chicks into a pouch/towel onto a heat source maintained around 31-34° (only if they have not fledged) and keep in an animal crate until transferred to a carer.</p>	<p>Juvenile</p> <p>Place animals in a suitable lined crate and keep covered in a dark quiet place. Refer to the wildlife contact list in your QFC Folder for a carer who specialises in reptiles.</p>	<p>Juvenile</p> <p>Place into a lined crate and keep covered in a dark and quiet location.</p>
<p>Adult</p> <p>Keep adult birds in a lined animal crate or cage and covered in a quiet area.</p>	<p>Adult</p> <p>Place animals in a suitable lined crate and keep covered in a dark quiet place. Refer to the wildlife contact list in your QFC Folder for a carer who specialises in reptiles.</p>	<p>Adult</p> <p>Place into a lined crate and keep covered in a dark and quiet location.</p>
<p>Feeding</p> <p>Providing food and water is generally not required during short periods (2-3 hrs) though this should be reconsidered if animals need to be held longer. Consult the vet and/or carer for further advice on how to proceed.</p>	<p>Feeding</p> <p>Newly hatched reptiles may require feeding if kept overnight. Consult with QFC for further advice. Snakes and turtles will not require feeding but water should be made available.</p>	<p>Feeding</p> <p>Providing food and water is generally not required during short periods (2-3 hrs) though this should be reconsidered if animals need to be held longer. Consult the carer for further advice on how to proceed.</p>

4.2 First Aid

Animals suffering from serious injuries or illness encountered on the project should be passed on to veterinary care as soon as possible. In the interim a licensed fauna spotter/catcher can provide first aid for the animal and organise suitable transportation.

If a seriously sick or injured animal is encountered the fauna spotter/catcher should:

1. Keep the animal calm by placing into an animal crate and keeping it covered in a dark and quiet location. Isolate any nearby threats such as domestic animals or predators.
2. Quickly and thoroughly inspect the animal for trauma. If the injuries are not serious enough to require euthanasia administer the basic first aid as a minimum (but only if capable to do so)

Representative first aid that may be administered by a fauna spotter/catcher is provided in *Table 5*.

Table 5: Wildlife First Aid

Ailment	First Aid
Bleeding	Using material that is clean and sanitary, apply direct pressure to the affected area. Bandages can be used to hold material in place until vet treatment can be sought. Veterinarian treatment should be sought for further assistance as soon as possible.
Broken limbs	House the animal in a suitably sized animal crate with towels under the animal for comfort. Keep the crate covered and in a quiet location. Proceed to a veterinarian for further assistance as soon as possible.
Injured tails	House the animal in a suitably sized animal crate with towels under the animal for comfort. Keep the crate covered and in a quiet location. Proceed to a veterinarian for further assistance as soon as possible.
Concussions	House the animal in a suitably sized animal crate with towels under the animal for comfort. Keep the crate covered and in a quiet location. Proceed to a veterinarian for further assistance as soon as possible.

4.3 Euthanasia

Section 12 of the code details how to determine when euthanasia is required and how to euthanise animals ethically. The following standards as listed under the code are to be followed when assessing whether euthanasia is required:

- The euthanasia of wildlife where required is to be provided for by all wildlife rehabilitators;
- Euthanasia without exception is to be carried out when:
 - Significant pain or suffering is to be alleviated where it is not able to be managed by a vet;
 - Further treatment is **not** practical or recovery is **not** expected in a way in which the animal can be successfully rehabilitated back to the wild;
 - Resources are not available to provide appropriate care or an acceptable quality of life throughout the likely rehabilitation period.
- Animals that are suffering and have a poor prognosis for survival must be euthanised rather than left to die from the injury or illness. Failure to undertake appropriate action is a breach of the *Animal Care and Protection Act 2001*.
- Unless permission has been granted by the Department of Environment and Heritage Protection for the animal to enter the Queensland Species Management Plan (QSMP) or otherwise advised by the DEHP Wildlife Management Director, animals must be euthanised when:
 - An orphaned animal is not viable or likely to be rehabilitated;
 - No suitable release locations are available;
 - The ability for an animal to reproduce is lost due to an injury, disease or surgical procedure;
 - The ability to move freely or normally (i.e. run, climb, crawl, hop, fly or swim) is permanently impaired. Examples are: a missing or impaired limb, wing, foot or tail that would significantly impair the animal's ability to survive in the wild;
 - The ability to sense environment (i.e. see, smell, feel, taste or hear) is permanently impaired. For example: missing or injured organ such as an eye, ear or nose that would significantly impair the animal's ability to survive in the wild;
 - The ability to catch, find or handle food is permanently impaired;
 - Its advanced age renders it unlikely to survive in the wild.

5. Wildlife Storage & Housing Plan

For wildlife requiring storage, temporary housing and transportation to release sites and/or to a wildlife carer or veterinarian, guidelines set out in the Code of Practice and QFC's Animal Ethics Permit will be followed.

Dependent on the species of animal and condition of the animal, temporary storage and housing of animals will be as follows:

Calico bags: Calico bags will be used to temporarily house fauna such as snakes, lizards and small mammals (including microbats), Bags will range in size from 200mm x 200mm to 600mm x 1800mm. Bag selection will vary according to the size of animals to be placed in them. In the case of snakes a "hoop bag" may be used to facilitate capture. The hoop is approximately 500mm in diameter attached to a handle. The bag is placed around the hoop ensuring a greater area in which to pass the snake through into the bag.

Plastic holding tubs/containers/animal crate: Plastic holding tubs/containers/crates will be used to temporarily house fauna such as snakes, lizards, frogs, small mammals and birds (Plastic holding tubs/containers/crates will range in size from 150mm x 150mm x 120mm to 500mm x 400mm x

400mm. Plastic holding tubs/containers/crates selection will vary according to the size and number of animals to be placed in them.

In addition to this, material is used to line the tub/crate to ensure the animals won't lose its footing. This may include folded towels on the bottom of the crate or a fitted pad. These items are washed between each use to reduce the spread of disease/parasites.

Section 9 of the Code relates to how transportation of wildlife should be undertaken. The following will be adhered to when transporting wildlife to the vet and/or carer:

- Additional pain or distress of the animal is to be avoided;
- Wildlife should only be transported when necessary;
- Transport containers must be appropriate for the species (size, strength and behaviour of species being moved);
- Transport containers must be designed and maintained in a way as to:
 - Prevent injury;
 - Prevent escape;
 - Prevent rolling/tipping during transit;
 - Prevent damage to plumage (feathers);
 - Be hygienic;
 - Minimise stress and
 - Be suitably ventilated.

- Non-compatible species must not be transported in a manner which allows for visual or physical contact;
- Containers must be secured to prevent movement and provide protection from direct sunlight, wind and rain;

Venomous, dangerous or potentially disease transmitting animals must be clearly marked with warning labels (i.e. Caution –‘venomous snake’ or ‘live bat’) and be locked and secured.

6. Wildlife Release & Disposal Plan

Spring Mountain Forest Park lies to the south of Village 11 and contains similar habitat types suitable for species likely to be encountered when clearing.

With the exception of highly mobile species such as birds and macropods where natural relocation may occur, it will be necessary for the fauna spotter/catcher to translocate the majority of fauna found into suitable habitat within these areas. A map of the intended release site can be viewed in Appendix B.

In regard to all fauna capture and disposal activities conducted on the project the following records will be made:

- a. species;
- b. identification name or number;
- c. sex (M, F, or unknown);
- d. approximate age or age class (neonate, juvenile, sub-adult, adult);
- e. time and date of capture;
- f. method of capture;
- g. exact point of capture (GPS point);
- h. state of health;
- i. incidents associated with capture likely to affect the animal;
- j. veterinary intervention or treatments;
- k. time held in captivity;
- l. disposal (euthanasia, re-release, translocation etc);
- m. date and time of disposal;
- n. details of disposal (if released, exact point of release GPS);
- o. for released animals: distance in metres from point of capture to point of release.

7. Post Works Impact Minimisation

As the project area will be cleared of all vegetation, post works impact monitoring and/or impact minimisation is deemed not necessary. It is unlikely the vast majority of wildlife will return to the area as all habitat and foraging resources will be removed and habitat connectivity is also not present.

In the event that fauna is found on site post-works, it is recommended personnel contact QFC and a licensed and experienced wildlife consultant can be dispatched to remove and relocate the animal should it be necessary. QFC wildlife consultants are available 24/7 for fauna related call-outs in relation to this project.

It is recommended that if any fauna, such as Kangaroos and Wallabies, are noted in the wider area and appear distressed post-works that QFC be contacted to further assess the situation.

8. Assessment, Conclusion and Fauna Management Recommendations

A number of conclusions and recommendations are presented, with the specific intention of providing a comprehensive management structure to facilitate minimal impact to fauna during the clearing of vegetation and subsequent disturbance of habitats. The directives given by Fauna Spotter Catchers should embrace a “best practice” approach which includes implementation of proven specific management techniques for identified habitat types and compliance with legislation relevant to the activity.

Fauna management is presented here specific to EVNT fauna, general terrestrial and arboreal fauna and aquatic fauna. Although each is treated separately, overlap does occur within target techniques providing a comprehensive approach for target species of all conservation significance.

9. References

Anstis, M (2013) *Tadpoles and Frogs of Australia*, New Holland Publishers, Sydney.

Curtis, LK, Dennis, AJ, McDonald, KR, Kyne, PM & Debus, SJS (2012), *Queensland's Threatened Animals*, CSIRO Publishing, Victoria.

Department of Environment & Heritage Protection, *Interim Hygiene Protocol for Handling Amphibians*, Technical Manual

Hanger, J & Nottidge B (2009), *Draft Queensland Code of Practice for the Welfare of Wild Animals Affected by Land-Clearing and other Habitat Impacts and Wildlife Spotter/Catchers*, Australian Wildlife Hospital, Australia Zoo, Beerwah.

Queensland Environmental Protection Agency and Queensland Parks and Wildlife Service (2006). *Nature Conservation (Koala) Conservation Plan 2006 and Management Plan 2006 – 2016*. Queensland Government – Environmental Protection Agency.

Queensland Fauna Consultancy (2017) *Fauna Spotter Catcher Pre-clearance and Survey and Wildlife Protection & Management Plan, Springfield Rise – Village 11, Spring Mountain, Queensland*, (QFC FHA WPMP Shadforths Springfield Rise Village 11 June 2017.doc).

References for nomenclature

Cogger, H. (2000) *Reptiles & Amphibians of Australia*. 6th edition, Sydney: Reed New Holland.

Leiper, G., Glazebrook, J., Cox, D. and Rathie, K. (2008) *Mangroves to Mountains: a Field Guide to the Native Plants of South-east Queensland*, Browns Plains: Logan River Branch Society for Growing Australian Plants.

Menkhorst, K. & Knight, F. (2011) *A Field Guide to the Mammals of Australia*, 3rd edition, South Melbourne: Oxford University Press.

Simpson, K. and Day, N. (2004) *Field Guide to the Birds of Australia*, Camberwell: Penguin Group Australia.

Strahan, R. (Ed) (1995) *The Mammals of Australia*. Sydney: New Holland Publishers.

Wilson, S. (2005) *A Field Guide to Reptiles of Queensland*. Sydney: New Holland Publishers.

Queensland Museum (2007) *Wildlife of Greater Brisbane*, 2nd edition, Brisbane: Queensland Museum Publishers.



June 2017

Fauna Spotter Catcher Pre-clearance Survey and Wildlife Protection & Management Plan

Springfield Rise – Village 11
Spring Mountain, Queensland
Report prepared for Shadforths Civil Contractors



Report prepared by
QLD Fauna Consultancy Pty Ltd
Phone: (07) 3376 9780
Fax: (07) 3376 9740
Email: fauna@qfc.com.au

Date:	04/07/17
Title:	Fauna Spotter Catcher Pre-clearance and Habitat Values Survey Springfield Rise – Village 11, Spring Mountain, Queensland
Author/s:	Bryan Robinson, Ramona Rohwedder
Reviewed by:	Bryan Robinson
Field personnel:	Jonathan Pickvance
Status:	Final Report
Filed as:	QFC FHA Shadforths Springfield Rise Village 11 June 2017.doc

The contents of this report and its appendices may not be used in any form by any party other than the Client. The reproduction, adaptation, use or communication of the information contained within this report may not be used without the written permission of Queensland Fauna Consultancy Pty Ltd. Neither the author/s nor the company (QFC Pty Ltd) accepts any liability or responsibility for the unauthorised use of any part of this document.

Contents

1. Introduction	4
1.1 Project Background	4
1.2 Project Location and Site Description	4
1.3 Current Permits and Authorities	6
2. Methodology	7
2.1 Specific methodology for Koalas <i>Phascolarctos cinereus</i>	7
3. Findings	8
3.1 Terrestrial Habitat Features	8
3.2 Arboreal Habitat Features	12
3.3 Aquatic Habitat Features.....	17
3.4 Endangered, Vulnerable and Near Threatened (EVNT) Species	18
4. Fauna Impacts	20
5. Assessment and Conclusion	21
6. References.....	22
7. Appendix A: Koala Habitat Values	24
8. Appendix B: EPBC Act Protected Matters Report	25
9. Appendix C: Wildlife Online extract.....	36

1. Introduction

1.1 Project Background

Queensland Fauna Consultancy Pty Ltd has been engaged by Shadforth's Civil Contractors to conduct a Fauna Spotter Catcher Pre-clearance and Habitat Values Survey and present a subsequent report for Village 11 of the Springfield Rise development located at Spring Mountain, Queensland. The site location is presented in Map 1.

The objective of this report is to summarise the existing fauna values present and assign mitigatory strategies applicable to probable species likely to be encountered during the clearing of identified habitats throughout or within specific localities of the site. Fauna species both common and of elevated conservation value have been considered within the parameters of onsite investigations and, where provided to QFC, include review of current fauna and floristic reports that may influence the assemblages expected to utilise the micro habitats evident within the site.

This review encompasses species identified under the provisions of the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* and the Queensland *Nature Conservation Act 1992*. Further consideration is given, where applicable, to species of iconic, cultural and/or regional significance identified under commonwealth, state or local planning instruments aimed at the persistence of biodiversity values within the area.

1.2 Project Location and Site Description

Village 11 is centrally located within the Springfield Rise precinct and adjoins the previously cleared areas of Village 12 to the north-east and Village 15 to the north-west. Village 11 will adjoin the conservation area to the south of the precinct.

Existing features exhibit a woodland vegetative complex with drainage features present due to an undulating topography. Dominant trees species across several vegetation types include *Corymbia henryi*, *C. citriodora*, *Eucalyptus crebra*, *E. siderophloia*, *Lophostemon confertus* and *Angophora leiocarpa*.



Map 1: Project Location

(Image extracted from Springfield Rise at Spring Mountain Concept Masterplan, LandPartners 2016)

1.3 Current Permits and Authorities

All activities conducted during the site investigations were implemented under the provisions of a number of permits issued to Queensland Fauna Consultancy Pty Ltd by the Department of Environment and Heritage Protection (DEHP) formerly the Department of Environment and Resource Management and the Department of Employment, Economic Development and Innovation (DEEDI). These permits and additional authorities are listed in Table 1.

Table 1: Current Permits and authorities issued to QFC

Permit/Authorisation	Permit Number	Expiry Date
Damage Mitigation Permit	WIMP17840916	5 th December 2019
Rehabilitation Permit	WIRP15052614	10 th September 2017
Scientific Purposes Permit	WISP16935816	14 th February 2021
Scientific User Registration	Registration Number 589	27 th February 2019
Animal Ethics	CA 2016/01/939	27 th February 2019

These permits and approvals enable QFC to conduct the investigation, observation and relocation of protected animals exposed to disturbance due to infrastructure expansion resulting in the destruction of natural and artificial habitats.

2. Methodology

A site inspection was carried out on 27th June 2017 by Qld Fauna Consultancy. A standard set of observational techniques aimed at maximising the detection of fauna and the probable habitats they may occupy were employed to ascertain and identify the current fauna values throughout the project area. Where species of elevated conservation significance were foreseen as potentially present targeted searches were instigated to further evaluate individual species habitat.

Due to the habitat variability expressed across the development site the composition of investigations may include a range of features that entail specific components indicative of the presence of particular species or faunal groups. This may include where evident, observation of activity or signs of both historical and current use.

These may include but are not limited to the following:

- Identification of terrestrial microhabitats such as ground hollows, rock, burrows, leaf litter, stands of heavy vegetation, fallen branches and bark exfoliations;
- Identification of arboreal micro habitats including basal, trunk and limb hollows, tree fissures, bark exfoliates and arboreal termitaria;
- Identification of constructed arboreal micro habitats including bird nests and Ringtail Possum dreys;
- Artificial habitats including, but not limited to ornamental gardens, discarded rubbish, human dwellings and other infrastructure;
- Observation and investigation of aquatic habitats including dams, soaks, creeks, rivers and seasonally inundated vegetation communities. Artificial aquatic habitats may include constructed drains and culverts. Further components of interest include bank profiles and undercuts, submerged and/or exposed timber and rock, immediate aquatic and riparian vegetation, surfacing animals, nesting and/or feeding birds;
- Direct observation of active or exposed fauna within terrestrial, aquatic and arboreal habitats;
- Identification of scats, tracks and scratchings to determine fauna potentially present or to have historically utilised the site for either transient or longer term life history purposes.

2.1 Specific methodology for Koalas *Phascolarctos cinereus*

Due to specific requirements and the cryptic nature of the Koala the following techniques were employed to assist in ascertaining the current and historical presence/absence status of the species at the site:

- Use of binoculars to inspect the crown, forks and trunk of trees for individuals currently occupying the site;
- 'Drip zone' searches at the base of known food trees for the presence of scats to a radius equal to that of the crown of individual trees;
- Inspection of trunks for scratchings indicative of use by Koalas.

3. Findings

The findings endeavor to demarcate the existing habitat profiles and the features present into three distinct groups: terrestrial, arboreal and aquatic. All habitat features present onsite are noted, however it is probable additional features will be present with these being accounted for during the Fauna Spotter Catcher process to be applied to all vegetation clearing across the site.

3.1 Terrestrial Habitat Features

The terrestrial fauna values of the site consist of different components and microhabitat features. These features include low level understorey exhibiting sparse to moderate vegetative cover (Figure 1) interspersed with dense grass (Figure 2).

The site is exhibitive of a moderate amount of ground timber and woody debris (Figure 3) with scattered surface rocks also present in several localities (Figure 4) further adding to its potential habitat value for small reptiles and amphibian species.

Leaf litter (Figure 4) is also a feature on site, being present in abundance and at variable depths, providing refugial opportunities and microhabitat connectivity that can be exploited by many different native terrestrial vertebrate and invertebrate species.

Terrestrial termite mounds (Figure 5) also feature throughout the site, with excavations observed in mounds sighted during the survey indicating fauna foraging activity. Mammal assemblages may comprise both native and introduced species. Potential native mammals occurring on site include the Northern Brown Bandicoot *Isodon macrourus* which may be present in localities with significant vegetative ground cover. Small mammal burrows were also observed in several localities (Figure 6).

These features collectively contribute to the potential presence of a variety of native fauna species utilising the area for refugial, foraging and other resources.

GPS coordinates for all indicative terrestrial habitat features are shown in Table 2. Localities for identified terrestrial habitat features are presented in Map 2.

A comprehensive list of fauna species recorded in the region can be viewed in Appendix C.

Table 2: Localities for identified terrestrial habitat features

Number	Habitat Feature	GPS Coordinates	
		Easting	Northing
1	Burrow	0489295	6936895
2	Long grass/dense vegetation	0489204	6936907
3	Rock piles/boulders	0489255	6936873
4	Terrestrial termite mound	0489053	6936924
5	Terrestrial termite mound (with excavations)	0489246	6936871
6	Woody debris	0489146	6937076



Figure 1: Sparse to moderate understory



Figure 2: Dense grass



Figure 3: Woody debris



Figure 4: Boulders with woody debris and leaf litter



Figure 5: Excavated termite mound



Figure 6: Burrow

Map 2: Localities for identified terrestrial habitat features



Key for habitat feature type:

Terrestrial termitaria	Rock piles/boulders	Woody debris	Burrows	Long grass/dense vegetation
------------------------	---------------------	--------------	---------	-----------------------------

3.2 Arboreal Habitat Features

The majority of the clearance area consists predominately of Eucalypt woodland (Figure 7) consisting of trees of varying height, species and density suitable for feeding and nesting resources. The intermittent contiguous canopy structure within the vegetation represented may be facilitative of arboreal progression for species such as Brushtail Possum *Trichosurus vulpecula* and Common Ringtail Possum *Pseudocheirus peregrinus*. Fauna scratches indicating possum activity were found on numerous tree trunks (Figure 8).

Arboreal termite mounds are present across the site (Figure 9). Several suitable mounds were located with the potential for use by species such as the Lace Monitor *Varanus varius* which utilises arboreal termitaria for egg deposition and long-term incubation.

Hollow bearing trees and stags (Figure 10) feature throughout the site providing habitat opportunities for several arboreal mammal and reptile species. No avian nests or Possum dreys were found during the survey however further inspections are recommended immediately prior to clearing commencement. Numerous trees exhibited exfoliating bark (Figure 11), which may provide refugial opportunities for reptile species including skinks and geckos.

GPS coordinates for all indicative arboreal habitat features are shown in Table 3. Localities for identified arboreal habitat features are presented in Map 3.

Primary and secondary Koala food trees located in the clearance area, however no evidence was observed to indicate recent use of these trees by koalas. No koala scats were found during 'drip zone' searches and characteristic scratchings were not found during trunk investigations. A Koala habitat values map for the clearance area is presented in Appendix A.

Table 3: Localities for identified arboreal habitat features

Number	Habitat Feature	GPS Coordinates	
		Easting	Northing
1	Arboreal termitaria	0489395	6936920
2	Arboreal termitaria	0489393	6936959
3	Arboreal termitaria	0489373	6936946
4	Arboreal termitaria	0489263	6936942
5	Arboreal termitaria	0489304	6936972
6	Arboreal termitaria	0489314	6936963
7	Arboreal termitaria	0489109	6937016
8	Arboreal termitaria	0489393	6936959
9	Arboreal termitaria	0489159	6937056
10	Arboreal termitaria	0489148	6937048
11	Arboreal termitaria	0489171	6936964
12	Arboreal termitaria	0489168	6936987
13	Arboreal termitaria	0489160	6937017
14	Arboreal termitaria	0489212	6936994
15	Arboreal termitaria	0489223	6937002
16	Arboreal termitaria	0489297	6937038
17	Arboreal termitaria	0489339	6937024
18	Bark exfoliations	0489178	6936871
19	Bark exfoliations	0489184	6936874
20	Bark exfoliations	0489167	6937044
21	Hollow bearing tree	0489211	6936920
22	Hollow bearing tree	0489131	6936938
23	Hollow bearing tree	0489157	6937071
24	Hollow bearing tree	0489580	6936920
25	Hollow bearing tree	0489148	6937048

26	Hollow bearing tree	0489168	6936940
27	Hollow bearing tree	0489163	6936961
28	Hollow bearing tree	0489126	6936971
29	Hollow bearing tree	0489177	6937032



Figure 7: Site overview – Eucalypt woodland



Figure 8: Fauna scratches on tree trunk



Figure 9: Arboreal termitaria




Figure 10: Stag with hollows



Figure 11: Exfoliating bark

Map 3: Localities for identified arboreal habitat features



Key for habitat feature type:

Hollow bearing trees	Arboreal termitaria	Exfoliating bark
----------------------	---------------------	------------------

3.3 Aquatic Habitat Features

Several existing ephemeral drainage features are present within the clearance site and may provide breeding opportunities for frogs during significant rainfall events when intermittent ponded features are created. Several localities exhibited pooled water at the time of the survey (Figure 12). A number of native species may exploit the various microhabitats present by these environmental features, particularly during times of rainfall, including Graceful Treefrog *Litoria gracilentia*, Keelback Snake *Tropidonophis mairii* and various mammals and birds as a water resource.



Figure 12: Drainage feature with pooled water

3.4 Endangered, Vulnerable and Near Threatened (EVNT) Species

It is not envisaged that any EVNT fauna species will be detrimentally impacted by the proposed works. However, six species identified within the Online EPBC Protected Matters Report and the Queensland Government Wildlife Online Search Tool were considered likely or possible to occur within the site and will require further mitigation during clearing activities.

Although no evidence was found during the site inspection of recent Koala use the species has previously been recorded in the area. Some areas within the site are identified as High Value Bushland under Koala Habitat in South East Queensland mapping sourced from the DEHP online search tool (see Appendix A). It is advised that dedicated methodologies be employed by a qualified Fauna Spotter specific to the detection of these species prior to vegetation clearing activities.

Table 4: Significant species deemed likely or possible to occur within the clearance survey area

Common Name Scientific Name	Species Information	Likelihood of Occurrence within the Clearance Survey area
Mammals		
Koala <i>Phascolarctos cinereus</i> EPBC: Vulnerable NCA: Vulnerable	Inhabits a range of open forest and woodland communities which may include any of the following noted food trees: <i>Eucalyptus</i> , <i>Corymbia</i> , <i>Melaleuca</i> , <i>Angophora</i> and <i>Lophostemon</i> .	Likely Known food trees for the transient Koala (<i>Phascolarctos cinereus</i>) occur on the clearance site and the species is well documented within the area.
Grey-headed Flying-fox <i>Pteropus poliocephalus</i> EPBC: Vulnerable NCA: Least Concern	The Grey-headed Flying-Fox roosts in aggregations of various sizes on exposed branches, commonly of emergent trees. Roost sites are typically located near water, such as lakes, rivers or the coast. Habitat includes open forests, woodlands, urban parks and gardens.	Possible Suitable vegetation communities containing both feeding and roosting resources occur on and adjacent to the clearance site.
Spotted-tail Quoll (SE Mainland Population) <i>Dasyurus maculatus maculatus</i> EPBC: Endangered NCA: Vulnerable	Currently known from the Granit Belt and Border Ranges though small numbers may occur from Gympie to the QLD border (Curtis <i>et al.</i> 2012). Inhabits vine-forest, wet and dry sclerophyll forests and woodlands containing boulder piles, fallen logs and hollow trees utilised as shelter sites (Curtis <i>et al.</i> 2012).	Possible Preferred habitat type and habitat features present and the species is documented within the area.
Greater Glider <i>Petauroides volans</i> EPBC: Vulnerable NCA: Least Concern	Largest of the gliders, the Great Glider is found along eastern Australia within a variety of eucalypt dominated forests and tall open woodlands (Lindenmayer 2002)	Possible Preferred habitat type and habitat features present and the species is documented within the area.

Birds		
<p>Powerful Owl <i>Ninox strenua</i></p> <p>EPBC: Not Listed NCA: Vulnerable</p>	<p>Inhabits forests and woodlands of eastern and south-eastern Australia (Beruldsen 2003). Breeds once per year in May to July or August. Nests in hollow trunks or limbs of large trees, usually at considerable height (Beruldsen 2003).</p>	<p>Possible Preferred habitat types present and the species is documented within the area.</p>
Reptiles		
<p>Collared Delma <i>Delma torquata</i></p> <p>EPBC: Vulnerable NCA: Vulnerable</p>	<p>Weathered loose rocks, flattish bedrock outcroppings, logs or mats of leaf litter, or in cracks and crevices among tussock grasses. Lays two eggs around December with hatching in February or March (Curtis <i>et al.</i> 2012)</p>	<p>Possible Preferred habitat type and habitat features present.</p>

4. Fauna Impacts

It is important to consider the existing and future residential developmental areas when investigation potential fauna impacts.

Impacts to fauna, as a result of vegetation clearance, will include the following:

- Loss of trees for foraging, roosting and nesting;
- Loss of hollow-bearing trees for nesting and refuge;
- Loss of habitat and foraging areas for terrestrial species;
- Loss of overall habitat;
- Potential loss of abundance of some local species.

Other impacts may include:

- Injury or death during felling of trees;
- Injury or death from machinery;
- Alteration of nesting, foraging and general activities due to disturbance.

5. Assessment and Conclusion

Overall the site contains medium value refugial opportunities for arboreal and terrestrial fauna species (see Section 3.1 and 3.2). The species expected within the site are likely to primarily reflect common fauna assemblages for the region; however, provisions will be proposed directly for common fauna and species of conservation significance.

The connectivity to adjacent conservation land in the south, in conjunction with sequential clearing methodologies, will aid in the movement of medium to large size fauna such as Koala and Kangaroos. Specific methodologies for these species will be detailed within the Wildlife and Habitat Impact Mitigation Plan (WHIMP).

A number of conclusions and recommendations will be presented in the WHIMP, with the specific intention of providing a comprehensive management structure to facilitate minimal impact to fauna during the clearing of vegetation and subsequent disturbance of habitats.

It is advised that all identified fauna habitats onsite be inspected by a DEHP approved Fauna Spotter prior to vegetation clearing and all vegetation removal activities be supervised during the clearing process. Terrestrial load reduction activities will be conducted ahead of the clearing front where possible. Fauna captured will be relocated to adjacent habitat consistent with the life history requirements of the species requiring translocation. The directives given by Fauna Spotter Catchers should embrace a “best practice” approach which includes implementation of proven specific management techniques for identified habitat types and compliance with legislation relevant to the activity.

It is recommended that in the event any nests which contain chicks are identified during clearing be left until fledged, and those that are in a construction phase should be dismantled to prevent further nesting activity. Any fertile eggs recovered will require incubation and subsequent rearing for latter release.

6. References

Anstis, M (2013) *Tadpoles and Frogs of Australia*, New Holland Publishers, Sydney.

Beruldsen, G. (2003) *Australian Birds their Nests and Eggs*, Kenmore Hills: Self Published.

Curtis, LK, Dennis, AJ, McDonald, KR, Kyne, PM & Debus, SJS (2012), Queensland's Threatened Animals, CSIRO Publishing, Victoria.

Department of Sustainability, Environment, Water, Population and Communities (2012) EPBC Act Protected Matters Report. Date created 4th July 2017.

Department of Environment and Heritage Protection (2012) Koala habitat map request form, <http://www.ehp.qld.gov.au/wildlife/koalas/mapping/maprequestform.php>
Date accessed 14th March 2017.

Department of Environment and Heritage Protection (2012) Wildlife and Ecosystems- Threatened Species, <http://www.ehp.qld.gov.au/wildlife/threatened-species/index.html>
Date created 4th July 2017.

LandPartners (2016). *Springfield Rise at Spring Mountain Concept Master Plan*. Plan prepared for Lend Lease Communities Pty Ltd, January 2016.

Lindenmayer, D. (2002) *Gliders of Australia – A Natural History*, UNSW Press Ltd, Sydney

Queensland Environmental Protection Agency and Queensland Parks and Wildlife Service (2006). *Nature Conservation (Koala) Conservation Plan 2006 and Management Plan 2006 – 2016*. Queensland Government – Environmental Protection Agency.

Van Dyck, S. & Strahan, R (2008). *The Mammals of Australia*, 3rd edn, Reed New Holland, Sydney.

References for nomenclature

Brooker, M.I.H. and Kleinig, D.A. (2004) *Field Guide to Eucalypts: Volume 3 Northern Australia*, 2nd edn, Melbourne: Blooming Books.

Churchill, S. (2008) *Australian Bats*, 2nd edition, Sydney: Allen & Unwin.

Cogger, H. (2000) *Reptiles & Amphibians of Australia*. 6th edition, Sydney: Reed New Holland.

Leiper, G., Glazebrook, J., Cox, D. and Rathie, K. (2008) *Mangroves to Mountains: a Field Guide to the Native Plants of South-east Queensland*, Browns Plains: Logan River Branch Society for Growing Australian Plants.

Menkhorst, K. & Knight, F. (2011) *A Field Guide to the Mammals of Australia*, 3rd edition, South Melbourne: Oxford University Press.


Strahan, R. (Ed) (1995) *The Mammals of Australia*. Sydney: New Holland Publishers.

Wilson, S. (2005) *A Field Guide to Reptiles of Queensland*. Sydney: New Holland Publishers.

Queensland Museum (2007) *Wildlife of Greater Brisbane*, 2nd edition, Brisbane: Queensland Museum Publishers.

Vanderduys, E. (2012) *Field Guide to the Frogs of Queensland*. Collingwood: CSIRO Publishing.

7. Appendix A: Koala Habitat Values



8. Appendix B: EPBC Act Protected Matters Report



Australian Government
Department of the Environment and Energy

EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 04/07/17 11:32:41

[Summary](#)

[Details](#)


[Matters of NES](#)

[Other Matters Protected by the EPBC Act](#)

[Extra Information](#)

[Caveat](#)

[Acknowledgements](#)



This map may contain data which are
©Commonwealth of Australia
(Geoscience Australia), ©PSMA 2010

[Coordinates](#)

[Buffer: 5.0Km](#)



Name	Status	Type of Presence
Poeophila cincta_cincta Southern Black-throated Finch [64447]	Endangered	Species or species habitat may occur within area
Rostratula australis Australian Painted Snipe [77037]	Endangered	Species or species habitat likely to occur within area
Turnix melanogaster Black-breasted Button-quail [923]	Vulnerable	Species or species habitat likely to occur within area
Insects		
Phylodes imperialis_smithersi Pink Underwing Moth [86084]	Endangered	Species or species habitat may occur within area
Mammals		
Chalinolobus dwyeri Large-eared Pled Bat, Large Pled Bat [183]	Vulnerable	Species or species habitat likely to occur within area
Dasyurus hallucatus Northern Quoll, Digul [331]	Endangered	Species or species habitat may occur within area
Dasyurus maculatus_maculatus (SE mainland population) Spot-tailed Quoll, Spotted-tail Quoll, Tiger Quoll (southeastern mainland population) [75184]	Endangered	Species or species habitat may occur within area
Petauroides volans Greater Glider [254]	Vulnerable	Species or species habitat known to occur within area
Petrogale penicillata Brush-tailed Rock-wallaby [225]	Vulnerable	Species or species habitat known to occur within area
Phascolarctos cinereus (combined populations of Qld, NSW and the ACT) Koala (combined populations of Queensland, New South Wales and the Australian Capital Territory) [85104]	Vulnerable	Species or species habitat known to occur within area
Pteropus poliocephalus Grey-headed Flying-fox [186]	Vulnerable	Foraging, feeding or related behaviour known to occur within area
Other		
Cycas ophiolitica [55797]	Endangered	Species or species habitat likely to occur within area
Plants		
Bosistoa transversa Three-leaved Bosistoa, Yellow Satinheart [16091]	Vulnerable	Species or species habitat likely to occur within area
Dichanthium setosum bluegrass [14159]	Vulnerable	Species or species habitat likely to occur within area
Macadamia integrifolia Macadamia Nut, Queensland Nut Tree, Smooth-shelled Macadamia, Bush Nut, Nut Oak [7326]	Vulnerable	Species or species habitat likely to occur within area
Macadamia tetraphylla Rough-shelled Bush Nut, Macadamia Nut, Rough-shelled Macadamia, Rough-leaved Queensland Nut [6581]	Vulnerable	Species or species habitat may occur within area
Notelaea lpsyclensis Cooneana Olive [81858]	Critically Endangered	Species or species habitat may occur within area

Extra Information

State and Territory Reserves [\[Resource Information \]](#)

Name	State
Stewartdale	QLD
White Rock	QLD

Invasive Species [\[Resource Information \]](#)

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resources Audit, 2001.

Name	Status	Type of Presence
------	--------	------------------

Birds

Acridotheres tristis Common Myna, Indian Myna [387]		Species or species habitat likely to occur within area
Anas platyrhynchos Mallard [974]		Species or species habitat likely to occur within area
Carduelis carduelis European Goldfinch [403]		Species or species habitat likely to occur within area
Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Lonchura punctulata Nutmeg Mannikin [399]		Species or species habitat likely to occur within area
Passer domesticus House Sparrow [405]		Species or species habitat likely to occur within area
Streptopelia chinensis Spotted Turtle-Dove [780]		Species or species habitat likely to occur within area
Sturnus vulgaris Common Starling [389]		Species or species habitat likely to occur within area

Frogs

Rhinella marina Cane Toad [83218]		Species or species habitat likely to occur within area
--------------------------------------	--	--

Mammals

Name	Status	Type of Presence
<i>Bos taurus</i> Domestic Cattle [16]		Species or species habitat likely to occur within area
<i>Canis lupus familiaris</i> Domestic Dog [82654]		Species or species habitat likely to occur within area
<i>Equus caballus</i> Horse [5]		Species or species habitat likely to occur within area
<i>Felis catus</i> Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
<i>Lepus capensis</i> Brown Hare [127]		Species or species habitat likely to occur within area
<i>Mus musculus</i> House Mouse [120]		Species or species habitat likely to occur within area
<i>Oryctolagus cuniculus</i> Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
<i>Rattus norvegicus</i> Brown Rat, Norway Rat [83]		Species or species habitat likely to occur within area
<i>Rattus rattus</i> Black Rat, Ship Rat [84]		Species or species habitat likely to occur within area
<i>Sus scrofa</i> Pig [6]		Species or species habitat likely to occur within area
<i>Vulpes vulpes</i> Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
<i>Cabomba caroliniana</i> Cabomba, Fanwort, Carolina Watershield, Fish Grass, Washington Grass, Watershield, Carolina Fanwort, Common Cabomba [5171]		Species or species habitat likely to occur within area
<i>Chrysanthemoides monilifera</i> Bitou Bush, Boneseed [18983]		Species or species habitat may occur within area
<i>Eichhornia crassipes</i> Water Hyacinth, Water Orchid, Nile Lily [13466]		Species or species habitat likely to occur within area
<i>Genista monspessulana</i> Montpellier Broom, Cape Broom, Canary Broom, Common Broom, French Broom, Soft Broom [20126]		Species or species habitat likely to occur within area
<i>Lantana camara</i> Lantana, Common Lantana, Kamara Lantana, Large-leaf Lantana, Pink Flowered Lantana, Red Flowered Lantana, Red-Flowered Sage, White Sage, Wild Sage [10892]		Species or species habitat likely to occur within area
<i>Parkinsonia aculeata</i> Parkinsonia, Jerusalem Thorn, Jelly Bean Tree, Horse Bean [12301]		Species or species habitat likely to occur within area
<i>Parthenium hysterophorus</i> Parthenium Weed, Bitter Weed, Carrot Grass, False Ragweed [19566]		Species or species habitat likely to occur

9. Appendix C: Wildlife Online Extract



Wildlife Online Extract

Search Criteria: Species List for a Specified Point
Species: Animals
Type: Native
Status: All
Records: All
Date: Since 1980
Latitude: -27.6915
Longitude: 152.8915
Distance: 5
Email: ramona@qfc.com.au
Date submitted: Tuesday 04 Jul 2017 11:32:57
Date extracted: Tuesday 04 Jul 2017 11:40:16

The number of records retrieved = 298

Disclaimer

As the DSITIA is still in a process of collating and vetting data, it is possible the information given is not complete. The information provided should only be used for the project for which it was requested and it should be appropriately acknowledged as being derived from Wildlife Online when it is used.

The State of Queensland does not invite reliance upon, nor accept responsibility for this information. Persons should satisfy themselves through independent means as to the accuracy and completeness of this information.

No statements, representations or warranties are made about the accuracy or completeness of this information. The State of Queensland disclaims all responsibility for this information and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you may incur as a result of the information being inaccurate or incomplete in any way for any reason.

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	amphibians	Hylidae	<i>Litoria fallax</i>	eastern sedgefrog		C		14
animals	amphibians	Hylidae	<i>Litoria nasuta</i>	striped rocketfrog		C		5
animals	amphibians	Hylidae	<i>Litoria dentata</i>	bleating treefrog		C		1
animals	amphibians	Hylidae	<i>Litoria rubella</i>	ruddy treefrog		C		8
animals	amphibians	Hylidae	<i>Cyclorana alboguttata</i>	greenstripe frog		C		1
animals	amphibians	Hylidae	<i>Litoria wilcoxii</i>	eastern stony creek frog		C		5
animals	amphibians	Hylidae	<i>Litoria gracilenta</i>	graceful treefrog		C		12
animals	amphibians	Hylidae	<i>Litoria latopalmata</i>	broad palmed rocketfrog		C		6
animals	amphibians	Hylidae	<i>Litoria brevipalmata</i>	green thighed frog		C		1
animals	amphibians	Hylidae	<i>Litoria caerulea</i>	common green treefrog		C		5
animals	amphibians	Limnodynastidae	<i>Limnodynastes tasmaniensis</i>	spotted grassfrog		C		2
animals	amphibians	Limnodynastidae	<i>Limnodynastes terraereginae</i>	scarlet sided pobblebonk		C		9
animals	amphibians	Limnodynastidae	<i>Platyplectrum ornatum</i>	ornate burrowing frog		C		22
animals	amphibians	Limnodynastidae	<i>Limnodynastes peronii</i>	striped marshfrog		C		9
animals	amphibians	Limnodynastidae	<i>Adelotus brevis</i>	tusked frog		V		1
animals	amphibians	Myobatrachidae	<i>Pseudophryne coriacea</i>	red backed broodfrog		C		1
animals	amphibians	Myobatrachidae	<i>Mixophyes fasciolatus</i>	great barred frog		C		8
animals	amphibians	Myobatrachidae	<i>Crinia parinsignifera</i>	beeping froglet		C		4
animals	amphibians	Myobatrachidae	<i>Pseudophryne raveni</i>	copper backed broodfrog		C		10
animals	amphibians	Myobatrachidae	<i>Pseudophryne major</i>	great brown broodfrog		C		1
animals	amphibians	Myobatrachidae	<i>Uperoleia rugosa</i>	chubby gungan		C		2
animals	birds	Acanthizidae	<i>Smicrorhnis brevirostris</i>	weebill		C		43
animals	birds	Acanthizidae	<i>Chthonicola sagittata</i>	speckled warbler		C		17
animals	birds	Acanthizidae	<i>Acanthiza nana</i>	yellow thornbill		C		8
animals	birds	Acanthizidae	<i>Gerygone mouki</i>	brown gerygone		C		1
animals	birds	Acanthizidae	<i>Acanthiza chrysorrhoa</i>	yellow-rumped thornbill		C		2
animals	birds	Acanthizidae	<i>Acanthiza pusilla</i>	brown thornbill		C		19
animals	birds	Acanthizidae	<i>Gerygone olivacea</i>	white-throated gerygone		C		47
animals	birds	Acanthizidae	<i>Acanthiza reguloides</i>	buff-rumped thornbill		C		24
animals	birds	Acanthizidae	<i>Sericornis frontalis</i>	white-browed scrubwren		C		34
animals	birds	Acanthizidae	<i>Acanthiza lineata</i>	striated thornbill		C		9
animals	birds	Accipitridae	<i>Accipiter cirrocephalus</i>	collared sparrowhawk		C		2
animals	birds	Accipitridae	<i>Accipiter novaehollandiae</i>	grey goshawk		C		2
animals	birds	Accipitridae	<i>Aquila audax</i>	wedge-tailed eagle		C		29
animals	birds	Accipitridae	<i>Elanus axillaris</i>	black-shouldered kite		C		9
animals	birds	Accipitridae	<i>Accipiter fasciatus</i>	brown goshawk		C		14
animals	birds	Accipitridae	<i>Accipiter subcristata</i>	Pacific baza		C		8
animals	birds	Accipitridae	<i>Haliaeetus leucogaster</i>	white-bellied sea-eagle		C		4
animals	birds	Accipitridae	<i>Hieraaetus morphnoides</i>	little eagle		C		2
animals	birds	Acrocephalidae	<i>Acrocephalus australis</i>	Australian reed-warbler		C		2
animals	birds	Aegothelidae	<i>Aegotheles cristatus</i>	Australian owllet-nightjar		C		13
animals	birds	Alcedinidae	<i>Ceyx azureus</i>	azure kingfisher		C		3
animals	birds	Alcedinidae	<i>Ceyx pusillus</i>	little kingfisher		C		1
animals	birds	Anatidae	<i>Cygnus atratus</i>	black swan		C		4
animals	birds	Anatidae	<i>Aythya australis</i>	hardhead		C		5
animals	birds	Anatidae	<i>Anas gracilis</i>	grey teal		C		4

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	birds	Anatidae	<i>Dendrocygna arcuata</i>	wandering whistling-duck		C		1
animals	birds	Anatidae	<i>Chenonetta jubata</i>	Australian wood duck		C		25
animals	birds	Anatidae	<i>Anas superciliosa</i>	Pacific black duck		C		22
animals	birds	Anhingidae	<i>Anhinga novaehollandiae</i>	Australasian darter		C		6
animals	birds	Anseranatidae	<i>Anseranas semipalmata</i>	magpie goose		C		1
animals	birds	Apodidae	<i>Hirundapus caudacutus</i>	white-throated needletail		SL		8
animals	birds	Ardeidae	<i>Ardea intermedia</i>	intermediate egret		C		4
animals	birds	Ardeidae	<i>Bubulcus ibis</i>	cattle egret		C		16
animals	birds	Ardeidae	<i>Ardea pacifica</i>	white-necked heron		C		3
animals	birds	Ardeidae	<i>Nycticorax caledonicus</i>	nankeen night-heron		C		1
animals	birds	Ardeidae	<i>Egretta novaehollandiae</i>	white-faced heron		C		19
animals	birds	Ardeidae	<i>Ardea alba modesta</i>	eastern great egret		C		2
animals	birds	Artamidae	<i>Artamus superciliosus</i>	white-browed woodswallow		C		1
animals	birds	Artamidae	<i>Cracticus nigrogularis</i>	piebald butcherbird		C		73
animals	birds	Artamidae	<i>Artamus leucorhynchus</i>	white-breasted woodswallow		C		4
animals	birds	Artamidae	<i>Cracticus torquatus</i>	grey butcherbird		C		54
animals	birds	Artamidae	<i>Artamus cyanopterus</i>	dusky woodswallow		C		9
animals	birds	Artamidae	<i>Cracticus tibicen</i>	Australian magpie		C		72
animals	birds	Artamidae	<i>Strepera graculina</i>	piebald currawong		C		63
animals	birds	Burhinidae	<i>Burhinus grallarius</i>	bush stone-curlew		C		1
animals	birds	Cacatuidae	<i>Calyptorhynchus lathami lathami</i>	glossy black-cockatoo (eastern)		V		2
animals	birds	Cacatuidae	<i>Calyptorhynchus banksii</i>	red-tailed black-cockatoo		C		5
animals	birds	Cacatuidae	<i>Eolophus roseicapilla</i>	galah		C		32
animals	birds	Cacatuidae	<i>Cacatua sanguinea</i>	little corella		C		2
animals	birds	Cacatuidae	<i>Cacatua galerita</i>	sulphur-crested cockatoo		C		41
animals	birds	Campephagidae	<i>Coracina novaehollandiae</i>	black-faced cuckoo-shrike		C		69
animals	birds	Campephagidae	<i>Coracina tenuirostris</i>	black-fronted dotterel		C		30
animals	birds	Campephagidae	<i>Coracina papuensis</i>	white-bellied cuckoo-shrike		C		9
animals	birds	Campephagidae	<i>Lalage tricolor</i>	white-winged triller		C		1
animals	birds	Campephagidae	<i>Lalage leucomela</i>	varied triller		C		12
animals	birds	Charadriidae	<i>Vanellus miles novaehollandiae</i>	masked lapwing (southern subspecies)		C		18
animals	birds	Charadriidae	<i>Vanellus miles</i>	masked lapwing		C		9
animals	birds	Charadriidae	<i>Elseyornis melanops</i>	black-fronted dotterel		C		2
animals	birds	Ciconiidae	<i>Ephippiorhynchus asiaticus</i>	black-necked stork		C		2
animals	birds	Cisticolidae	<i>Cisticola exilis</i>	golden-headed cisticola		C		21
animals	birds	Climacteridae	<i>Cormobates leucophaea metastasis</i>	white-throated treecreeper (southern)		C		47
animals	birds	Climacteridae	<i>Cormobates leucophaea</i>	white-throated treecreeper		C		8
animals	birds	Climacteridae	<i>Climacteris affinis</i>	white-browed treecreeper		C		1
animals	birds	Columbidae	<i>Lopholaimus antarcticus</i>	topknot pigeon		C		7
animals	birds	Columbidae	<i>Leucosarcia melanoleuca</i>	wonga pigeon		C		1
animals	birds	Columbidae	<i>Macropygia amboinensis</i>	brown cuckoo-dove		C		19
animals	birds	Columbidae	<i>Geopelia humeralis</i>	bar-shouldered dove		C		39
animals	birds	Columbidae	<i>Chalcophaps indica</i>	emerald dove		C		6
animals	birds	Columbidae	<i>Phaps chalcoptera</i>	common bronzewing		C		21
animals	birds	Columbidae	<i>Ocyphaps lophotes</i>	crested pigeon		C		32
animals	birds	Columbidae	<i>Geopelia striata</i>	peaceful dove		C		39

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	birds	Coraciidae	<i>Eurystomus orientalis</i>	dollarbird		C		35
animals	birds	Corvidae	<i>Corvus coronoides</i>	Australian raven		C		1
animals	birds	Corvidae	<i>Corvus orru</i>	Torresian crow		C		127
animals	birds	Cuculidae	<i>Eudynamis orientalis</i>	eastern koel		C		22
animals	birds	Cuculidae	<i>Chalcites minutillus barnardi</i>	little bronze-cuckoo		C		1
animals	birds	Cuculidae	<i>Scythrops novaehollandiae</i>	channel-billed cuckoo		C		22
animals	birds	Cuculidae	<i>Cacomantis flabelliformis</i>	fan-tailed cuckoo		C		30
animals	birds	Cuculidae	<i>Centropus phasianinus</i>	pheasant coucal		C		21
animals	birds	Cuculidae	<i>Cacomantis variolosus</i>	brush cuckoo		C		15
animals	birds	Cuculidae	<i>Chalcites basalus</i>	Horsfield's bronze-cuckoo		C		9
animals	birds	Cuculidae	<i>Cuculus optatus</i>	oriental cuckoo		SL		5
animals	birds	Cuculidae	<i>Chalcites lucidus</i>	shining bronze-cuckoo		C		11
animals	birds	Dicruridae	<i>Dicrurus bracteatus bracteatus</i>	spangled drongo (eastern Australia)		C		1
animals	birds	Dicruridae	<i>Dicrurus bracteatus</i>	spangled drongo		C		40
animals	birds	Estrildidae	<i>Neochmia temporalis</i>	red-browed finch		C		53
animals	birds	Estrildidae	<i>Lonchura castaneothorax</i>	chestnut-breasted mannikin		C		8
animals	birds	Estrildidae	<i>Taeniopygia bichenovii</i>	double-barred finch		C		26
animals	birds	Eurostopodidae	<i>Eurostopodus mystacalis</i>	white-throated nightjar		C		14
animals	birds	Falconidae	<i>Falco hypoleucos</i>	grey falcon		V		1
animals	birds	Falconidae	<i>Falco longipennis</i>	Australian hobby		C		3
animals	birds	Falconidae	<i>Falco cenchroides</i>	nankeen kestrel		C		13
animals	birds	Falconidae	<i>Falco peregrinus</i>	peregrine falcon		C		13
animals	birds	Halcyonidae	<i>Dacelo novaeguineae</i>	laughing kookaburra		C		88
animals	birds	Halcyonidae	<i>Todiramphus macleayii</i>	forest kingfisher		C		15
animals	birds	Halcyonidae	<i>Todiramphus sanctus</i>	sacred kingfisher		C		28
animals	birds	Hirundinidae	<i>Hirundo neoxena</i>	welcome swallow		C		28
animals	birds	Hirundinidae	<i>Petrochelidon ariel</i>	fairy martin		C		8
animals	birds	Hirundinidae	<i>Cheramoeca leucosterna</i>	white-backed swallow		C		8
animals	birds	Hirundinidae	<i>Petrochelidon nigricans</i>	tree martin		C		14
animals	birds	Jacaniidae	<i>Irediparra gallinacea</i>	comb-crested jacana		C		6
animals	birds	Maluridae	<i>Malurus lamberti</i>	variegated fairy-wren		C		55
animals	birds	Maluridae	<i>Malurus cyaneus</i>	superb fairy-wren		C		32
animals	birds	Maluridae	<i>Malurus melanocephalus</i>	red-backed fairy-wren		C		69
animals	birds	Megaluridae	<i>Megalurus timoriensis</i>	tawny grassbird		C		8
animals	birds	Megapodiidae	<i>Alectura lathamii</i>	Australian brush-turkey		C		9
animals	birds	Meliphagidae	<i>Acanthorhynchus tenuirostris</i>	eastern spinebill		C		22
animals	birds	Meliphagidae	<i>Plectorhyncha lanceolata</i>	striped honeyeater		C		13
animals	birds	Meliphagidae	<i>Melithreptus albogularis</i>	white-throated honeyeater		C		73
animals	birds	Meliphagidae	<i>Philemon citreogularis</i>	little friarbird		C		21
animals	birds	Meliphagidae	<i>Anthochaera chrysoptera</i>	little wattlebird		C		8
animals	birds	Meliphagidae	<i>Ptilotula fusca</i>	fuscous honeyeater		C		13
animals	birds	Meliphagidae	<i>Meliphaga lewinii</i>	Lewin's honeyeater		C		52
animals	birds	Meliphagidae	<i>Caligavis chrysops</i>	yellow-faced honeyeater		C		89
animals	birds	Meliphagidae	<i>Entomyzon cyanotis</i>	blue-faced honeyeater		C		24
animals	birds	Meliphagidae	<i>Lichmera indistincta</i>	brown honeyeater		C		54
animals	birds	Meliphagidae	<i>Melithreptus gularis</i>	black-chinned honeyeater		C		6

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	birds	Meliphagidae	<i>Meliphaga lunata</i>	white-naped honeyeater		C		5
animals	birds	Meliphagidae	<i>Philemon corniculatus</i>	noisy friarbird		C		90
animals	birds	Meliphagidae	<i>Lichenostomus melanops</i>	yellow-tufted honeyeater		C		11
animals	birds	Meliphagidae	<i>Myzomela sanguinolenta</i>	scarlet honeyeater		C		85
animals	birds	Meliphagidae	<i>Manorina melanocephala</i>	noisy miner		C		74
animals	birds	Meropidae	<i>Merops ornatus</i>	rainbow bee-eater		C		60
animals	birds	Monarchidae	<i>Grallina cyanoleuca</i>	maggpie-lark		C		44
animals	birds	Monarchidae	<i>Myiagra cyanoleuca</i>	satin flycatcher		SL		1
animals	birds	Monarchidae	<i>Myiagra inquieta</i>	restless flycatcher		C		6
animals	birds	Monarchidae	<i>Symphoricarpha trivirgatus</i>	spectacled monarch		SL		8
animals	birds	Monarchidae	<i>Monarcha melanopsis</i>	black-faced monarch		SL		15
animals	birds	Monarchidae	<i>Myiagra rubecula</i>	leaden flycatcher		C		38
animals	birds	Motacillidae	<i>Anthus novaeseelandiae</i>	Australasian pipit		C		3
animals	birds	Nectariniidae	<i>Dicaeum hirundinaceum</i>	mistletoebird		C		42
animals	birds	Neosittidae	<i>Daphoenositta chrysoptera</i>	varied sittella		C		35
animals	birds	Oriolidae	<i>Sphecotheres vieilloti</i>	Australasian figbird		C		18
animals	birds	Oriolidae	<i>Oriolus sagittatus</i>	olive-backed oriole		C		34
animals	birds	Pachycephalidae	<i>Falcunculus frontatus</i>	crested shrike-tit		C		1
animals	birds	Pachycephalidae	<i>Pachycephala rufiventris</i>	rufous whistler		C		68
animals	birds	Pachycephalidae	<i>Colluricincla megarhyncha</i>	little shrike-thrush		C		12
animals	birds	Pachycephalidae	<i>Colluricincla harmonica</i>	grey shrike-thrush		C		64
animals	birds	Pachycephalidae	<i>Pachycephala pectoralis</i>	golden whistler		C		42
animals	birds	Pardalotidae	<i>Pardalotus striatus</i>	striated pardalote		C		101
animals	birds	Pardalotidae	<i>Pardalotus punctatus</i>	spotted pardalote		C		40
animals	birds	Pelecanidae	<i>Pelecanus conspicillatus</i>	Australian pelican		C		1
animals	birds	Petroicidae	<i>Microeca fascians</i>	jacky winter		C		22
animals	birds	Petroicidae	<i>Eopsaltria australis</i>	eastern yellow robin		C		59
animals	birds	Petroicidae	<i>Petroica rosea</i>	rose robin		C		27
animals	birds	Phalacrocoracidae	<i>Microcarbo melanoleucos</i>	little pied cormorant		C		9
animals	birds	Phalacrocoracidae	<i>Phalacrocorax sulcirostris</i>	little black cormorant		C		4
animals	birds	Phasianidae	<i>Coturnix ypsilophora</i>	brown quail		C		18
animals	birds	Podargidae	<i>Podargus strigoides</i>	tawny frogmouth		C		22
animals	birds	Podicipedidae	<i>Tachybaptus novaehollandiae</i>	Australasian grebe		C		9
animals	birds	Pomatostomidae	<i>Pomatostomus temporalis</i>	grey-crowned babbler		C		12
animals	birds	Psittacidae	<i>Alisterus scapularis</i>	Australian king-parrot		C		17
animals	birds	Psittacidae	<i>Parvipsitta pusilla</i>	little lorikeet		C		45
animals	birds	Psittacidae	<i>Trichoglossus haematodus moluccanus</i>	rainbow lorikeet		C		74
animals	birds	Psittacidae	<i>Platycercus adscitus palliceps</i>	pale-headed rosella (southern form)		C		2
animals	birds	Psittacidae	<i>Platycercus eximius</i>	eastern rosella		C		13
animals	birds	Psittacidae	<i>Platycercus adscitus</i>	pale-headed rosella		C		46
animals	birds	Psittacidae	<i>Trichoglossus chlorolepidotus</i>	scaly-breasted lorikeet		C		62
animals	birds	Psophodidae	<i>Psophodes olivaceus</i>	eastern whipbird		C		52
animals	birds	Psophodidae	<i>Cinclosoma punctatum</i>	spotted quail-thrush		C		11
animals	birds	Ptilonorhynchidae	<i>Ptilonorhynchus maculatus</i>	spotted bowerbird		C		1
animals	birds	Ptilonorhynchidae	<i>Sericulus chrysocephalus</i>	regent bowerbird		C		1
animals	birds	Rallidae	<i>Fulica atra</i>	Eurasian coot		C		8

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	birds	Rallidae	<i>Gallinula tenebrosa</i>	dusky moorhen		C		15
animals	birds	Rallidae	<i>Porphyrio melanotus</i>	purple swamphen		C		7
animals	birds	Recurvirostridae	<i>Himantopus himantopus</i>	black-winged stilt		C		1
animals	birds	Rhipiduridae	<i>Rhipidura albiscapa</i>	grey fantail		C		74
animals	birds	Rhipiduridae	<i>Rhipidura leucophrys leucophrys</i>	willie wagtail (southern)		C		1
animals	birds	Rhipiduridae	<i>Rhipidura leucophrys</i>	willie wagtail		C		50
animals	birds	Rhipiduridae	<i>Rhipidura rufifrons</i>	rufous fantail		SL		25
animals	birds	Strigidae	<i>Ninox boobook</i>	southern boobook		C		28
animals	birds	Strigidae	<i>Ninox strenua</i>	powerful owl		V		6
animals	birds	Threskiomithidae	<i>Threskiornis molucca</i>	Australian white ibis		C		8
animals	birds	Threskiomithidae	<i>Threskiornis spinicollis</i>	straw-necked ibis		C		10
animals	birds	Threskiomithidae	<i>Platalea regia</i>	royal spoonbill		C		2
animals	birds	Timaliidae	<i>Zosterops lateralis</i>	silveryeye		C		69
animals	birds	Timaliidae	<i>Zosterops lateralis cornwalli</i>	silveryeye (eastern)		C		1
animals	birds	Turnicidae	<i>Turnix varius</i>	painted button-quail		C		14
animals	birds	Tytonidae	<i>Tyto novaehollandiae novaehollandiae</i>	masked owl (southern subspecies)		C		1
animals	insects	Hesperiidae	<i>Neohesperilla xanthomera</i>	yellow grass-skipper				1
animals	insects	Lycaenidae	<i>Candalides cyprotus pallescens</i>	copper pencilled-blue				1
animals	insects	Lycaenidae	<i>Acrodipsas brisbanensis</i>	bronze ant-blue				2
animals	insects	Lycaenidae	<i>Ogyris oroetes oroetes</i>	silky azure				1
animals	insects	Lycaenidae	<i>Ogyris zosine zosine</i>	northern purple azure (southern subspecies)				1
animals	insects	Nymphalidae	<i>Charaxes sempronius sempronius</i>	tailed emperor				1
animals	insects	Nymphalidae	<i>Acraea andromacha andromacha</i>	glasswing				7
animals	insects	Nymphalidae	<i>Tirumala hamata hamata</i>	blue tiger				1
animals	insects	Nymphalidae	<i>Junonia villida calybe</i>	meadow argus				1
animals	insects	Nymphalidae	<i>Melanitis leda bankia</i>	common evening-brown				3
animals	insects	Nymphalidae	<i>Vanessa kershawi</i>	Australian painted lady				2
animals	insects	Nymphalidae	<i>Danaus plexippus</i>	monarch				7
animals	insects	Nymphalidae	<i>Danaus petilia</i>	lesser wanderer				4
animals	insects	Nymphalidae	<i>Euploea corinna</i>	common crow				3
animals	insects	Papilionidae	<i>Graphium choredon</i>	blue triangle				3
animals	insects	Pieridae	<i>Eurema hecabe</i>	large grass-yellow				4
animals	insects	Pieridae	<i>Eurema smilax</i>	small grass-yellow				1
animals	insects	Pieridae	<i>Delias nigrina</i>	black jezebel				2
animals	insects	Pieridae	<i>Catopsilia pomona</i>	lemon migrant				1
animals	insects	Pieridae	<i>Belenois java teutonia</i>	caper white				1
animals	insects	Pieridae	<i>Eurema brigitta australis</i>	no-brand grass-yellow				1
animals	mammals	Acrobatidae	<i>Acrobatodes pygmaeus</i>	feathertail glider		C		1
animals	mammals	Canidae	<i>Canis lupus dingo</i>	dingo				6
animals	mammals	Dasyuridae	<i>Antechinus flavipes flavipes</i>	yellow-footed antechinus (south-east Queensland)		C		5
animals	mammals	Dasyuridae	<i>Dasyurus maculatus maculatus</i>	spotted-tailed quoll (southern subspecies)		V	E	1
animals	mammals	Dasyuridae	<i>Antechinus stuartii</i>	brown antechinus		C		1
animals	mammals	Dasyuridae	<i>Sminthopsis murina</i>	common dunnart		C		2

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	mammals	Dasyuridae	<i>Planigale maculata</i>	common planigale			C	1
animals	mammals	Macropodidae	<i>Macropus rufogriseus</i>	red-necked wallaby			C	19
animals	mammals	Macropodidae	<i>Macropus dorsalis</i>	black-striped wallaby			C	2
animals	mammals	Macropodidae	<i>Macropus giganteus</i>	eastern grey kangaroo			C	20
animals	mammals	Macropodidae	<i>Petrogale penicillata</i>	brush-tailed rock-wallaby		V	V	1
animals	mammals	Macropodidae	<i>Macropus robustus</i>	common wallaroo			C	1
animals	mammals	Macropodidae	<i>Wallabia bicolor</i>	swamp wallaby			C	10/1
animals	mammals	Macropodidae	<i>Macropus parryi</i>	whiptail wallaby			C	4
animals	mammals	Miniopteridae	<i>Miniopterus schreibersii oceanensis</i>	eastern bent-wing bat			C	1
animals	mammals	Molossidae	<i>Tadarida australis</i>	white-striped freetail bat			C	10
animals	mammals	Molossidae	<i>Mormopterus sp.</i>				C	1
animals	mammals	Molossidae	<i>Mormopterus lumsdenae</i>	northern free-tailed bat			C	1
animals	mammals	Muridae	<i>Rattus fuscipes</i>	bush rat			C	2
animals	mammals	Muridae	<i>Rattus tunneyi</i>	pale field-rat			C	2
animals	mammals	Peramelidae	<i>Isodon macrourus</i>	northern brown bandicoot			C	5
animals	mammals	Petauridae	<i>Petaurus australis australis</i>	yellow-bellied glider (southern subspecies)			C	1
animals	mammals	Petauridae	<i>Petaurus norfolcensis</i>	squirrel glider			C	23
animals	mammals	Petauridae	<i>Petaurus breviceps</i>	sugar glider			C	4
animals	mammals	Phalangeridae	<i>Trichosurus vulpecula</i>	common brushtail possum			C	22
animals	mammals	Phascolarctidae	<i>Phascolarctos cinereus</i>	koala		V	V	51
animals	mammals	Potoroidae	<i>Aepyprymnus rufescens</i>	rufous bettong			C	1
animals	mammals	Pseudocheiridae	<i>Pseudocheirus peregrinus</i>	common ringtail possum			C	5
animals	mammals	Pseudocheiridae	<i>Petauroides volans volans</i>	southern greater glider		V	V	15
animals	mammals	Pteropodidae	<i>Pteropus sp.</i>				C	2
animals	mammals	Pteropodidae	<i>Pteropus scapulatus</i>	little red flying-fox			C	9
animals	mammals	Pteropodidae	<i>Pteropus poliocephalus</i>	grey-headed flying-fox			C	8
animals	mammals	Tachyglossidae	<i>Tachyglossus aculeatus</i>	short-beaked echidna			SL	3
animals	mammals	Vespertilionidae	<i>Nyctophilus gouldi</i>	Gould's long-eared bat			C	2
animals	mammals	Vespertilionidae	<i>Scotorepens sp.</i>				C	2
animals	mammals	Vespertilionidae	<i>Scotorepens orion</i>	south-eastern broad-nosed bat			C	3
animals	mammals	Vespertilionidae	<i>Scotorepens greyii</i>	little broad-nosed bat			C	1
animals	reptiles	Agamidae	<i>Pogona barbata</i>	bearded dragon			C	13
animals	reptiles	Agamidae	<i>Diporiphora australis</i>	tommy roundhead			C	5
animals	reptiles	Agamidae	<i>Intellagama lesueurii</i>	eastern water dragon			C	7
animals	reptiles	Boidae	<i>Morelia spilota</i>	carpet python			C	2
animals	reptiles	Colubridae	<i>Boiga irregularis</i>	brown tree snake			C	1
animals	reptiles	Colubridae	<i>Dendrelaphis punctulatus</i>	green tree snake			C	6
animals	reptiles	Colubridae	<i>Tropidonophis mairii</i>	freshwater snake			C	1
animals	reptiles	Diplodactylidae	<i>Diplodactylus vittatus</i>	wood gecko			C	2
animals	reptiles	Diplodactylidae	<i>Oedura tryoni</i>	southern spotted velvet gecko			C	5
animals	reptiles	Diplodactylidae	<i>Nebulifera robusta</i>	robust velvet gecko			C	1
animals	reptiles	Elapidae	<i>Cryptophis nigrescens</i>	eastern small-eyed snake			C	5
animals	reptiles	Elapidae	<i>Pseudonaja textilis</i>	eastern brown snake			C	3
animals	reptiles	Elapidae	<i>Pseudechis porphyriacus</i>	red-bellied black snake			C	6
animals	reptiles	Elapidae	<i>Brachyurophis australis</i>	coral snake			C	2

Kingdom	Class	Family	Scientific Name	Common Name	I	Q	A	Records
animals	reptiles	Elapidae	<i>Vermicella annulata</i>	bandy-bandy		C		1
animals	reptiles	Elapidae	<i>Furina diadema</i>	red-naped snake		C		1
animals	reptiles	Elapidae	<i>Demansia psammophis</i>	yellow-faced whipsnake		C		12
animals	reptiles	Elapidae	<i>Pseudechis guttatus</i>	spotted black snake		C		2
animals	reptiles	Gekkonidae	<i>Gehyra dubia</i>	dubious dtella		C		3
animals	reptiles	Pygopodidae	<i>Lialis burtonis</i>	Burton's legless lizard		C		6
animals	reptiles	Scincidae	<i>Ctenotus spaldingi</i>	straight-browed ctenotus		C		3
animals	reptiles	Scincidae	<i>Tiliqua scincoides</i>	eastern blue-tongued lizard		C		1
animals	reptiles	Scincidae	<i>Lygisaurus foliorum</i>	tree-base litter-skink		C		7
animals	reptiles	Scincidae	<i>Ctenotus taeniolatus</i>	copper-tailed skink		C		2
animals	reptiles	Scincidae	<i>Lampropholis amicula</i>	friendly sunskink		C		2
animals	reptiles	Scincidae	<i>Anomalopus verreauxii</i>	three-clawed worm-skink		C		3
animals	reptiles	Scincidae	<i>Lampropholis delicata</i>	dark-flecked garden sunskink		C		13
animals	reptiles	Scincidae	<i>Morethia taeniopleura</i>	fire-tailed skink		C		1
animals	reptiles	Scincidae	<i>Calyptotis scutirostrum</i>	scute-snouted calyptotis		C		5
animals	reptiles	Scincidae	<i>Ophioscincus ophioscincus</i>	yolk-bellied snake-skink		C		1
animals	reptiles	Scincidae	<i>Carlia pectoralis sensu lato</i>			C		3
animals	reptiles	Scincidae	<i>Cryptoblepharus pulcher pulcher</i>	elegant snake-eyed skink		C		26
animals	reptiles	Scincidae	<i>Carlia munda</i>	shaded-litter rainbow-skink		C		1
animals	reptiles	Scincidae	<i>Carlia schmeltzii</i>	robust rainbow-skink		C		3
animals	reptiles	Scincidae	<i>Concinnia martini</i>	dark bar-sided skink		C		1
animals	reptiles	Scincidae	<i>Carlia pectoralis</i>	open-litter rainbow skink		C		1
animals	reptiles	Scincidae	<i>Ctenotus arcanus</i>	arcane ctenotus		C		1
animals	reptiles	Scincidae	<i>Concinnia tenuis</i>	bar-sided skink		C		1
animals	reptiles	Scincidae	<i>Carlia vivax</i>	tussock rainbow-skink		C		18
animals	reptiles	Varanidae	<i>Varanus varius</i>	lace monitor		C		10

CODES

I - Y indicates that the taxon is introduced to Queensland and has naturalised.

Q - Indicates the Queensland conservation status of each taxon under the *Nature Conservation Act 1992*. The codes are Extinct in the Wild (PE), Endangered (E), Vulnerable (V), Near Threatened (NT), Least Concern (C) or Not Protected ().

A - Indicates the Australian conservation status of each taxon under the *Environment Protection and Biodiversity Conservation Act 1999*. The values of EPBC are Conservation Dependent (CD), Critically Endangered (CE), Endangered (E), Extinct (EX), Extinct in the Wild (XW) and Vulnerable (V).

Records – The first number indicates the total number of records of the taxon for the record option selected (i.e. All, Confirmed or Specimens).

This number is output as 99999 if it equals or exceeds this value. The second number located after the / indicates the number of specimen records for the taxon.

This number is output as 999 if it equals or exceeds this value.

ATTACHMENT 5– Contactor Environmental Awareness Acknowledgement

ENVIRONMENTAL AWARENESS

CONTRACTOR ACKNOWLEDGEMENT

I Tong Hooper, the Contractor (or the Contractor Representative), appointed by Lendlease Communities, acknowledge receipt and acceptance of the Lendlease Communities rules and policies in the **Springfield Rise Site Based Management Plan**. By signing below, I acknowledge that there are mechanisms in place to ensure all material provided within this SBMP will be read and understood by all site contractors and sub-contractors prior to commencing works on site.

Shadforth
Company Name (Please print)

[Signature]
Signature (Contractor / Contractor Representative)

Tong Hooper
Name (Please print)

Project Manager
Title / Position

10/9/19
Date

ATTACHMENT 6– ICC Correspondence

Keira Grundy

From: Christo Louw <Christo.Louw@arcadis.com>
Sent: Wednesday, 5 July 2017 3:18 PM
To: Keira Grundy
Cc: Sam Schroter; Tony Hooper; AA008228; Shane Miley; Jaco le Roux; Daniel O'Malley
Subject: FW: 2.6 Interim Uses App. No. 3136/2017/OW - Clearing and Bulk earthworks

Hi Keira

Please see below advice from ICC that they do not require a pre start meeting to the current clearing extents (the interim 2.6 Approval extents) for Village 7 and Village 11. This requirement can be listed as N/A (waived by ICC) on the pre clearing checklists.

Regards

Christo Louw | Senior Civil Engineer | BENH (Civil) | Christo.Louw@arcadis.com
Arcadis | Level 5/120 Edward Street/ Brisbane 4000
T. + 61 7 3337 0846 | M. 041 828 5709
www.arcadis.com/au



Be green, leave it on the screen.



From: Christo Louw
Sent: 4 July 2017 7:50 AM
To: Shane Miley <Shane.Miley@arcadis.com>; Tom Duffy <Tom.Duffy@lendlease.com>; Joel Salmon <Joel.Salmon@lendlease.com>; Graeme Knox <Graeme.Knox@lendlease.com>; AA008228 <AA008228@arcadis.com>
Subject: Fwd: 2.6 Interim Uses App. No. 3136/2017/OW - Clearing and Bulk earthworks

FYI

Sent from my iPhone

Begin forwarded message:

From: Mark Dillon <Mark.Dillon@ipswich.qld.gov.au>
Date: 4 July 2017 at 7:33:39 am AEST
To: "Christo Louw (Christo.Louw@arcadis.com)" <Christo.Louw@arcadis.com>
Cc: Daniel O'Malley (Daniel.OMalley@arcadis.com) <Daniel.OMalley@arcadis.com>
Subject: 2.6 Interim Uses App. No. 3136/2017/OW - Clearing and Bulk earthworks

Christo,

Reference is made to the receipt by Council of various pre-start documentation for the above project via HIGHTAIL transfer. Given the ongoing nature of the works within the Lend Lease Springfield Rise Development and that various personnel have not changed, it is considered that a formal prestart meeting is not required in this instance.

Regards,



Mark Dillon | Senior Technical Officer (Engineering)
Engineering and Environment Branch
Planning and Development Department

IPSWICH CITY COUNCIL T | 07 3810 7738



City of Ipswich

The City of Ipswich has a bold plan that embraces new ways of living, learning and working as Australia's most liveable and prosperous Smart City.

Confidential Communication | [Email Disclaimer](#)

This email and any files transmitted with it are the property of Arcadis and its affiliates. All rights, including without limitation copyright, are reserved. This email contains information that may be confidential and may also be privileged. It is for the exclusive use of the intended recipient(s). If you are not an intended recipient, please note that any form of distribution, copying or use of this communication or the information in it is strictly prohibited and may be unlawful. If you have received this communication in error, please return it to the sender and then delete the email and destroy any copies of it. While reasonable precautions have been taken to ensure that no software or viruses are present in our emails, we cannot guarantee that this email or any attachment is virus free or has not been intercepted or changed. Any opinions or other information in this email that do not relate to the official business of Arcadis are neither given nor endorsed by it.